

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 September 2004 (30.09.2004)

PCT




(10) International Publication Number
WO 2004/082638 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number:
PCT/US2004/008342
- (22) International Filing Date: 17 March 2004 (17.03.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/455,437 17 March 2003 (17.03.2003) US
60/531,203 19 December 2003 (19.12.2003) US
- (71) Applicant (for all designated States except US): **SYRRX, INC.** [US/US]; 10410 Science Center Drive, San Diego, CA 92121 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BRESSI, Jerome, C.** [US/US]; 3799 Ibis Street, San Diego, CA 92103 (US). **BROWN, Jason, W.** [US/US]; 864 Bangor Street, San Diego, CA 92106 (US). **CAO, Sheldon, X.** [US/US]; 10512 Sand Crab Place, San Diego, CA 92130 (US).

- GANGLOFF, Anthony, R.** [US/US]; 4969 Dick Street, San Diego, CA 92115 (US). **JENNINGS, Andrew, J.** [GB/US]; 605 Bonair Way, La Jolla, 92037 (US). **STAFFORD, Jeffrey, A.** [US/US]; 12752 Sandy Crest Court, San Diego, CA 92130 (US). **VU, Phong, H.** [US/US]; 9310 Galvin Avenue, San Diego, CA 92126 (US). **XIAO, Xiao-Yi** [US/US]; 3851 Torrey Hill Lane, San Diego, CA 92130 (US).
- (74) Agent: **WEITZ, David, J.**; Syrrx, Inc., 10410 Science Center Drive, San Diego, CA 92121 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: HISTONE DEACETYLASE INHIBITORS



BEST AVAILABLE COPY

(57) Abstract: Compounds that may be used to inhibit histone deacetylase having the formula (I) or formula (II) wherein M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound; and Z and Q are as defined herein.

WO 2004/082638 A2



GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

HISTONE DEACETYLASE INHIBITORS

FIELD OF THE INVENTION

[0001] The invention relates to compounds that may be used to inhibit deacetylases and, in one variation, histone deacetylases (HDACs), as well as compositions of matter and kits comprising these compounds. The present invention also relates to methods for inhibiting deacetylases, such as HDAC, as well as treatment methods using compounds according to the present invention.

DESCRIPTION OF RELATED ART

[0002] DNA in eukaryotic cells is tightly complexed with proteins (histones) to form chromatin. Histones are small, positively charged proteins that are rich in basic amino acids (positively charged at physiological pH), which contact the phosphate groups (negatively charged at physiological pH) of DNA. There are five main classes of histones H1, H2A, H2B, H3, and H4. The amino acid sequences of H2A, H2B, H3, and H4 show remarkable conservation between species, wherein H1 varies somewhat and in some cases is replaced by another histone, e.g., H5. Four pairs of each of H2A, H2B, H3 and H4 together form a disk-shaped octomeric protein core, around which DNA (about 140 base pairs) is wound to form a nucleosome. Individual nucleosomes are connected by short stretches of linker DNA associated with another histone molecule to form a structure resembling a beaded string, which is itself arranged in a helical stack, known as a solenoid.

[0003] The majority of histones are synthesized during the S phase of the cell cycle, and newly synthesized histones quickly enter the nucleus to become associated with DNA. Within minutes of its synthesis, new DNA becomes associated with histones in nucleosomal structures.

[0004] A small fraction of histones, more specifically, the amino acid side chains thereof, are enzymatically modified by post-translational addition of methyl, acetyl, or phosphate groups, neutralizing the positive charge of the side chain, or converting it to a negative charge. For example, lysine and arginine groups may be methylated, lysine groups may be acetylated, and serine groups may be phosphorylated. For lysine, the $-(CH_2)_4-NH_2$ sidechain may be acetylated, for example by an acetyltransferase enzyme to give the amide $-(CH_2)_4-$

NHC(=O)CH₃. Methylation, acetylation, and phosphorylation of amino termini of histones that extend from the nucleosomal core affects chromatin structure and gene expression. Spencer and Davie 1999. *Gene* 240:1 1-12.

[0005] Acetylation and deacetylation of histones is associated with transcriptional events leading to cell proliferation and/or differentiation. Regulation of the function of transcriptional factors is also mediated through acetylation. Recent reviews on histone deacetylation include Kouzarides, et al., 1999. *Curr. Opin. Genet. Dev.* 9:1, 40-48 and Pazin, et al. 1997. 89:3 325-328.

[0006] The correlation between acetylation status of histones and the transcription of genes has been known for quite some time. Certain enzymes, specifically acetylases (e.g., histone acetyltransferases (HAT)) and deacetylases (histone deacetylases or HDACs), which regulate the acetylation state of histones have been identified in many organisms and have been implicated in the regulation of numerous genes, confirming a link between acetylation and transcription. In general, histone acetylation is believed to correlate with transcriptional activation, whereas histone deacetylation is believed to be associated with gene repression.





[0007] A growing number of histone deacetylases (HDACs) have been identified. HDACs function as part of large multiprotein complexes, which are tethered to the promoter and repress transcription. Well characterized transcriptional repressors such as MAD, nuclear receptors and YY1 associate with HDAC complexes to exert their repressor function.

[0008] Studies of HDAC inhibitors have shown that these enzymes play an important role in cell proliferation and differentiation. HDACs are believed to be associated with a variety of different disease states including, but not limited to cell proliferative diseases and conditions (Marks, P.A., Richon, V.M., Breslow, R. and Rifkind, R.A., *J. Natl. Cancer Inst. (Bethesda)* 92, 1210-1215, 2000) such as leukemia (Lin et al. 1998. *Nature* 391: 811-814; Grignani, et al. 1998. *Nature* 391: 815-818; Warrell et al. 1998. *J. Natl. Cancer Inst.* 90:1621-1625; Gelmetti et al. 1998. *Mol. Cell Biol.* 18:7185-7191; Wang et al. 1998. *PNAS* 95: 10860-10865), melanomas/squamous cell carcinomas (Gillenwater, et al., 1998, *Int. J. Cancer* 75:217-224; Saunders, et al., 1999, *Cancer Res.* 59:399-404), breast cancer, prostate cancer, bladder cancer (Gelmetti et al. 1998. *Mol. Cell Biol.* 18:7185-7191; Wang et al. 1998. *PNAS* 95: 10860-10865), lung cancer, ovarian cancer and colon cancer (Hassig, et al., 1997, *Chem. Biol.*

4:783-789; Archer, et al., 1998, PNAS, 956791-6796; Swendeman, et al., 1999, Proc. Amer. Assoc. Cancer Res. 40, Abstract #3836).

[0009] Histone deacetylase inhibitors are potent inducers of growth arrest, differentiation, or apoptotic cell death in a variety of transformed cells in culture and in tumor bearing animals (*Histone deacetylase inhibitors as new cancer drugs*, Marks, P.A., Richon, V.M., Breslow, R. and Rifkind, R.A., Current Opinions in Oncology, 2001, Nov. 13 (6): 477-83; *Histone deacetylases and cancer: causes and therapies*, Marks, P., Rifkind, R.A., Richon, V.M., Breslow, R., Miller, T. and Kelly, W.K., Nat. Rev. Cancer 2001 Dec.1 (3):194-202). In addition, HDAC inhibitors are useful in the treatment or prevention of protozoal diseases (US Patent 5,922,837) and psoriasis (PCT Publication No. WO 02/26696).

[0010] A variety of inhibitors of HDAC have been reported. Some of these inhibitors are described in the following table:

Inhibitors	References
Butyrates 	Marks PA, et al., J. Natl. Cancer Inst. 2000, 92:1210-1216; Weidle UH, et al., Anticancer Res. 2000, 20:1471-1486; Gore SD, et al., Exp. Opin. Invest. Drugs 2000, 9:2923-2934; Sowa Y, et al., Biofactors 2000, 12:283-287.
Trichostatin A 	Marks PA, et al., J. Natl. Cancer Inst. 2000, 92:1210-1216; Weidle UH, et al., Anticancer Res. 2000, 20:1471-1486; Nervi C, et al., Cancer Res. 2001, 61:1247-1249; Suzuki T, et al., Int. J. Cancer 2000, 88:992-997.
Suberoylanilidine hydroxamic acid 	Marks PA, et al., J. Natl. Cancer Inst. 2000, 92:1210-1216; Kelly WK, et al., Proc. Amer. Soc. Clin. Oncol. 2001, 20:87a; Butler LM, et al., Cancer Res. 2000, 60:5165-5170.
MS-275 	Lee BI, et al., Cancer Res. 2001, 61:931-934.

[0011] Additional examples of HDAC inhibitors can be found in Marks PA, et al., J. Natl. Cancer Inst. 2000, 92:1210-1216 & Weidle UH, et al., Anticancer Res. 2000, 20:1471-1486 and PCT Publication Nos. WO 02/26696, WO 02/062773, and WO 01/18171.

[0012] Despite the various HDAC inhibitors that have been reported to date, a need continues to exist for new and more effective inhibitors of HDACs.

SUMMARY OF THE INVENTION

[0013] The present invention relates to compounds that have activity for inhibiting HDACs.

[0014] The present invention also provides compositions, articles of manufacture and kits comprising these compounds.

[0015] In one embodiment, a pharmaceutical composition is provided that comprises a HDAC inhibitor according to the present invention as an active ingredient. Pharmaceutical compositions according to the invention may optionally comprise 0.001%-100% of one or more HDAC inhibitors of this invention. These pharmaceutical compositions may be administered or coadministered by a wide variety of routes, including for example, orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery (for example by catheter or stent), subcutaneously, intraadiposally, intraarticularly, or intrathecally. The compositions may also be administered or coadministered in slow release dosage forms.

[0016] The invention is also directed to kits and other articles of manufacture for treating disease states associated with HDAC.

[0017] In one embodiment, a kit is provided that comprises a composition comprising at least one HDAC inhibitor of the present invention in combination with instructions. The instructions may indicate the disease state for which the composition is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition. The kit may also comprise packaging materials. The packaging material may comprise a container for housing the composition. The kit may also optionally comprise additional components, such as syringes for administration of the composition. The kit may comprise the composition in single or multiple dose forms.

[0018] In another embodiment, an article of manufacture is provided that comprises a composition comprising at least one HDAC inhibitor of the present invention in combination with packaging materials. The packaging material may comprise a container for housing the composition. The container may optionally comprise a label indicating the disease state for which the composition is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition. The kit may also optionally comprise additional components, such as syringes for administration of the composition. The kit may comprise the composition in single or multiple dose forms.

[0019] Also provided are methods for preparing compounds, compositions and kits according to the present invention. For example, several synthetic schemes are provided herein for synthesizing compounds according to the present invention.

[0020] Also provided are methods for using compounds, compositions, kits and articles of manufacture according to the present invention.

[0021] In one embodiment, the compounds, compositions, kits and articles of manufacture are used to inhibit HDAC.

[0022] In one embodiment, the compounds, compositions, kits and articles of manufacture are used to treat a disease state for which HDAC possesses activity that contributes to the pathology and/or symptomology of the disease state.

[0023] In another embodiment, a compound is administered to a subject wherein HDAC activity within the subject is altered, preferably reduced.

[0024] In another embodiment, a prodrug of a compound is administered to a subject that is converted to the compound *in vivo* where it inhibits HDAC.

[0025] In another embodiment, a method of inhibiting HDAC is provided that comprises contacting HDAC with a compound according to the present invention.

[0026] In another embodiment, a method of inhibiting HDAC is provided that comprises causing a compound according to the present invention to be present in a subject in order to inhibit HDAC *in vivo*.

[0027] In another embodiment, a method of inhibiting HDAC is provided that comprises administering a first compound to a subject that is converted *in vivo* to a second compound wherein the second compound inhibits HDAC *in vivo*.

[0028] In another embodiment, a therapeutic method is provided that comprises administering a compound according to the present invention.

[0029] In another embodiment, a method of inhibiting cell proliferation is provided that comprises contacting a cell with an effective amount of a compound according to the present invention.

[0030] In another embodiment, a method of inhibiting cell proliferation in a patient is provided that comprises administering to the patient a therapeutically effective amount of a compound according to the present invention.

[0031] In another embodiment, a method of treating a condition in a patient which is known to be mediated by HDAC, or which is known to be treated by HDAC inhibitors, comprising administering to the patient a therapeutically effective amount of a compound according to the present invention.

[0032] In another embodiment, a method is provided for using a compound according to the present invention in order to manufacture a medicament for use in the treatment of disease state which is known to be mediated by HDAC, or which is known to be treated by HDAC inhibitors.

[0033] In another embodiment, a method is provided for treating a disease state for which HDAC possesses activity that contributes to the pathology and/or symptomology of the disease state, the method comprising: causing a compound according to the present invention to be present in a subject in a therapeutically effective amount for the disease state.

[0034] In another embodiment, a method is provided for treating a disease state for which HDAC possesses activity that contributes to the pathology and/or symptomology of the disease state, the method comprising: administering a first compound to a subject that is converted *in vivo* to a second compound such that the second compound is present in the subject in a therapeutically effective amount for the disease state.

[0035] In another embodiment, a method is provided for treating a disease state for which HDAC possesses activity that contributes to the pathology and/or symptomology of the disease state, the method comprising: administering a compound according to the present invention to a subject such that the compound is present in the subject in a therapeutically effective amount for the disease state.

[0036] In another embodiment, a method is provided for treating a cell proliferative disease state comprising treating cells with a compound according to the present invention in combination with an anti-proliferative agent, wherein the cells are treated with the compound according to the present invention before, at the same time, and/or after the cells are treated with the anti-proliferative agent, referred to herein as combination therapy. It is noted that treatment of one agent before another is referred to herein as sequential therapy, even if the agents are also administered together. It is noted that combination therapy is intended to cover when agents are administered before or after each other (sequential therapy) as well as when the agents are administered at the same time.

[0037] Examples of diseases that may be treated by administration of compounds and compositions according to the present invention include, but are not limited to protozoal diseases and cell proliferative diseases and conditions such as leukemia, melanomas, squamous cell carcinomas, breast cancer, prostate cancer, bladder cancer, lung cancer including non small-cell lung cancer and small-cell lung cancer, ovarian cancer, colon cancer, squamous cell carcinoma, astrocytoma, Kaposi's sarcoma, glioblastoma, bladder cancer, head and neck cancer, glioma, colorectal cancer, genitourinary cancer and gastrointestinal cancer.

[0038] It is noted in regard to all of the above embodiments that the present invention is intended to encompass pharmaceutically acceptable salts, biohydrolyzable esters, biohydrolyzable amides, biohydrolyzable carbamates, and solvates (e.g., hydrates) of the compounds, regardless of whether such salts, esters, amides, carbamates and solvates are specified since it is well known in the art to administer pharmaceutical agents in a salt, ester, amide, carbamate or solvated form. It is further noted that prodrugs may also be administered which are altered *in vivo* and become a compound according to the present invention. For example, the compound optionally comprises a substituent that is convertible *in vivo* to a different substituent, such as hydrogen. Accordingly, for example, an inhibitor comprising a hydroxy group may be administered as an ester that is converted by hydrolysis *in vivo* to the hydroxy compound. Suitable esters that may be converted *in vivo* into hydroxy compounds include acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis-b-hydroxynaphthoates, gentisates, isethionates, di-*p*-toluoyltartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, *p*-toluenesulfonates, cyclohexylsulfamates, quinate, esters of amino acids, and the like.

Similarly, an inhibitor comprising an amine group may be administered as an amide that is converted by hydrolysis *in vivo* to the amine compound.

[0039] The various methods of using the compounds of the present invention are intended, regardless of whether prodrug delivery is specified, to encompass the administration of a prodrug that is converted *in vivo* into a compound according to the present invention.

BRIEF DESCRIPTION OF THE FIGURES

[0040] Figure 1 illustrates a ribbon diagram overview of the structure of HDAC8, highlighting the secondary structural elements of the protein.

[0041] Figure 2A illustrates particular examples of substituent R₁ that may be employed in the Z moiety.

[0042] Figure 2B illustrates particular examples of Z moieties that the compounds of the present invention may comprise.

[0043] Figure 2C illustrates examples of moieties, Q, that the leader group may comprise to link the leader group (L) to the remainder of the compound.

[0044] Figure 2D illustrates particular examples of moieties that the leader groups may comprise.

[0045] It is noted in regard to Figures 2A-2D that the squiggle line is intended to indicate a bond to an adjacent moiety. It is also noted that the substituents shown may optionally be further substituted beyond what is shown. Further, one or more heteroatoms may optionally be substituted for the carbon atoms shown. In regard to Figure 2D, it is noted that the leader groups moieties may be incorporated into the leader group in either possible orientation.

[0046] Figure 3 illustrates residues 1-482 of HDAC1 and a 6-histidine tag at the N-terminus (SEQ. I.D. No. 1).

[0047] Figure 4 illustrates the DNA sequence (SEQ. I.D. No. 2) that was used to encode SEQ. I.D. No. 1.

[0048] Figure 5 illustrates residues 1-488 of HDAC2 and a 6-histidine tag at the C-terminus (SEQ. I.D. No. 3).

[0049] Figure 6 illustrates the DNA sequence (SEQ. I.D. No. 4) that was used to encode SEQ. I.D. No. 3.

[0050] Figure 7 illustrates residues 73-845 of HDAC6 and a 6-histidine tag at the C-terminus (SEQ. I.D. No. 5).

[0051] Figure 8 illustrates the DNA sequence (SEQ. I.D. No. 6) that was used to encode SEQ. I.D. No. 5.

[0052] Figure 9 illustrates residues 1-377 of HDAC8 and a 6-histidine tag at the N-terminus (SEQ. I.D. No. 7).

[0053] Figure 10 illustrates the DNA sequence (SEQ. I.D. No. 8) that was used to encode SEQ. I.D. No. 7.

DEFINITIONS

[0054] Unless otherwise stated, the following terms used in the specification and claims shall have the following meanings for the purposes of this Application.

[0055] "Alicyclic" means a moiety comprising a non-aromatic ring structure. Alicyclic moieties may be saturated or partially unsaturated with one or more double or triple bonds. Alicyclic moieties may also optionally comprise heteroatoms such as nitrogen, oxygen and sulfur. Examples of alicyclic moieties include, but are not limited to moieties with C3 - C8 rings such as cyclopropyl, cyclohexane, cyclopentane, cyclopentene, cyclopentadiene, cyclohexene, cyclohexadiene, cycloheptane, cycloheptene, cycloheptadiene, cyclooctane, cyclooctene, and cyclooctadiene.

[0056] "Aliphatic" means a moiety characterized by a straight or branched chain arrangement of constituent carbon atoms and may be saturated or partially unsaturated with one or more double or triple bonds.

[0057] "Alkenyl" represented by itself means a straight or branched, unsaturated, aliphatic radical having a chain of carbon atoms having at least one double bond between adjacent carbon atoms. C_X alkenyl and C_{X-Y} alkenyl are typically used where X and Y indicate the number of carbon atoms in the chain. For example, C₂₋₆ alkenyl includes alkenyls that have a chain of between 2 and 6 carbons.

[0058] "Alkoxy" means an oxygen moiety having a further alkyl substituent.

[0059] "Alkyl" represented by itself means a straight or branched, saturated or unsaturated, aliphatic radical having a chain of carbon atoms, optionally with oxygen (See "oxaalkyl") or nitrogen atoms (See "aminoalkyl") between the carbon atoms. C_X alkyl and

C_{X-Y} alkyl are typically used where X and Y indicate the number of carbon atoms in the chain. For example, C_{1-6} alkyl includes alkyls that have a chain of between 1 and 6 carbons (e.g., methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, isobutyl, *tert*-butyl, vinyl, allyl, 1-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-methylallyl, ethynyl, 1-propynyl, 2-propynyl, and the like). Alkyl represented along with another radical (e.g., as in arylalkyl) means a straight or branched, saturated or unsaturated aliphatic divalent radical having the number of atoms indicated or when no atoms are indicated means a bond (e.g., (C_{6-10}) aryl (C_{0-3}) alkyl includes phenyl, benzyl, phenethyl, 1-phenylethyl 3-phenylpropyl, and the like).

[0060] "Alkylene", unless indicated otherwise, means a straight or branched, saturated or unsaturated, aliphatic, divalent radical. C_X alkylene and C_{X-Y} alkylene are typically used where X and Y indicate the number of carbon atoms in the chain. For example, C_{1-6} alkylene includes methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{CH}_2-$), trimethylene ($-\text{CH}_2\text{CH}_2\text{CH}_2-$), tetramethylene ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$), 2-butenylene ($-\text{CH}_2\text{CH}=\text{CHCH}_2-$), 2-methyltetramethylene ($-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2-$), pentamethylene ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$) and the like).

[0061] "Alkylidene" means a straight or branched unsaturated, aliphatic, divalent radical having a general formula $=\text{CR}_a\text{R}_b$. C_X alkylidene and C_{X-Y} alkylidene are typically used where X and Y indicate the number of carbon atoms in the chain. For example, C_{1-6} alkylidene includes methylenidene ($=\text{CH}_2$), ethylenidene ($=\text{CHCH}_3$), isopropylidene ($=\text{C}(\text{CH}_3)_2$), propylidene ($=\text{CHCH}_2\text{CH}_3$), allylidene ($=\text{CH}-\text{CH}=\text{CH}_2$), and the like).

[0062] "Alkynyl" represented by itself means a straight or branched, unsaturated, aliphatic radical having a chain of carbon atoms having at least one triple bond between adjacent carbon atoms. C_X alkynyl and C_{X-Y} alkynyl are typically used where X and Y indicate the number of carbon atoms in the chain. For example, C_{2-6} alkynyl includes alkynyls that have a chain of between 2 and 6 carbons.

[0063] "Amino" means a nitrogen moiety having two further substituents where each substituent has a hydrogen or carbon atom alpha bonded to the nitrogen. Unless indicated otherwise, the compounds of the invention containing amino moieties may include protected derivatives thereof. Suitable protecting groups for amino moieties include acetyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, and the like.

[0064] "Aminoalkyl" means an alkyl, as defined above, except where one or more substituted or unsubstituted nitrogen atoms (-N-) are positioned between carbon atoms of the alkyl. For example, an (C₂₋₆) aminoalkyl refers to a chain comprising between 2 and 6 carbons and one or more nitrogen atoms positioned between the carbon atoms.

[0065] "Animal" includes humans, non-human mammals (e.g., dogs, cats, rabbits, cattle, horses, sheep, goats, swine, deer, and the like) and non-mammals (e.g., birds, and the like).

[0066] "Aromatic" means a moiety wherein the constituent atoms make up an unsaturated ring system, all atoms in the ring system are *sp*² hybridized and the total number of pi electrons is equal to 4n+2. An aromatic ring may be such that the ring atoms are only carbon atoms or may include carbon and non-carbon atoms (see Heteroaryl).

[0067] "Aryl" means a monocyclic or fused bicyclic ring assembly wherein each ring is aromatic or when fused with a second ring forms an aromatic ring assembly. If one or more ring atoms is not carbon (e.g., N, S), the aryl is a heteroaryl. C_X aryl and C_{X-Y} aryl are typically used where X and Y indicate the number of atoms in the ring.

[0068] "Bicycloalkyl" means a saturated or partially unsaturated fused bicyclic or bridged polycyclic ring assembly.

[0069] "Bicycloaryl" means a bicyclic ring assembly wherein the rings are linked by a single bond or fused and at least one of the rings comprising the assembly is aromatic. C_X bicycloaryl and C_{X-Y} bicycloaryl are typically used where X and Y indicate the number of carbon atoms in the bicyclic ring assembly and directly attached to the ring.

[0070] "Carbamoyl" means the radical -OC(O)NR_aR_b where R_a and R_b are each independently two further substituents where a hydrogen or carbon atom is alpha to the nitrogen. It is noted that carbamoyl moieties may include protected derivatives thereof. Examples of suitable protecting groups for carbamoyl moieties include acetyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, and the like. It is noted that both the unprotected and protected derivatives fall within the scope of the invention.

[0071] "Carbocycle" means a ring consisting of carbon atoms.

[0072] "Carbocyclic ketone derivative" means a carbocyclic derivative having a -C(O)- substituent.

[0073] "Carbonyl" means the radical $-C(O)-$. It is noted that the carbonyl radical may be further substituted with a variety of substituents to form different carbonyl groups including acids, acid halides, amides, esters, and ketones.

[0074] "Carboxy" means the radical $-C(O)O-$. It is noted that compounds of the invention containing carboxy moieties may include protected derivatives thereof, i.e., where the oxygen is substituted with a protecting group. Suitable protecting groups for carboxy moieties include benzyl, *tert*-butyl, and the like.

[0075] "Cyano" means the radical $-CN$.

[0076] "Cycloalkyl" means a non-aromatic, saturated or partially unsaturated, monocyclic, fused bicyclic or bridged polycyclic ring assembly. C_X cycloalkyl and C_{X-Y} cycloalkyl are typically used where X and Y indicate the number of carbon atoms in the ring assembly. For example, C_{3-10} cycloalkyl includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, 2,5-cyclohexadienyl, bicyclo[2.2.2]octyl, adamantan-1-yl, decahydronaphthyl, oxocyclohexyl, dioxocyclohexyl, thiocyclohexyl, 2-oxobicyclo[2.2.1]hept-1-yl, and the like.

[0077] "Cycloalkylene" means a divalent saturated or partially unsaturated, monocyclic ring or bridged polycyclic ring assembly. C_X cycloalkylene and C_{X-Y} cycloalkylene are typically used where X and Y indicate the number of carbon atoms in the ring assembly.

[0078] "Disease" specifically includes any unhealthy condition of an animal or part thereof and includes an unhealthy condition that may be caused by, or incident to, medical or veterinary therapy applied to that animal, i.e., the "side effects" of such therapy.

[0079] "Halo" means fluoro, chloro, bromo or iodo.

[0080] "Halo-substituted alkyl", as an isolated group or part of a larger group, means "alkyl" substituted by one or more "halo" atoms, as such terms are defined in this Application. Halo-substituted alkyl includes haloalkyl, dihaloalkyl, trihaloalkyl, perhaloalkyl and the like (e.g. halo-substituted (C_{1-3}) alkyl includes chloromethyl, dichloromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, perfluoroethyl, 2,2,2-trifluoro-1,1-dichloroethyl, and the like).

[0081] "Heteroatom" refers to an atom that is not a carbon atom. Particular examples of heteroatoms include, but are not limited to nitrogen, oxygen, sulfur and halogens.

[0082] "Heteroatom moiety" includes a moiety where the atom by which the moiety is attached is not a carbon. Examples of heteroatom moieties include -N=, -NR_c-, -N⁺(O⁻)=, -O-, -S- or -S(O)₂-, wherein R_c is further substituent.

[0083] "Heterobicycloalkyl" means bicycloalkyl, as defined in this Application, provided that one or more of the atoms forming the ring is a heteroatom. For example hetero(C₉₋₁₂)bicycloalkyl as used to define Z in this application includes, but is not limited to, 3-aza-bicyclo[4.1.0]hept-3-yl, 2-aza-bicyclo[3.1.0]hex-2-yl, 3-aza-bicyclo[3.1.0]hex-3-yl, and the like.

[0084] "Heterocycloalkylene" means cycloalkylene, as defined in this Application, provided that one or more of the ring member carbon atoms indicated, is replaced by a heteroatom.

[0085] "Heteroaryl" means an aryl ring, as defined in this Application, where one or more of the atoms forming the ring is a heteroatom.

[0086] "Heterobicycloaryl" means bicycloaryl, as defined in this Application, provided that one or more of the atoms forming the ring is a heteroatom. For example, hetero(C₈₋₁₀)bicycloaryl as used in this Application includes, but is not limited to, 2-amino-4-oxo-3,4-dihydropteridin-6-yl, and the like.

[0087] "Heterocycloalkyl" means cycloalkyl, as defined in this Application, provided that one or more of the atoms forming the ring is a heteroatom.

[0088] "Hydroxy" means the radical -OH.

[0089] "Imine derivative" means a derivative comprising the moiety -C(NR)-, wherein R comprises a hydrogen or carbon atom alpha to the nitrogen.

[0090] "Isomers" mean any compound having an identical molecular formulae but differing in the nature or sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and stereoisomers that are nonsuperimposable mirror images are termed "enantiomers" or sometimes "optical isomers". A carbon atom bonded to four nonidentical substituents is termed a "chiral center". A compound with one chiral center has two enantiomeric forms of opposite chirality. A mixture of the two enantiomeric forms is termed a "racemic mixture". A compound that has more than one chiral center has 2ⁿ⁻¹ enantiomeric

pairs, where n is the number of chiral centers. Compounds with more than one chiral center may exist as either an individual diastereomers or as a mixture of diastereomers, termed a "diastereomeric mixture". When one chiral center is present a stereoisomer may be characterized by the absolute configuration of that chiral center. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral center. Enantiomers are characterized by the absolute configuration of their chiral centers and described by the *R*- and *S*-sequencing rules of Cahn, Ingold and Prelog. Conventions for stereochemical nomenclature, methods for the determination of stereochemistry and the separation of stereoisomers are well known in the art (e.g., see "Advanced Organic Chemistry", 4th edition, March, Jerry, John Wiley & Sons, New York, 1992).

[0091] "Nitro" means the radical -NO_2 .

[0092] "Oxaalkyl" means an alkyl, as defined above, except where one or more oxygen atoms (-O-) are positioned between carbon atoms of the alkyl. For example, an (C_{2-6}) oxaalkyl refers to a chain comprising between 2 and 6 carbons and one or more oxygen atoms positioned between the carbon atoms.

[0093] "Oxoalkyl" means an alkyl, further substituted with a carbonyl group. The carbonyl group may be an aldehyde, ketone, ester, amide, acid or acid chloride.

[0094] "Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary use as well as human pharmaceutical use.

[0095] "Pharmaceutically acceptable salts" means salts of inhibitors of the present invention which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity. Such salts include acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or with organic acids such as acetic acid, propionic acid, hexanoic acid, heptanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartatic acid, citric acid, benzoic acid, *o*-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, *p*-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, *p*-toluenesulfonic acid,

camphorsulfonic acid, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid and the like.

[0096] Pharmaceutically acceptable salts also include base addition salts which may be formed when acidic protons present are capable of reacting with inorganic or organic bases. Acceptable inorganic bases include sodium hydroxide, sodium carbonate, potassium hydroxide, aluminum hydroxide and calcium hydroxide. Acceptable organic bases include ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine and the like.

[0097] "Prodrug" means a compound that is convertible *in vivo* metabolically into an inhibitor according to the present invention. The prodrug itself may or may not also have HDAC inhibitory activity. For example, an inhibitor comprising a hydroxy group may be administered as an ester that is converted by hydrolysis *in vivo* to the hydroxy compound. Suitable esters that may be converted *in vivo* into hydroxy compounds include acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis-*b*-hydroxynaphthoates, gentisates, isethionates, di-*p*-toluoyltartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, *p*-toluenesulfonates, cyclohexylsulfamates and quinate.

[0098] "Protected derivatives" means derivatives of inhibitors in which a reactive site or sites are blocked with protecting groups. Protected derivatives are useful in the preparation of inhibitors or in themselves may be active as inhibitors. A comprehensive list of suitable protecting groups can be found in T.W. Greene, *Protecting Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons, Inc. 1999.

[0099] "Substituted or unsubstituted" means that a given moiety may consist of only hydrogen substituents through available valencies (unsubstituted) or may further comprise one or more non-hydrogen substituents through available valencies (substituted) that are not otherwise specified by the name of the given moiety. For example, isopropyl is an example of an ethylene moiety that is substituted by -CH₃. In general, a non-hydrogen substituent may be any substituent that may be bound to an atom of the given moiety that is specified to be substituted. Examples of substituents include, but are not limited to, aldehyde, alicyclic,

aliphatic, alkyl, alkylene, alkylidene, amide, amino, aminoalkyl, aromatic, aryl, bicycloalkyl, bicycloaryl, carbamoyl, carbocycle, carboxy, carbonyl group, cycloalkyl, cycloalkylene, ester, halo, heterobicycloalkyl, heterocycloalkylene, heteroaryl, heterobicycloaryl, heterocycloalkyl, hydroxy, iminoketone, ketone, nitro, oxaalkyl, and oxoalkyl moieties, each of which may optionally also be substituted or unsubstituted.

[0100] "Sulfinyl" means the radical $-S(O)-$. It is noted that the sulfinyl radical may be further substituted with a variety of substituents to form different sulfinyl groups including sulfinic acids, sulfinamides, sulfinyl esters, and sulfoxides.

[0101] "Sulfonyl" means the radical $-S(O)(O)-$. It is noted that the sulfonyl radical may be further substituted with a variety of substituents to form different sulfonyl groups including sulfonic acids, sulfonamides, sulfonate esters, and sulfones.

[0102] "Therapeutically effective amount" means that amount which, when administered to an animal for treating a disease, is sufficient to effect such treatment for the disease.

[0103] "Thiocarbonyl" means the radical $-C(S)-$. It is noted that the thiocarbonyl radical may be further substituted with a variety of substituents to form different thiocarbonyl groups including thioacids, thioamides, thioesters, and thioketones.

[0104] "Treatment" or "treating" means any administration of a compound of the present invention and includes:

(1) preventing the disease from occurring in an animal which may be predisposed to the disease but does not yet experience or display the pathology or symptomatology of the disease,

(2) inhibiting the disease in an animal that is experiencing or displaying the pathology or symptomatology of the disease (i.e., arresting further development of the pathology and/or symptomatology), or

(3) ameliorating the disease in an animal that is experiencing or displaying the pathology or symptomatology of the diseased (i.e., reversing the pathology and/or symptomatology).

[0105] It is noted in regard to all of the definitions provided herein that the definitions should be interpreted as being open ended in the sense that further substituents beyond those specified may be included. Hence, a C_1 alkyl indicates that there is one carbon atom but does not indicate what are the substituents on the carbon atom. Hence, a C_1 alkyl comprises methyl

(i.e., $-\text{CH}_3$) as well as $-\text{CR}_a\text{R}_b\text{R}_c$ where R_a , R_b , and R_c may each independently be hydrogen or any other substituent where the atom alpha to the carbon is a heteroatom or cyano. Hence, CF_3 , CH_2OH and CH_2CN are all C_1 alkyls.

DETAILED DESCRIPTION OF THE INVENTION

[0106] The present invention relates to compounds, compositions, kits and articles of manufacture that may be used to inhibit histone deacetylases (referred to herein as HDACs). The compounds may optionally be more particularly used as inhibitors of Class I HDACs such as HDAC1, HDAC2, HDAC6 and HDAC8.

[0107] At least seventeen human genes that encode proven or putative HDACs have been identified to date, some of which are described in Johnstone, R. W., "Histone-Deacetylase Inhibitors: Novel Drugs for the Treatment of Cancer", Nature Reviews, Volume I, pp. 287-299, (2002) and PCT Publication Nos. 00/10583, 01/18045, 01/42437 and 02/08273.

[0108] HDACs have been categorized into three distinct classes based on their relative size and sequence homology. The different HDACs (Homo sapiens), HDAC classes, sequences and references describing the different HDACs are provided in Tables 1 - 3.

TABLE 1: CLASS I HDACs

HDAC	GenBank Accession Number	Reference
1	NP_004955	Histone deacetylase: a regulator of transcription, Wolffe, A.P., Science 272 (5260), 371-372 (1996)
2	NP_001518	Isolation and mapping of a human gene (RPD3L1) that is homologous to RPD3, a transcription factor in Saccharomyces cerevisiae; Furukawa, Y., Kawakami, T., Sudo, K., Inazawa, J., Matsumine, A., Akiyama, T. and Nakamura, Y., Cytogenet. Cell Genet. 73 (1-2), 130-133 (1996)
3	NP_003874	Isolation and characterization of cDNAs corresponding to an additional member of the human histone deacetylase gene family, Yang, W.M., Yao, Y.L., Sun, J.M., Davie, J.R. and Seto, E., J. Biol. Chem. 272 (44), 28001-28007 (1997)
8	NP_060956	Buggy, J.J., Sideris, M.L., Mak, P., Lorimer, D.D., McIntosh, B. and Clark, J.M. Biochem. J. 350 Pt 1, 199-205 (2000)

HDAC	GenBank Accession Number	Reference
11	NP_079103	Cloning and Functional Characterization of HDAC11, a Novel Member of the Human Histone Deacetylase Family, Gao,L., Cueto,M.A., Asselbergs,F. and Atadja,P., J. Biol. Chem. 277 (28), 25748-25755 (2002)

TABLE 2: CLASS II HDACs

HDAC	GenBank Accession Number	Reference
4	NP_006028	Transcriptional control. Sinful repression, Wolffe,A.P., Nature 387 (6628), 16-17 (1997)
5	NP_631944	Prediction of the coding sequences of unidentified human genes. IX. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro, Nagase,T., Ishikawa,K., Miyajima,N., Tanaka,A., Kotani,H., Nomura,N. and Ohara,O., DNA Res. 5 (1), 31-39 (1998)
6	NP_006035	Transcriptional control. Sinful repression, Wolffe,A.P., Nature 387 (6628), 16-17 (1997)
7	NP_057680	Isolation of a novel histone deacetylase reveals that class I and class II deacetylases promote SMRT-mediated repression, Kao,H.Y., Downes,M., Ordentlich,P. and Evans,R.M., Genes Dev. 14 (1), 55-66 (2000)
9	NP_478056	MEF-2 function is modified by a novel co-repressor, MITR, Sparrow,D.B., Miska,E.A., Langley,E., Reynaud-Deonauth,S., Kotecha,S., Towers,N., Spohr,G., Kouzarides,T. and Mohun,T.J., EMBO J. 18 (18), 5085-5098 (1999)
10	NP_114408	Isolation and characterization of mammalian HDAC10, a novel histone deacetylase, Kao,H.Y., Lee,C.H., Komarov,A., Han,C.C. and Evans,R.M., J. Biol. Chem. 277 (1), 187-193 (2002)

TABLE 3: CLASS III HDACs

HDAC	GenBank Accession Number	Reference
Sirtuin 1	NP_036370	Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity; Frye,R.A.;

HDAC	GenBank Accession Number	Reference
		Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)
Sirtuin 2	NP_085096 / NP_036369	A 'double adaptor' method for improved shotgun library construction; Andersson,B., Wentland,M.A., Ricafrente,J.Y., Liu,W. and Gibbs,R.A.; Anal. Biochem. 236 (1), 107-113 (1996)
Sirtuin 3	NP_036371	Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity; Frye,R.A.; Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)
Sirtuin 4	NP_036372	Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity; Frye,R.A.; Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)
Sirtuin 5	NP_112534 / NP_036373	Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity; Frye,R.A.; Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)
Sirtuin 6	NP_057623	Phylogenetic classification of prokaryotic and eukaryotic Sir2-like proteins; Frye,R.A.; Biochem. Biophys. Res. Commun. 273 (2), 793-798 (2000)
Sirtuin 7	NP_057622	Phylogenetic classification of prokaryotic and eukaryotic Sir2-like proteins; Frye,R.A.; Biochem. Biophys. Res. Commun. 273 (2), 793-798 (2000)

[0109] Of particular note are Class I HDACs. All Class I HDACs appear to be sensitive to inhibition by trichostatin A (TSA). Also of particular note is HDAC8, a protein whose crystal structure Applicants determined and used in conjunction with arriving at the present invention.

[0110] HDAC8 is a 377 residue, 42kDa protein localized to the nucleus of a wide array of tissues, as well as several human tumor cell lines. The wild-type form of full length HDAC8 is described in GenBank Accession Number NP 060956; Buggy,J.J., Sideris,M.L., Mak,P., Lorimer,D.D., McIntosh,B. and Clark,J.M., Cloning and characterization of a novel human

histone deacetylase, HDAC8, Biochem. J. 350 Pt 1, 199-205 (2000). Zn^{2+} is likely native to the protein and required for HDAC8 activity.


1. CRYSTAL STRUCTURE FOR HDAC

[0111] Syrrx, Inc. in San Diego, California recently solved the crystal structure for HDAC8. Knowledge of the crystal structure was used to guide the design of the HDAC inhibitors provided herein.

[0112] Figure 1 illustrates a ribbon diagram overview of the structure of HDAC8, highlighting the secondary structural elements of the protein. HDAC8 was found to have a single domain structure belonging to the open α/β class of folds. The structure consists of a central 8-stranded parallel β -sheet sandwiched between layers of α -helices. The ligand binding clefts lie almost in the plane of the central β -sheet, and are formed primarily by loops emanating from the carboxy-terminal ends of the β -strands comprising the sheet. There are two large structural extensions, which occur beyond the core of the α/β motif, off the second and last β -strands of the central β -sheet. Residues contained in the extension off the second β -strand form a globular "cap" over the core of the protein, play an important role in defining the shape of the ligand binding pockets, and are involved in a number of key interactions with the bound ligands.


2. HDAC INHIBITORS

[0113] In one embodiment, HDAC inhibitors of the present invention are provided that comprise the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen, and


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0114] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;

R₂, R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted; and

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a

metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0115] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;

R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro,


and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound.

[0116] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;


R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound.

[0117] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;

R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl,

aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0118] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_1 is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_1 is not halo, cyano, nitro and thio in the case where the ring atom to which R_1 is bound is nitrogen;

R_{10} and R_{11} are taken together to form a substituted or unsubstituted aromatic ring;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl,


aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0119] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein


R_1 is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_1 is not halo, cyano, nitro and thio in the case where the ring atom to which R_1 is bound is nitrogen;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the remainder of the compound.

[0120] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z-Q- is selected from the group consisting of




R₁ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

R₂, R₃, R₄, and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the remainder of the compound.

[0121] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein


Z-Q- is selected from the group consisting of



R₁ is selected from the group consisting of (C₁₋₄)alkyl, phenyl, 1-piperidin-4-ylmethyl, 2-morpholin-4-yl-ethyl, 2-halo-phenyl, 2-halo-phen(C₁₋₄)alkyl, 3-halo-phen(C₁₋₄)alkyl, 2-CF₃O-phen(C₁₋₄)alkyl, 3-CF₃O-phen(C₁₋₄)alkyl, 3-halo-phenyl, 4-halo-phenyl, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 4-phenoxy-phenyl, 4-benzyloxyphenyl, 4-pyrazol-1-yl-benzyl, 1-p-tolyl-ethyl, pyrrolidin-3-yl, 1-(C₁₋₄)alkyl-pyrrolidin-2-yl, 1-(C₁₋₄)alkyl-pyrrolidin-2-yl; 2-di(C₁₋₄)alkylamino-ethyl, 2-di(C₁₋₄)alkylamino-1-methyl-ethyl, 2-di(C₁₋₄)alkylamino-ethyl, 2-hydroxy-2-phenyl-ethyl, 2-pyridin-2-yl-ethyl, 2-pyridin-3-yl-ethyl, 2-pyridin-4-yl-ethyl, 2-(1H-indol-3-yl)-ethyl, 3-indolyl(C₁₋₄)alkyl, 1-indan-2-yl, *R*-α-(HOCH₂)-phen(C₁₋₄)alkyl, *S*-α-(HOCH₂)-phen(C₁₋₄)alkyl, *S*-β-(HOCH₂)-phen(C₁₋₄)alkyl, *R*-β-(CH₃)-phen(C₁₋₄)alkyl, 6-propylsulfanyl, *trans*-4-hydroxy-cyclohexyl, 1-aza-bicyclo[2.2.2]oct-2-yl, 1-(C₁₋₄)alkyl-piperidin-3-yl, 1-(2,2-difluoro-ethyl)-piperidin-3-yl, (2-di(C₁₋₄)alkylamino-2-phenyl-ethyl), 1-benzyl-piperidin-3-yl, 1-allyl-piperidin-3-yl, 1-acetyl-piperidin-3-yl, piperidin-3-yl, and phen(C₁₋₄)alkyl;


R₂, R₃, R₄, and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, cyano, and nitro;

M is selected from the group consisting of:



and L is E, Z or mixtures of E/Z -CH₂=CH₂-.

[0122] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;


each Y is independently selected from the group consisting of O, S and NR_{12} ;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the ring.

[0123] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;


each X is independently selected from the group consisting of CR_{12} and N;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between M and the ring.

[0124] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where

at least one V is NR₁₂;

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0125] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of C(R₁₂)₂ and NR₁₂ where

at least one V is NR₁₂;

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


[0126] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of





wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_2 , R_3 , R_4 and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0127] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;


R_6 , R_7 , R_8 , and R_9 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

[0128] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;


R_6 , R_7 , R_8 , and R_9 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

[0129] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;


R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

[0130] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_{10} and R_{11} are taken together to form a substituted or unsubstituted aromatic ring;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0131] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the remainder of the compound.

[0132] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z-Q- is selected from the group consisting of



each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;


R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 2-10 atoms separation between M and the Q substituent.

[0133] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z-Q- is selected from the group consisting of




each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12}

where at least one V is NR_{12} ;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, cyano, and nitro;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is selected from the group consisting of



and L is E, Z or mixtures of E/Z $-CH_2=CH_2-$.

[0134] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the ring.

[0135] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the ring.

[0136] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0137] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₂, R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted; and

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy,


heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0138] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl


group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

[0139] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

[0140] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and


thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

[0141] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁₀ and R₁₁ are taken together to form a substituted or unsubstituted aromatic ring;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;


R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

[0142] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or

unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound.

[0143] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z-Q- is selected from the group consisting of



wherein


R₂, R₃, R₄, and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 2-10 atoms separation between the M substituent and the Q substituent.

[0144] In another embodiment, HDAC inhibitors of the present invention are provided that comprise the formula:



wherein

Z-Q- is selected from the group consisting of




wherein

R₂, R₃, R₄, and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, cyano, and nitro;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is selected from the group consisting of




and L is E, Z or mixtures of E/Z -CH₂=CH₂-.

[0145] In one variation of any of the above embodiments comprising Q, Q is a substituted or unsubstituted phenyl ring. In another variation of any of the above embodiments comprising Q, Q is a substituted or unsubstituted heteroaryl. In still another variation of any of the above embodiments comprising Q, Q is a substituted or unsubstituted heteroaryl selected from the group consisting of furan, thiophene, pyrrole, pyrazole, triazole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, benzofuran, isobenzofuran, benzothiophene, isobenzothiophene, indole, isobenzazole, quinoline, isoquinoline, cinnoline, quinazoline, naphthyridine, pyridopyridine, quinoxaline, phthalazine, benthiazole, and triazine.


[0146] In one variation of any of the above embodiments and variations comprising X, at least one X in the six membered ring is a substituted carbon atom. In another variation of any of the above embodiments and variations comprising X, at least one X in the six membered ring is -CF.

[0147] In one variation of any of the above embodiments and variations comprising X, any two adjacent X moieties may optionally be CR₁₂ where the R₁₂ substituents are taken together to form a ring.


[0148] In one variation of any of the above embodiments and variations comprising M, M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols (-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group. In another variation of any of the above embodiments and variations comprising M, M comprises a hydroxamic acid. In yet another variation of any of the above embodiments and variations comprising M, M is selected from the group consisting of:




[0149] In one variation of any of the above embodiments and variations comprising QLM, QLM is




[0150] In another variation of any of the above embodiments and variations comprising QLM, QLM is



[0151] In one variation of any of the above embodiments and variations comprising LM, LM is




[0152] In another variation of any of the above embodiments and variations comprising LM, LM is



[0153] In one variation of any of the above embodiments and variations comprising R₂, R₃, R₄, and R₅, at least one of R₂, R₃, R₄, or R₅ is fluorine.

[0154] In one variation of any of the above embodiments and variations comprising R₆, R₇, R₈, and R₉, at least one of R₆, R₇, R₈, or R₉ is fluorine.

[0155] In one variation of any of the above embodiments and variations comprising R₆, R₇, R₈, and R₉, it is noted that R₆, R₇, R₈ or R₉ may be selected such that the phenyl ring linking the five membered ring and the L group comprise one or two fluorines as indicated in the structural subunit below:



[0156] In one variation of any of the above embodiments and variations comprising R₆, R₇, R₈, and R₉, it is also noted in regard to the R₆, R₇, R₈ and R₉ substituents that any two adjacent substituents may be taken together to form a ring.

[0157] In one variation of any of the above embodiments and variations comprising R₁₀ and R₁₁, it is noted that the substituted or unsubstituted aromatic ring formed when R₁₀ and R₁₁ are taken together may optionally be a substituted or unsubstituted aryl or a heteroaryl.

[0158] In one variation of any of the above embodiments and variations comprising V, each V is selected so that the ring is an unsubstituted or substituted piperidin-3-yl moiety.

[0159] In one variation of any of the above embodiments and variations comprising R₁₄, R₁₄ comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.


[0160] In another variation of any of the above embodiments and variations comprising R₁₄, R₁₄ is a substituted or unsubstituted C₁₋₆ alkyl. In still another variation of any of the above embodiments and variations comprising R₁₄, R₁₄ is a substituted or unsubstituted -C(O)C₁₋₆ alkyl. In a further variation of any of the above embodiments and variations comprising R₁₄, R₁₄ is selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted. In yet another variation of any of the above embodiments and variations comprising R₁₄, R₁₄ is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.

[0161] It is noted in regard to each of the above embodiments that a given alkyl, alkoxy, aryloxy, heteroaryloxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, amino, thio, or carbonyl group substituent may optionally be further substituted. As also noted, such two substituents may be taken together to form a ring. Examples of further substituted alkyl groups include, but are not limited to, those selected from the group consisting of haloalkyl, cycloalkyl, aminoalkyl, oxaalkyl, heteroaralkyl, and aralkyl, each of which may optionally be further substituted. Examples of further substituted alkoxy aryloxy, and heteroaryloxy groups include, but are not limited to, those selected from the group consisting of haloalkoxy, haloaryloxy, and haloheteroaryloxy, each of which may optionally be further substituted. Examples of further substituted aminosulfonyl, alkylsulfonyl, arylsulfonyl, and heteroarylsulfonyl groups include, but are not limited to, those selected from the group consisting of alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, heteroaralkylsulfonyl, and aralkylsulfonyl, each of which may optionally be further substituted. Examples of further substituted amino groups include, but are not limited to, those selected from the group consisting of alkylamino, arylamino, and acylamino, each of which may optionally be further substituted. Examples of further substituted thio groups include, but are not limited to, those selected from the group consisting of alkylthio, arylthio, and heteroarylthio, each of which may optionally be further substituted. Examples of further substituted carbonyl groups include, but are not limited to, acids, acid halides, amides, esters, and ketones. For example, the carbonyl groups may be an alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, aminocarbonyl, alkoxycarbonyl, aralkoxycarbonyl, or heteroaralkoxycarbonyl, each of which may optionally be further substituted.

[0162] It is noted that the preceding lists of examples are not intended to be limiting as other forms of alkyl, alkoxy, aryloxy, heteroaryloxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, amino, and thio groups may also be formed with the addition of other substituents to the base group, some of which are described herein and all of which are intended to fall within the scope of the present invention.

Substituent R₁

[0163] Figure 2A illustrates particular examples of moieties that may be used as a R₁ substituent. The below table also provides examples of different compounds having different R₁ substituents.

**R₁ Substituents**

H	(S)-1-isopropyl-piperidin-3-yl
Me	1-piperidin-4-ylmethyl
Et	phenyl
isopropyl	2-chlorophenyl
(±)-1-methyl-piperidin-3-yl	3-chlorophenyl
(R)-1-methyl-piperidin-3-yl	4-chlorophenyl
(S)-1-methyl-piperidin-3-yl	2-methoxy-phenyl
(±)-1-ethyl-piperidin-3-yl	3-methoxy-phenyl
(R)-1-ethyl-piperidin-3-yl	4-methoxy-phenyl
(S)-1-ethyl-piperidin-3-yl	4-phenoxy-phenyl
(±)-1-isopropyl-piperidin-3-yl	benzyl
(R)-1-isopropyl-piperidin-3-yl	2-chlorobenzyl
Pyrrolidin-3-yl	3-chlorobenzyl
2-trifluoromethoxy-benzyl	4-chlorobenzyl
3-trifluoromethoxy-benzyl	4-pyrazol-1-yl-benzyl
4-trifluoromethoxy-benzyl	(R)-1-phenyl-ethyl
(R)-1-(p-tolyl)-ethyl	(S)-1-phenyl-ethyl
(R)-1-(4-Fluoro-phenyl)-ethyl	2-phenyl-ethyl
(2-fluoro-phenyl)-ethyl	(2-methoxy-phenyl)-ethyl
(3-fluoro-phenyl)-ethyl	(3-methoxy-phenyl)-ethyl
(4-fluoro-phenyl)-ethyl	(4-methoxy-phenyl)-ethyl
(2-chloro-phenyl)-ethyl	(R)-1-(2-phenyl-propyl)
(3-chloro-phenyl)-ethyl	(S)-1-(2-phenyl-propyl)
(4-chloro-phenyl)-ethyl	(R)-1-hydroxymethyl-2-phenylethyl
(±)-2-(4-fluoro-phenyl)-1-methyl-ethyl	(S)-1-hydroxymethyl-2-phenylethyl
(R)-2-hydroxy-2-phenyl-ethyl	2-(1H-indol-3-yl)-ethyl
(S)-2-hydroxy-2-phenyl-ethyl	indan-2-yl
2-pyridin-2-yl-ethyl	3-phenyl-propyl
3-pyridin-2-yl-ethyl	1-(4-Fluoro-phenyl)-ethyl

4-pyridin-2-yl-ethyl	2-piperidin-1-yl-ethyl
4-benzyloxy-phenyl	<i>trans</i> -4-hydroxy-cyclohexyl
tert-butyl	Cyclohexyl
(±)-1-BOC-piperidin-3-yl	2-Diethylamino-ethyl
(±)-piperidin-3-yl	(±)-1-(2-hydroxy-ethyl)-piperidin-3-yl
(R)-piperidin-3-yl	(±)-1-ethyl-pyrrolidin-2-ylmethyl
(S)-piperidin-3-yl	(±)-1-ethyl-pyrrolidin-3-yl
2-dimethylamino-ethyl	1-aza-bicyclo[2.2.2]oct-2-yl
2-dimethylamino-1-methyl-ethyl	1-(2,2-difluoro-ethyl)-piperidin-3-yl
2-diisopropylamino-ethyl	2-dimethylamino-2-phenyl-ethyl
(±)-1-benzyl-piperidin-3-yl	(±)-1-propyl-piperidin-3-yl
(±)-1-allyl-piperidin-3-yl	(±)-1-isobutyl-piperidin-3-yl
(±)-1-acetyl-piperidin-3-yl	

[0164] It should be recognized that the compounds described in the above table where the R₁ substituent is varied may each be further substituted by replacing one or more of the hydrogens implicitly depicted in the structure with non-hydrogen substituents. Such further substituents may optionally form additional fused rings, as is also taught herein.

[0165] In one variation, R₁ is a substituted alkyl where the carbon of R₁ alpha to the ring atom is a tertiary carbon, i.e., in addition to the bond to the ring atom, the carbon atom has two non-hydrogen substituents. It is believed that substitution of the carbon alpha to the ring atom in this manner may reduce oxidation of that alpha carbon, particularly when the ring atom is nitrogen, thus adding to the stability of the compound.

Substituents R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉

[0166] R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ may each independently be selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted.

[0167] It is noted that R₂ and R₃; R₃ and R₄; and R₄ and R₅ may each optionally be taken together to form a ring. The ring formed may optionally be a 5 or 6 membered ring. In one variation, the ring formed is an aryl or heteroaryl ring.

[0168] It is also noted that R₆ and R₇; R₇ and R₈; and R₈ and R₉ may each optionally be taken together to form a ring. The ring formed may optionally be a 5 or 6 membered ring. In one variation, the ring formed is an aryl or heteroaryl ring.

Substituent Z

[0169] Figure 2B illustrates particular examples of Z moieties that the compounds of the present invention may comprise. In one particular embodiment, the Z moiety is a substituted or unsubstituted benzimidazole or imidazole.

[0170] It is noted that the examples of Z moieties shown in Figure 2B may optionally be further substituted as has been specified herein. For example, the various R₁ substituents that may be appended to the ring are not specified in Figure 2B.

[0171] Also, it is noted that Figure 2B is intended only to be exemplary and that other Z substituents may be employed in the compounds according to the present invention consistent with the teachings herein.


Substituent Q

[0172] As noted above, Q may be a substituted or unsubstituted aromatic ring. The substituents of the aromatic ring can vary widely and may optionally be such that one or more additional rings are fused to the core aromatic ring of Q.


[0173] Q may optionally be a 5 or 6 membered aromatic ring. When Q is a 6 membered aromatic ring, moieties Z and L may be meta or para substituents relative to each other on the 6 membered aromatic ring. In one variation, moieties Z and L are meta substituted relative to each other.

[0174] In one variation where Q is a phenyl ring, the phenyl ring may have substituents R₆, R₇, R₈, and R₉. As indicated above, these substituents may each optionally be independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted. It is noted that other substituents may additionally be appended to the phenyl ring without departing from the intended scope of the present invention.

[0175] In another variation, Q is a 5 and 6 membered aromatic ring comprising heteroatoms, i.e., a heteroaryl. For example, the heteroaryl ring may optionally have the formula



or



where a, b, c, d and e are each independently nitrogen (N) or carbon (C), with a proviso that when a and c are both nitrogen, then c is carbon. When a, b, c, d and/or e are carbon, the given carbon atom may be substituted. Examples of substituents include, but are not limited to members selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted. It is noted that other substituents may additionally be appended to the heteroaryl ring without departing from the intended scope of the present invention.

[0176] Examples of rings comprising heteroatoms, including 5 and 6 membered aromatic rings comprising heteroatoms are illustrated in Figure 2C. It is noted that the rings shown in Figure 2C are unsubstituted and that further substitutions may optionally be added as has been specified.

[0177] Further particular examples of rings that may be comprised in the Q substituent include, but are not limited to furan, thiophene, pyrrole, pyrazole, triazole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, benzofuran, isobenzofuran, benzothiophene, isobenzothiophene, indole, isobenzazole, quinoline, isoquinoline, cinnoline, quinoxaline, naphthyridine, pyridopyridine, quinoxaline, phthalazine, benthiazole, and triazine.


[0178] Surprisingly, it was determined that when group Q is a meta substituted aryl or heteroaryl group, the resulting inhibitors show improved biological activities over that of the corresponding para substituted aryl or heteroaryl groups. Preferably, the meta substituted aryl is a meta substituted phenyl moiety that is substituted or unsubstituted. Without being bound by any particular theory, it is believed that the meta substitution serves to direct the zinc

complexing substituent M to a more favorable position so as to allow the zinc complexing substituent to interact with the zinc ion while the remainder of the compound maintains its interaction with hydrophobic regions in the binding pocket of the histone deacetylase.

Metal Ion Complexing Substituent, M

[0179] In regard to each of the above embodiments, substituent M may be a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion, and optionally more particularly a zinc ion since a zinc ion is known to be present in the catalytic site of deacetylases. Hence, the M substituent may facilitate inhibitor binding by complexing with the zinc ion present in the catalytic site of deacetylases. In one particular variation, M is a substituent capable of complexing with a histone deacetylase catalytic site and/or a metal ion.

[0180] Examples of substituents capable of complexing with a zinc ion that may be used as the M substituent include, but are not limited to trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkyloxy group. Particular examples of such substituents include:



[0181] In one particular variation, M is a hydroxamic acid ($-\text{C}(\text{O})-\text{NHOH}$), also shown above. It is noted that hydroxamic acids, such as trichostatin A, have been shown to be effective inhibitors against histone deacetylases by complexing with the zinc ion present in the catalytic site of histone deacetylases.

Leader group, L

[0182] In regard to each of the above embodiments, the leader group, L, may be any substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound. The number of atoms separating the M substituent and the remainder of the compound serves to extend the zinc complexing substituent, M, a sufficient

distance away from the remainder of the compound so as to allow the zinc complexing substituent to interact with the zinc ion while the remainder of the compound interacts with hydrophobic regions in the binding pocket of the deacetylase.

[0183] In one embodiment, the leader group, L, provides between 1-10 atoms that extend from the M substituent to remainder of the compound, optionally 3-9 and optionally 4-8 atoms. In one variation, the number of atoms separating the M substituent from the remainder of the compound is 3, 4, 5, 6, 7, 8 or 9 atoms.

[0184] It is noted that the atoms of the leader group extending between the M substituent and the remainder of the compound may consist only of carbon atoms. Alternatively, the atoms of the leader group extending between the M substituent and the remainder of the compound may also comprise non-carbon atoms such as nitrogen, oxygen and sulfur.

[0185] It is also noted that the bonds between the atoms of the leader group extending between the M substituent and the remainder of the compound may be saturated, partially unsaturated, or fully unsaturated. For example, the leader group may comprise one or more alkene ($-\text{CH}=\text{CH}-$) or alkyne ($-\text{C}\equiv\text{C}-$) bonds.

[0186] A variety of different moieties may be incorporated into the leader groups of the HDAC inhibitors of the present invention. Examples of such moieties are shown in Figure 2D.

[0187] The atoms forming the backbone of the leader group, L, may optionally comprise one or more members of the group consisting of: $-(\text{CH}_2)_n-$, where n is an integer from 1 to 10; $-\text{CH}(\text{CH}_3)-$; $-\text{CH}(\text{CH}_3)\text{CH}_2-$ and $-\text{CH}_2\text{CH}(\text{CH}_3)-$; $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)-$; $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)-$; $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CHCH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)-$; $-\text{CH}(\text{CH}_2\text{CH}_3)-$; $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2-$ and $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)-$; $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)-$; $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)-$; $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2-$, and $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$; $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CHCH}(\text{CH}_2\text{CH}_3)$; $-\text{CH}=\text{CH}-$; $-\text{CH}=\text{CHCH}_2-$ and $-\text{CH}_2\text{CH}=\text{CH}-$; $-\text{CH}=\text{CHCHCH}_2-$, $-\text{CH}_2\text{CH}=\text{CHCH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}-$; -

CH=CHCH₂CH₂CH₂-, -CH₂CH=CHCH₂CH₂-, -CH₂CH₂CH=CHCH₂-, and -H₂CH₂CH₂CH=CH-; -CH=CHCHCH₂CH₂CH₂-, -CH₂CH=CHCH₂CH₂CH₂-, -CH₂CH₂CH=CHCH₂CH₂-, -CH₂CH₂CH₂CH=CHCH₂-, and -CH₂CH₂CH₂CHCH=CH-; -C(CH₃)=CH- and -CH=C(CH₃)-; -C(CH₃)=CHCH₂-, -CH=C(CH₃)CH₂-, and -CH=CHCH(CH₃)-; -CH(CH₃)CH=CH-, -CH₂C(CH₃)=CH-, and -CH₂CH=C(CH₃)-; -CH=CHCH=CH-; -CH=CHCH=CHCH₂-, -CH₂CH=CHCH=CH-, and -CH=CHCH₂CH=CH-; -CH=CHCH=CHCH₂CH₂-, -CH=CHCH₂CH=CHCH₂-, and -CH=CHCH₂CH₂CH=CH-, -CH₂CH=CHCH=CHCH₂-, -CH₂CH=CHCH₂CH=CH-, and -CH₂CH₂CH=CHCH=CH-; -C(CH₃)=CHCH=CH-, -CH=C(CH₃)CH=CH-, -CH=CHC(CH₃)=CH-, and -CH=CHCH=C(CH₃)-; -C≡C-, -C≡CCH₂-, -CH₂C≡C-, -C≡CCH(CH₃)-, and -CH(CH₃)C≡C-; -C≡CCH₂CH₂-, -CH₂C≡CCH₂-, and -CH₂CH₂C≡C-; -C≡CCH(CH₃)CH₂- and -C≡CCH₂CH(CH₃)-; -CH(CH₃)C≡CCH₂- and -CH₂C≡CCH(CH₃)-; -CH(CH₃)CH₂C≡C- and -CH₂CH(CH₃)C≡C-; -C≡CCH=CH-, -CH=CHC≡C-, and -C≡CC≡C-; -C≡CCH₂CH₂CH₂- and -CH₂CH₂CH₂C≡C-; -C≡CCH₂CH₂CH₂CH₂- and -CH₂CH₂CH₂CH₂C≡C-; -C≡CCH=CHCH=CH-, -CH=CHC≡C-CH=CH-, and -CH=CHCH=CHC≡C-; -C(CH₃)=CHC≡C-, -CH=C(CH₃)C≡C-, -C≡CC(CH₃)=CH-, and -C≡CCH=C(CH₃). L may also be E, Z or mixtures of E/Z -CH₂=CH₂-. It is noted that the hydrogen atoms of above possible portions of the leader group may optionally be substituted with further substituents.

[0188] It is also noted that the leader group may comprise one or more substituents extending from one or more atoms of the leader group backbone. In one variation, two substituents extending from the atoms extending between the carbon alpha to the leader group and the M substituent to form one or more three, four, five, six, seven, eight or nine membered rings. The atoms of the leader group forming the ring may be separated from each other by 0, 1, 2, 3, or 4 atoms.

[0189] The rings may be saturated or partially unsaturated (i.e., comprise one or two double bonds). The rings may also be aromatic, referred to herein as aryl and heteroaryl rings. The rings may optionally be further substituted. These further ring substituents may combine to form additional rings that are fused to the rings forming a portion of the backbone, e.g., bicycloaryl and bicycloheteroaryl.

[0190] Examples of cycloalkyl rings that may be formed by one or more leader group backbone atoms include, but are not limited to: cyclopropyl, cyclohexane, cyclopentane,

cyclopentene, cyclopentadiene, cyclohexane, cyclohexene, cyclohexadiene, phenyl, cycloheptane, cycloheptene, cycloheptadiene, cyclooctane, cyclooctene, and cyclooctadiene.

[0191] Examples of heteroaryl rings that may be formed by one or more leader group backbone atoms include, but are not limited to: furan, thiofuran, pyrrole, isopyrrole, 3-isopyrrole, pyrazole, isoimidazole, triazole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, benzofuran, isobenzofuran, benzothiofuran, isobenzothiophene, indole, isobenzazole, quinoline, isoquinoline, cinnoline, quinazoline, naphthyridine, and pyridopyridine.

[0192] It is noted that the inhibitors may include one or more chiral centers. The chiral centers may be either the R or S enantiomers, depending on the substituents.

[0193] Synthetic scheme for synthesizing compounds according to these various embodiments are provided in the Examples. Particular examples of HDAC inhibitors according to these embodiments are provided in the examples.

A. Salts, Hydrates, and Prodrugs of HDAC Inhibitors

[0194] It should be recognized that the compounds of the present invention may be present and optionally administered in the form of salts, hydrates and prodrugs that are converted *in vivo* into the compounds of the present invention. For example, it is within the scope of the present invention to convert the compounds of the present invention into and use them in the form of their pharmaceutically acceptable salts derived from various organic and inorganic acids and bases in accordance with procedures well known in the art.

[0195] When the compounds of the present invention possess a free base form, the compounds can be prepared as a pharmaceutically acceptable acid addition salt by reacting the free base form of the compound with a pharmaceutically acceptable inorganic or organic acid, e.g., hydrohalides such as hydrochloride, hydrobromide, hydroiodide; other mineral acids and their corresponding salts such as sulfate, nitrate, phosphate, etc.; and alkyl- and monoarylsulfonates such as ethanesulfonate, toluenesulfonate and benzenesulfonate; and other organic acids and their corresponding salts such as acetate, tartrate, maleate, succinate, citrate, benzoate, salicylate and ascorbate. Further acid addition salts of the present invention include, but are not limited to: adipate, alginate, arginate, aspartate, benzenesulfonate (besylate), bisulfate, bisulfite, bromide, butyrate, camphorate, camphorsulfonate, caprylate, chloride, chlorobenzoate, cyclopentanepropionate, digluconate, dihydrogenphosphate, dinitrobenzoate,

dodecylsulfate, ethanesulfonate, fumarate, galacterate (from mucic acid), galacturonate, glucoheptaoate, gluconate, glutamate, glycerophosphate, hemisuccinate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, iodide, isethionate, iso-butyrate, lactate, lactobionate, malate, malonate, mandelate, metaphosphate, methanesulfonate, methylbenzoate, monohydrogenphosphate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, oleate, pamoate, pectinate, persulfate, phenylacetate, 3-phenylpropionate, phosphate, phosphonate and phthalate. It should be recognized that the free acid forms will typically differ from their respective salt forms somewhat in physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid forms for the purposes of the present invention.

[0196] When the compounds of the present invention possess a free base form, a pharmaceutically acceptable base addition salt can be prepared by reacting the free acid form of the compound with a pharmaceutically acceptable inorganic or organic base. Examples of such bases are alkali metal hydroxides including potassium, sodium and lithium hydroxides; alkaline earth metal hydroxides such as barium and calcium hydroxides; alkali metal alkoxides, e.g. potassium ethanolate and sodium propanolate; and various organic bases such as ammonium hydroxide, piperidine, diethanolamine and N-methylglutamine. Also included are the aluminum salts of the compounds of the present invention. Further base salts of the present invention include, but are not limited to: copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium and zinc salts. Organic base salts include, but are not limited to, salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, e.g., arginine, betaine, caffeine, chlorprocaine, choline, N, N'-dibenzylethylenediamine (benzathine), dicyclohexylamine, diethanolamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, iso-propylamine, lidocaine, lysine, meglumine, N-methyl-D-glucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethanolamine, triethylamine, trimethylamine, tripropylamine and tris-(hydroxymethyl)-methylamine (tromethamine). It should be recognized that the free base forms will typically differ from their respective salt forms

somewhat in physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free base forms for the purposes of the present invention.

[0197] Compounds of the present invention, which comprise basic nitrogen-containing groups, may be quaternized with such agents as (C₁₋₄) alkyl halides, e.g., methyl, ethyl, isopropyl and tert-butyl chlorides, bromides and iodides; di (C₁₋₄) alkyl sulfates, e.g., dimethyl, diethyl and diamyl sulfates; (C₁₀₋₁₈) alkyl halides, e.g., decyl, dodecyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; and aryl (C₁₋₄) alkyl halides, e.g., benzyl chloride and phenethyl bromide. Such salts permit the preparation of both water-soluble and oil-soluble compounds of the present invention.

[0198] *N*-oxides of compounds according to the present invention can be prepared by methods known to those of ordinary skill in the art. For example, *N*-oxides can be prepared by treating an unoxidized form of the compound with an oxidizing agent (e.g., trifluoroperacetic acid, permaleic acid, perbenzoic acid, peracetic acid, *meta*-chloroperoxybenzoic acid, or the like) in a suitable inert organic solvent (e.g., a halogenated hydrocarbon such as dichloromethane) at approximately 0 °C. Alternatively, the *N*-oxides of the compounds can be prepared from the *N*-oxide of an appropriate starting material.

[0199] Prodrug derivatives of compounds according to the present invention can be prepared by modifying substituents of compounds of the present invention that are then converted *in vivo* to a different substituent. It is noted that in many instances, the prodrugs themselves also fall within the scope of the range of compounds according to the present invention. For example, prodrugs can be prepared by reacting a compound with a carbamylating agent (e.g., 1,1-acyloxyalkylcarbonochloridate, *para*-nitrophenyl carbonate, or the like) or an acylating agent. Further examples of methods of making prodrugs are described in Saulnier *et al.* (1994), *Bioorganic and Medicinal Chemistry Letters*, Vol. 4, p. 1985.

[0200] Protected derivatives of compounds of the present invention can also be made. Examples of techniques applicable to the creation of protecting groups and their removal can be found in T.W. Greene, *Protecting Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons, Inc. 1999.

[0201] Compounds of the present invention may also be conveniently prepared, or formed during the process of the invention, as solvates (e.g. hydrates). Hydrates of compounds of the present invention may be conveniently prepared by recrystallization from an aqueous/organic solvent mixture, using organic solvents such as dioxane, tetrahydrofuran or methanol.

[0202] A "pharmaceutically acceptable salt", as used herein, is intended to encompass any compound according to the present invention that is utilized in the form of a salt thereof, especially where the salt confers on the compound improved pharmacokinetic properties as compared to the free form of compound or a different salt form of the compound. The pharmaceutically acceptable salt form may also initially confer desirable pharmacokinetic properties on the compound that it did not previously possess, and may even positively affect the pharmacodynamics of the compound with respect to its therapeutic activity in the body. An example of a pharmacokinetic property that may be favorably affected is the manner in which the compound is transported across cell membranes, which in turn may directly and positively affect the absorption, distribution, biotransformation and excretion of the compound. While the route of administration of the pharmaceutical composition is important, and various anatomical, physiological and pathological factors can critically affect bioavailability, the solubility of the compound is usually dependent upon the character of the particular salt form thereof, which it utilized. One of skill in the art will appreciate that an aqueous solution of the compound will provide the most rapid absorption of the compound into the body of a subject being treated, while lipid solutions and suspensions, as well as solid dosage forms, will result in less rapid adsorption of the compound.

3. PREPARATION OF HDAC INHIBITORS

[0203] Various methods may be developed for synthesizing compounds according to the present invention. Representative methods for synthesizing these compounds are provided in the Examples. It is noted, however, that the compounds of the present invention may also be synthesized by other synthetic routes that others may devise.

[0204] It will be readily recognized that certain compounds according to the present invention have atoms with linkages to other atoms that confer a particular stereochemistry to the compound (e.g., chiral centers). It is recognized that synthesis of compounds according to the present invention may result in the creation of mixtures of different stereoisomers

(enantiomers, diastereomers). Unless a particular stereochemistry is specified, recitation of a compound is intended to encompass all of the different possible stereoisomers.

[0205] Various methods for separating mixtures of different stereoisomers are known in the art. For example, a racemic mixture of a compound may be reacted with an optically active resolving agent to form a pair of diastereoisomeric compounds. The diastereomers may then be separated in order to recover the optically pure enantiomers. Dissociable complexes may also be used to resolve enantiomers (e.g., crystalline diastereoisomeric salts). Diastereomers typically have sufficiently distinct physical properties (e.g., melting points, boiling points, solubilities, reactivity, etc.) that they can be readily separated by taking advantage of these dissimilarities. For example, diastereomers can typically be separated by chromatography or by separation/resolution techniques based upon differences in solubility. A more detailed description of techniques that can be used to resolve stereoisomers of compounds from their racemic mixture can be found in Jean Jacques Andre Collet, Samuel H. Wilen, *Enantiomers, Racemates and Resolutions*, John Wiley & Sons, Inc. (1981).

4. INDICATIONS FOR USE OF HDAC INHIBITORS

[0206] HDAC is believed to contribute to the pathology and/or symptomology of several different diseases such that reduction of the activity of HDAC in a subject through inhibition may be used to therapeutically address these disease states. Examples of various diseases that may be treated using the HDAC inhibitors of the present invention are described herein. It is noted that additional diseases beyond those disclosed herein may be later identified as the biological roles that HDAC play in various pathways becomes more fully understood.

A. Undesirable or Uncontrolled Cell Proliferation

[0207] One set of indications that HDAC inhibitors of the present invention may be used to treat are those involving undesirable or uncontrolled cell proliferation. Such indications include benign tumors, various types of cancers such as primary tumors and tumor metastasis, restenosis (e.g. coronary, carotid, and cerebral lesions), abnormal stimulation of endothelial cells (atherosclerosis), insults to body tissue due to surgery, abnormal wound healing, abnormal angiogenesis, diseases that produce fibrosis of tissue, repetitive motion disorders, disorders of tissues that are not highly vascularized, and proliferative responses associated with organ transplants. More specific indications for HDAC inhibitors include, but are not

limited to prostate cancer, lung cancer, acute leukemia, multiple myeloma, bladder carcinoma, renal carcinoma, breast carcinoma, colorectal carcinoma, neuroblastoma and melanoma.

[0208] In one embodiment, a method is provided for treating diseases associated with undesired and uncontrolled cell proliferation. The method comprises administering to a subject suffering from uncontrolled cell proliferation a therapeutically effective amount of a HDAC inhibitor according to the present invention, such that said uncontrolled cell proliferation is reduced. The particular dosage of the inhibitor to be used will depend on the severity of the disease state, the route of administration, and related factors that can be determined by the attending physician. Generally, acceptable and effective daily doses are amounts sufficient to effectively slow or eliminate uncontrolled cell proliferation.

[0209] HDAC inhibitors according to the present invention may also be used in conjunction with other agents to inhibit undesirable and uncontrolled cell proliferation. Examples of other anti-cell proliferation agents that may be used in conjunction with the HDAC inhibitors of the present invention include, but are not limited to, retinoid acid and derivatives thereof, 2-methoxyestradiol, ANGIOSTATIN™ protein, ENDOSTATIN™ protein, suramin, squalamine, tissue inhibitor of metalloproteinase-I, tissue inhibitor of metalloproteinase-2, plasminogen activator inhibitor-1, plasminogen activator inhibitor-2, cartilage-derived inhibitor, paclitaxel, platelet factor 4, protamine sulfate (clupeine), sulfated chitin derivatives (prepared from queen crab shells), sulfated polysaccharide peptidoglycan complex (sp-pg), staurosporine, modulators of matrix metabolism, including for example, proline analogs ((1-azetidine-2-carboxylic acid (LACA), cishydroxyproline, d,l-3,4-dehydroproline, thiaproline), beta.-aminopropionitrile fumarate, 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; methotrexate, mitoxantrone, heparin, interferons, 2 macroglobulin-serum, chimp-3, chymostatin, beta.-cyclodextrin tetradecasulfate, eponemycin; fumagillin, gold sodium thiomalate, d-penicillamine (CDPT), beta.-1-anticollagenase-serum, alpha.2-antiplasmin, bisantrene, lobenzarit disodium, n-(2-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA", thalidomide; angostatic steroid, carboxyaminoimidazole; metalloproteinase inhibitors such as BB94. Other anti-angiogenesis agents that may be used include antibodies, preferably monoclonal antibodies against these angiogenic growth factors: bFGF, aFGF, FGF-5, VEGF isoforms, VEGF-C, HGF/SF and Ang-1/Ang-2. Ferrara N. and

Alitalo, K. "Clinical application of angiogenic growth factors and their inhibitors" (1999) Nature Medicine 5:1359-1364.

[0210] Generally, cells in benign tumors retain their differentiated features and do not divide in a completely uncontrolled manner. A benign tumor is usually localized and nonmetastatic. Specific types of benign tumors that can be treated using HDAC inhibitors of the present invention include hemangiomas, hepatocellular adenoma, cavernous haemangioma, focal nodular hyperplasia, acoustic neuromas, neurofibroma, bile duct adenoma, bile duct cystanoma, fibroma, lipomas, leiomyomas, mesotheliomas, teratomas, myxomas, nodular regenerative hyperplasia, trachomas and pyogenic granulomas.

[0211] In the case of malignant tumors, cells become undifferentiated, do not respond to the body's growth control signals, and multiply in an uncontrolled manner. Malignant tumors are invasive and capable of spreading to distant sites (metastasizing). Malignant tumors are generally divided into two categories: primary and secondary. Primary tumors arise directly from the tissue in which they are found. Secondary tumors, or metastases, are tumors that originated elsewhere in the body but have now spread to distant organs. Common routes for metastasis are direct growth into adjacent structures, spread through the vascular or lymphatic systems, and tracking along tissue planes and body spaces (peritoneal fluid, cerebrospinal fluid, etc.).

[0212] Specific types of cancers or malignant tumors, either primary or secondary, that can be treated using the HDAC inhibitors of the present invention include, but are not limited to, leukemia, breast cancer, skin cancer, bone cancer, prostate cancer, liver cancer, lung cancer, brain cancer, cancer of the larynx, gallbladder, pancreas, rectum, parathyroid, thyroid, adrenal, neural tissue, head and neck, colon, stomach, bronchi, kidneys, basal cell carcinoma, squamous cell carcinoma of both ulcerating and papillary type, metastatic skin carcinoma, osteo sarcoma, Ewing's sarcoma, veticulum cell sarcoma, myeloma, giant cell tumor, small-cell lung tumor, gallstones, islet cell tumor, primary brain tumor, acute and chronic lymphocytic and granulocytic tumors, hairy-cell tumor, adenoma, hyperplasia, medullary carcinoma, pheochromocytoma, mucosal neuronms, intestinal ganglioneuromas, hyperplastic corneal nerve tumor, marfanoid habitus tumor, Wilm's tumor, seminoma, ovarian tumor, leiomyomater tumor, cervical dysplasia and in situ carcinoma, neuroblastoma, retinoblastoma, soft tissue sarcoma, malignant carcinoid, topical skin lesion, mycosis fungoide,

rhabdomyosarcoma, Kaposi's sarcoma, osteogenic and other sarcoma, malignant hypercalcemia, renal cell tumor, polycythemia vera, adenocarcinoma, glioblastoma multiforma, leukemias, lymphomas, malignant melanomas, epidermoid carcinomas, and other carcinomas and sarcomas.

[0213] The HDAC inhibitors of the present invention may also be used to treat abnormal cell proliferation due to insults to body tissue during surgery. These insults may arise as a result of a variety of surgical procedures such as joint surgery, bowel surgery, and cheloid scarring. Diseases that produce fibrotic tissue include emphysema. Repetitive motion disorders that may be treated using the present invention include carpal tunnel syndrome. An example of a cell proliferative disorder that may be treated using the invention is a bone tumor.

[0214] Proliferative responses associated with organ transplantation that may be treated using HDAC inhibitors of the invention include proliferative responses contributing to potential organ rejections or associated complications. Specifically, these proliferative responses may occur during transplantation of the heart, lung, liver, kidney, and other body organs or organ systems.

[0215] Abnormal angiogenesis that may be may be treated using this invention include those abnormal angiogenesis accompanying rheumatoid arthritis, ischemic-reperfusion related brain edema and injury, cortical ischemia, ovarian hyperplasia and hypervascularity, (polycystic ovary syndrome), endometriosis, psoriasis, diabetic retinopathy, and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neurovascular glaucoma and Oster Webber syndrome.

[0216] Examples of diseases associated with uncontrolled angiogenesis that may be treated according to the present invention include, but are not limited to retinal/choroidal neovascularization and corneal neovascularization. Examples of retinal/choroidal neovascularization include, but are not limited to, Best's disease, myopia, optic pits, Stargardt's disease, Paget's disease, vein occlusion, artery occlusion, sickle cell anemia, sarcoid, syphilis, pseudoxanthoma elasticum carotid artery obstructive diseases, chronic uveitis/vitritis, mycobacterial infections, Lyme's disease, systemic lupus erythematosus, retinopathy of prematurity, Eales disease, diabetic retinopathy, macular degeneration, Behcet's disease, infections causing a retinitis or chorioiditis, presumed ocular histoplasmosis, pars planitis,

chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, trauma and post-laser complications, diseases associated with rubesis (neovascularization of the angle) and diseases caused by the abnormal proliferation of fibrovascular or fibrous tissue including all forms of proliferative vitreoretinopathy. Examples of corneal neovascularization include, but are not limited to, epidemic keratoconjunctivitis, Vitamin A deficiency, contact lens overwear, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, Mooren ulcer, Terrien's marginal degeneration, marginal keratolysis, polyarteritis, Wegener sarcoidosis, Scleritis, periphigoid radial keratotomy, neovascular glaucoma and retrolental fibroplasia, syphilis, Mycobacteria infections, lipid degeneration, chemical burns, bacterial ulcers, fungal ulcers, Herpes simplex infections, Herpes zoster infections, protozoan infections and Kaposi sarcoma.

[0217] Chronic inflammatory diseases associated with uncontrolled angiogenesis may also be treated using HDAC inhibitors of the present invention. Chronic inflammation depends on continuous formation of capillary sprouts to maintain an influx of inflammatory cells. The influx and presence of the inflammatory cells produce granulomas and thus maintains the chronic inflammatory state. Inhibition of angiogenesis using a HDAC inhibitor alone or in conjunction with other anti-inflammatory agents may prevent the formation of the granulomas and thus alleviate the disease. Examples of chronic inflammatory diseases include, but are not limited to, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, psoriasis, sarcoidosis, and rheumatoid arthritis.

[0218] Inflammatory bowel diseases such as Crohn's disease and ulcerative colitis are characterized by chronic inflammation and angiogenesis at various sites in the gastrointestinal tract. For example, Crohn's disease occurs as a chronic transmural inflammatory disease that most commonly affects the distal ileum and colon but may also occur in any part of the gastrointestinal tract from the mouth to the anus and perianal area. Patients with Crohn's disease generally have chronic diarrhea associated with abdominal pain, fever, anorexia, weight loss and abdominal swelling. Ulcerative colitis is also a chronic, nonspecific, inflammatory and ulcerative disease arising in the colonic mucosa and is characterized by the presence of bloody diarrhea. These inflammatory bowel diseases are generally caused by chronic granulomatous inflammation throughout the gastrointestinal tract, involving new

capillary sprouts surrounded by a cylinder of inflammatory cells. Inhibition of angiogenesis by these inhibitors should inhibit the formation of the sprouts and prevent the formation of granulomas. Inflammatory bowel diseases also exhibit extra intestinal manifestations, such as skin lesions. Such lesions are characterized by inflammation and angiogenesis and can occur at many sites other the gastrointestinal tract. Inhibition of angiogenesis by HDAC inhibitors according to the present invention can reduce the influx of inflammatory cells and prevent lesion formation.

[0219] Sarcoidosis, another chronic inflammatory disease, is characterized as a multisystem granulomatous disorder. The granulomas of this disease can form anywhere in the body. Thus, the symptoms depend on the site of the granulomas and whether the disease is active. The granulomas are created by the angiogenic capillary sprouts providing a constant supply of inflammatory cells. By using HDAC inhibitors according to the present invention to inhibit angiogenesis, such granulomas formation can be inhibited. Psoriasis, also a chronic and recurrent inflammatory disease, is characterized by papules and plaques of various sizes. Treatment using these inhibitors alone or in conjunction with other anti-inflammatory agents should prevent the formation of new blood vessels necessary to maintain the characteristic lesions and provide the patient relief from the symptoms.

[0220] Rheumatoid arthritis (RA) is also a chronic inflammatory disease characterized by non-specific inflammation of the peripheral joints. It is believed that the blood vessels in the synovial lining of the joints undergo angiogenesis. In addition to forming new vascular networks, the endothelial cells release factors and reactive oxygen species that lead to pannus growth and cartilage destruction. The factors involved in angiogenesis may actively contribute to, and help maintain, the chronically inflamed state of rheumatoid arthritis. Treatment using HDAC inhibitors according to the present invention alone or in conjunction with other anti-RA agents may prevent the formation of new blood vessels necessary to maintain the chronic inflammation and provide the RA patient relief from the symptoms.

5. COMPOSITIONS COMPRISING HDAC INHIBITORS

[0221] A wide variety of compositions and administration methods may be used in conjunction with the HDAC inhibitors of the present invention. Such compositions may include, in addition to the HDAC inhibitors of the present invention, conventional

pharmaceutical excipients, and other conventional, pharmaceutically inactive agents. Additionally, the compositions may include active agents in addition to the HDAC inhibitors of the present invention. These additional active agents may include additional compounds according to the invention, or one or more other pharmaceutically active agents.

[0222] The compositions may be in gaseous, liquid, semi-liquid or solid form, formulated in a manner suitable for the route of administration to be used. For oral administration, capsules and tablets are typically used. For parenteral administration, reconstitution of a lyophilized powder, prepared as described herein, is typically used.

[0223] Compositions comprising HDAC inhibitors of the present invention may be administered or coadministered orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery (for example by catheter or stent), subcutaneously, intraadiposally, intraarticularly, or intrathecally. The compounds and/or compositions according to the invention may also be administered or coadministered in slow release dosage forms.

[0224] The HDAC inhibitors and compositions comprising them may be administered or coadministered in any conventional dosage form. Coadministration in the context of this invention is intended to mean the administration of more than one therapeutic agents, one of which includes a HDAC inhibitor, in the course of a coordinated treatment to achieve an improved clinical outcome. Such coadministration may also be coextensive, that is, occurring during overlapping periods of time.

[0225] Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application may optionally include one or more of the following components: a sterile diluent, such as water for injection, saline solution, fixed oil, polyethylene glycol, glycerine, propylene glycol or other synthetic solvent; antimicrobial agents, such as benzyl alcohol and methyl parabens; antioxidants, such as ascorbic acid and sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid (EDTA); buffers, such as acetates, citrates and phosphates; agents for the adjustment of tonicity such as sodium chloride or dextrose, and agents for adjusting the acidity or alkalinity of the composition, such as alkaline or acidifying agents or buffers like carbonates, bicarbonates, phosphates, hydrochloric acid, and organic acids like acetic and citric acid. Parenteral preparations may optionally be enclosed in ampules,

disposable syringes or single or multiple dose vials made of glass, plastic or other suitable material.

[0226] When HDAC inhibitors according to the present invention exhibit insufficient solubility, methods for solubilizing the compounds may be used. Such methods are known to those of skill in this art, and include, but are not limited to, using cosolvents, such as dimethylsulfoxide (DMSO), using surfactants, such as TWEEN, or dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such as prodrugs of the compounds may also be used in formulating effective pharmaceutical compositions.

[0227] Upon mixing or adding HDAC inhibitors according to the present invention to a composition, a solution, suspension, emulsion or the like may be formed. The form of the resulting composition will depend upon a number of factors, including the intended mode of administration, and the solubility of the compound in the selected carrier or vehicle. The effective concentration needed to ameliorate the disease being treated may be empirically determined.

[0228] Compositions according to the present invention are optionally provided for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, dry powders for inhalers, granules, sterile parenteral solutions or suspensions, and oral solutions or suspensions, and oil-water emulsions containing suitable quantities of the compounds, particularly the pharmaceutically acceptable salts, preferably the sodium salts, thereof. The pharmaceutically therapeutically active compounds and derivatives thereof are typically formulated and administered in unit-dosage forms or multiple-dosage forms. Unit-dose forms, as used herein, refers to physically discrete units suitable for human and animal subjects and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of the therapeutically active compound sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes individually packaged tablet or capsule. Unit-dose forms may be administered in fractions or multiples thereof. A multiple-dose form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dose form. Examples of multiple-dose forms include vials, bottles of tablets or capsules or bottles of pint or gallons. Hence, multiple dose form is a multiple of unit-doses that are not segregated in packaging.

[0229] In addition to one or more HDAC inhibitors according to the present invention, the composition may comprise: a diluent such as lactose, sucrose, dicalcium phosphate, or carboxymethylcellulose; a lubricant, such as magnesium stearate, calcium stearate and talc; and a binder such as starch, natural gums, such as gum acaciagelatin, glucose, molasses, polyvinylpyrrolidone, celluloses and derivatives thereof, povidone, crospovidones and other such binders known to those of skill in the art. Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of auxiliary substances such as wetting agents, emulsifying agents, or solubilizing agents, pH buffering agents and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Actual methods of preparing such dosage forms are known in the art, or will be apparent, to those skilled in this art; for example, see Remington: The Science and Practice of Pharmacy, A. Gennaro, ed., 20th edition, Lippincott, Williams & Wilkins, Philadelphia, PA, 2000. The composition or formulation to be administered will, in any event, contain a sufficient quantity of a HDAC inhibitor of the present invention to reduce HDAC activity *in vivo*, thereby treating the disease state of the subject.

[0230] Dosage forms or compositions may optionally comprise one or more HDAC inhibitors according to the present invention in the range of 0.005% to 100% (weight/weight) with the balance comprising additional substances such as those described herein. For oral administration, a pharmaceutically acceptable composition may optionally comprise any one or more commonly employed excipients, such as, for example pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, talcum, cellulose derivatives, sodium crosscarmellose, glucose, sucrose, magnesium carbonate, sodium saccharin, talcum. Such compositions include solutions, suspensions, tablets, capsules, powders, dry powders for inhalers and sustained release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as collagen, ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic

acid and others. Methods for preparing these formulations are known to those skilled in the art. The compositions may optionally contain 0.01%-100% (weight/weight) of one or more HDAC inhibitors, optionally 0.1-95%, and optionally 1-95%.

[0231] Salts, preferably sodium salts, of the HDAC inhibitors may be prepared with carriers that protect the compound against rapid elimination from the body, such as time release formulations or coatings. The formulations may further include other active compounds to obtain desired combinations of properties.

A. Formulations For Oral Administration

[0232] Oral pharmaceutical dosage forms may be as a solid, gel or liquid. Examples of solid dosage forms include, but are not limited to tablets, capsules, granules, and bulk powders. More specific examples of oral tablets include compressed, chewable lozenges and tablets that may be enteric-coated, sugar-coated or film-coated. Examples of capsules include hard or soft gelatin capsules. Granules and powders may be provided in non-effervescent or effervescent forms. Each may be combined with other ingredients known to those skilled in the art.

[0233] In certain embodiments, HDAC inhibitors according to the present invention are provided as solid dosage forms, preferably capsules or tablets. The tablets, pills, capsules, troches and the like may optionally contain one or more of the following ingredients, or compounds of a similar nature: a binder; a diluent; a disintegrating agent; a lubricant; a glidant; a sweetening agent; and a flavoring agent.

[0234] Examples of binders that may be used include, but are not limited to, microcrystalline cellulose, gum tragacanth, glucose solution, acacia mucilage, gelatin solution, sucrose and starch paste.

[0235] Examples of lubricants that may be used include, but are not limited to, talc, starch, magnesium or calcium stearate, lycopodium and stearic acid.

[0236] Examples of diluents that may be used include, but are not limited to, lactose, sucrose, starch, kaolin, salt, mannitol and dicalcium phosphate.

[0237] Examples of glidants that may be used include, but are not limited to, colloidal silicon dioxide.

[0238] Examples of disintegrating agents that may be used include, but are not limited to, crosscarmellose sodium, sodium starch glycolate, alginic acid, corn starch, potato starch, bentonite, methylcellulose, agar and carboxymethylcellulose.

[0239] Examples of coloring agents that may be used include, but are not limited to, any of the approved certified water soluble FD and C dyes, mixtures thereof; and water insoluble FD and C dyes suspended on alumina hydrate.

[0240] Examples of sweetening agents that may be used include, but are not limited to, sucrose, lactose, mannitol and artificial sweetening agents such as sodium cyclamate and saccharin, and any number of *spray-dried* flavors.

[0241] Examples of flavoring agents that may be used include, but are not limited to, natural flavors extracted from plants such as fruits and synthetic blends of compounds that produce a pleasant sensation, such as, but not limited to peppermint and methyl salicylate.

[0242] Examples of wetting agents that may be used include, but are not limited to, propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate and polyoxyethylene lauryl ether.

[0243] Examples of anti-emetic coatings that may be used include, but are not limited to, fatty acids, fats, waxes, shellac, ammoniated shellac and cellulose acetate phthalates.

[0244] Examples of film coatings that may be used include, but are not limited to, hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate.

[0245] If oral administration is desired, the salt of the compound may optionally be provided in a composition that protects it from the acidic environment of the stomach. For example, the composition can be formulated in an *enteric-coating* that maintains its integrity in the stomach and releases the active compound in the intestine. The composition may also be formulated in combination with an antacid or other such ingredient.

[0246] When the dosage unit form is a capsule, it may optionally additionally comprise a liquid carrier such as a fatty oil. In addition, dosage unit forms may optionally additionally comprise various other materials that modify the physical form of the dosage unit, for example, coatings of sugar and other enteric agents.

[0247] Compounds according to the present invention may also be administered as a component of an elixir, suspension, syrup, wafer, sprinkle, chewing gum or the like. A syrup

may optionally comprise, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

[0248] The HDAC inhibitors of the present invention may also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, such as antacids, H₂ blockers, and diuretics. For example, if a compound is used for treating asthma or hypertension, it may be used with other bronchodilators and antihypertensive agents, respectively.

[0249] Examples of pharmaceutically acceptable carriers that may be included in tablets comprising HDAC inhibitors of the present invention include, but are not limited to binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, and wetting agents. Enteric-coated tablets, because of the enteric-coating, resist the action of stomach acid and dissolve or disintegrate in the neutral or alkaline intestines. Sugar-coated tablets may be compressed tablets to which different layers of pharmaceutically acceptable substances are applied. Film-coated tablets may be compressed tablets that have been coated with polymers or other suitable coating. Multiple compressed tablets may be compressed tablets made by more than one compression cycle utilizing the pharmaceutically acceptable substances previously mentioned. Coloring agents may also be used in tablets. Flavoring and sweetening agents may be used in tablets, and are especially useful in the formation of chewable tablets and lozenges.

[0250] Examples of liquid oral dosage forms that may be used include, but are not limited to, aqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules and effervescent preparations reconstituted from effervescent granules.

[0251] Examples of aqueous solutions that may be used include, but are not limited to, elixirs and syrups. As used herein, elixirs refer to clear, sweetened, hydroalcoholic preparations. Examples of pharmaceutically acceptable carriers that may be used in elixirs include, but are not limited to solvents. Particular examples of solvents that may be used include glycerin, sorbitol, ethyl alcohol and syrup. As used herein, syrups refer to concentrated aqueous solutions of a sugar, for example, sucrose. Syrups may optionally further comprise a preservative.

[0252] Emulsions refer to two-phase systems in which one liquid is dispersed in the form of small globules throughout another liquid. Emulsions may optionally be oil-in-water or water-in-oil emulsions. Examples of pharmaceutically acceptable carriers that may be used in emulsions include, but are not limited to non-aqueous liquids, emulsifying agents and preservatives.

[0253] Examples of pharmaceutically acceptable substances that may be used in non-effervescent granules, to be reconstituted into a liquid oral dosage form, include diluents, sweeteners and wetting agents.

[0254] Examples of pharmaceutically acceptable substances that may be used in effervescent granules, to be reconstituted into a liquid oral dosage form, include organic acids and a source of carbon dioxide.

[0255] Coloring and flavoring agents may optionally be used in all of the above dosage forms.

[0256] Particular examples of preservatives that may be used include glycerin, methyl and propylparaben, benzoic acid, sodium benzoate and alcohol.

[0257] Particular examples of non-aqueous liquids that may be used in emulsions include mineral oil and cottonseed oil.

[0258] Particular examples of emulsifying agents that may be used include gelatin, acacia, tragacanth, bentonite, and surfactants such as polyoxyethylene sorbitan monooleate.

[0259] Particular examples of suspending agents that may be used include sodium carboxymethylcellulose, pectin, tragacanth, Veegum and acacia. Diluents include lactose and sucrose. Sweetening agents include sucrose, syrups, glycerin and artificial sweetening agents such as sodium cyclamate and saccharin.

[0260] Particular examples of wetting agents that may be used include propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate and polyoxyethylene lauryl ether.

[0261] Particular examples of organic acids that may be used include citric and tartaric acid.

[0262] Sources of carbon dioxide that may be used in effervescent compositions include sodium bicarbonate and sodium carbonate. Coloring agents include any of the approved certified water soluble FD and C dyes, and mixtures thereof.

[0263] Particular examples of flavoring agents that may be used include natural flavors extracted from plants such fruits, and synthetic blends of compounds that produce a pleasant taste sensation.

[0264] For a solid dosage form, the solution or suspension, in for example propylene carbonate, vegetable oils or triglycerides, is preferably encapsulated in a gelatin capsule. Such solutions, and the preparation and encapsulation thereof, are disclosed in U.S. Pat. Nos. 4,328,245; 4,409,239; and 4,410,545. For a liquid dosage form, the solution, egg., for example, in a polyethylene glycol, may be diluted with a sufficient quantity of a pharmaceutically acceptable liquid carrier, e.g. water, to be easily measured for administration.

[0265] Alternatively, liquid or semi-solid oral formulations may be prepared by dissolving or dispersing the active compound or salt in vegetable oils, glycols, triglycerides, propylene glycol esters (e.g. propylene carbonate) and other such carriers, and encapsulating these solutions or suspensions in hard or soft gelatin capsule shells. Other useful formulations include those set forth in U.S. Pat. Nos. Re 28,819 and 4,358,603.

B. Injectables, Solutions and Emulsions

[0266] The present invention is also directed to compositions designed to administer the HDAC inhibitors of the present invention by parenteral administration, generally characterized by injection, either subcutaneously, intramuscularly or intravenously. Injectables may be prepared in any conventional form, for example as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions.

[0267] Examples of excipients that may be used in conjunction with injectables according to the present invention include, but are not limited to water, saline, dextrose, glycerol or ethanol. The injectable compositions may also optionally comprise minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents, stabilizers, solubility enhancers, and other such agents, such as for example, sodium acetate, sorbitan monolaurate, triethanolamine oleate and cyclodextrins. Implantation of a slow-release or sustained-release system, such that a constant level of dosage is maintained (see, e.g., U.S. Pat. No. 3,710,795) is also contemplated herein. The percentage of active compound contained in such parenteral compositions is highly dependent on the specific nature thereof, as well as the activity of the compound and the needs of the subject.

[0268] Parenteral administration of the formulations includes intravenous, subcutaneous and intramuscular administrations. Preparations for parenteral administration include sterile solutions ready for injection, sterile dry soluble products, such as the lyophilized powders described herein, ready to be combined with a solvent just prior to use, including hypodermic tablets, sterile suspensions ready for injection, sterile dry insoluble products ready to be combined with a vehicle just prior to use and sterile emulsions. The solutions may be either aqueous or nonaqueous.

[0269] When administered intravenously, examples of suitable carriers include, but are not limited to physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, and polypropylene glycol and mixtures thereof.

[0270] Example of pharmaceutically acceptable carriers that may optionally be used in parenteral preparations include, but are not limited to aqueous vehicles, nonaqueous vehicles, antimicrobial agents, isotonic agents, buffers, antioxidants, local anesthetics, suspending and dispersing agents, emulsifying agents, sequestering or chelating agents and other pharmaceutically acceptable substances.

[0271] Examples of aqueous vehicles that may optionally be used include Sodium Chloride Injection, Ringers Injection, Isotonic Dextrose Injection, Sterile Water Injection, Dextrose and Lactated Ringers Injection.

[0272] Examples of nonaqueous parenteral vehicles that may optionally be used include fixed oils of vegetable origin, cottonseed oil, corn oil, sesame oil and peanut oil.

[0273] Antimicrobial agents in bacteriostatic or fungistatic concentrations may be added to parenteral preparations, particularly when the preparations are packaged in multiple-dose containers and thus designed to be stored and multiple aliquots to be removed. Examples of antimicrobial agents that may be used include phenols or cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl p-hydroxybenzoic acid esters, thimerosal, benzalkonium chloride and benzethonium chloride.

[0274] Examples of isotonic agents that may be used include sodium chloride and dextrose. Examples of buffers that may be used include phosphate and citrate. Examples of antioxidants that may be used include sodium bisulfate. Examples of local anesthetics that may be used include procaine hydrochloride. Examples of suspending and dispersing agents

that may be used include sodium carboxymethylcellulose, hydroxypropyl methylcellulose and polyvinylpyrrolidone. Examples of emulsifying agents that may be used include Polysorbate 80 (Tween 80). A sequestering or chelating agent of metal ions include EDTA.

[0275] Pharmaceutical carriers may also optionally include ethyl alcohol, polyethylene glycol and propylene glycol for water miscible vehicles and sodium hydroxide, hydrochloric acid, citric acid or lactic acid for pH adjustment.

[0276] The concentration of a HDAC inhibitor in the parenteral formulation may be adjusted so that an injection administers a pharmaceutically effective amount sufficient to produce the desired pharmacological effect. The exact concentration of a HDAC inhibitor and/or dosage to be used will ultimately depend on the age, weight and condition of the patient or animal as is known in the art.

[0277] Unit-dose parenteral preparations may be packaged in an ampoule, a vial or a syringe with a needle. All preparations for parenteral administration should be sterile, as is known and practiced in the art.

[0278] Injectables may be designed for local and systemic administration. Typically a therapeutically effective dosage is formulated to contain a concentration of at least about 0.1% w/w up to about 90% w/w or more, preferably more than 1% w/w of the HDAC inhibitor to the treated tissue(s). The HDAC inhibitor may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment will be a function of the location of where the composition is parenterally administered, the carrier and other variables that may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test data. It is to be noted that concentrations and dosage values may also vary with the age of the individual treated. It is to be further understood that for any particular subject, specific dosage regimens may need to be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations. Hence, the concentration ranges set forth herein are intended to be exemplary and are not intended to limit the scope or practice of the claimed formulations.

[0279] The HDAC inhibitor may optionally be suspended in micronized or other suitable form or may be derivatized to produce a more soluble active product or to produce a prodrug. The form of the resulting mixture depends upon a number of factors, including the intended

mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for ameliorating the symptoms of the disease state and may be empirically determined.

C. Lyophilized Powders

[0280] The HDAC inhibitors of the present invention may also be prepared as lyophilized powders, which can be reconstituted for administration as solutions, emulsions and other mixtures. The lyophilized powders may also be formulated as solids or gels.

[0281] Sterile, lyophilized powder may be prepared by dissolving the sodium salt in a sodium phosphate buffer solution containing dextrose or other suitable excipient. Subsequent sterile filtration of the solution followed by lyophilization under standard conditions known to those of skill in the art provides the desired formulation. Briefly, the lyophilized powder may optionally be prepared by dissolving dextrose, sorbitol, fructose, corn syrup, xylitol, glycerin, glucose, sucrose or other suitable agent, about 1-20%, preferably about 5 to 15%, in a suitable buffer, such as citrate, sodium or potassium phosphate or other such buffer known to those of skill in the art at, typically, about neutral pH. Then, a HDAC inhibitor is added to the resulting mixture, preferably above room temperature, more preferably at about 30-35 °C, and stirred until it dissolves. The resulting mixture is diluted by adding more buffer to a desired concentration. The resulting mixture is sterile filtered or treated to remove particulates and to insure sterility, and apportioned into vials for lyophilization. Each vial may contain a single dosage or multiple dosages of the HDAC inhibitor.

D. Topical Administration

[0282] The HDAC inhibitors of the present invention may also be administered as topical mixtures. Topical mixtures may be used for local and systemic administration. The resulting mixture may be a solution, suspension, emulsions or the like and are formulated as creams, gels, ointments, emulsions, solutions, elixirs, lotions, suspensions, tinctures, pastes, foams, aerosols, irrigations, sprays, suppositories, bandages, dermal patches or any other formulations suitable for topical administration.

[0283] The HDAC inhibitors may be formulated as aerosols for topical application, such as by inhalation (see, U.S. Pat. Nos. 4,044,126, 4,414,209, and 4,364,923, which describe aerosols for delivery of a steroid useful for treatment inflammatory diseases, particularly asthma). These formulations for administration to the respiratory tract can be in the form of

an aerosol or solution for a nebulizer, or as a microfine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will typically have diameters of less than 50 microns, preferably less than 10 microns.

[0284] The HDAC inhibitors may also be formulated for local or topical application, such as for topical application to the skin and mucous membranes, such as in the eye, in the form of gels, creams, and lotions and for application to the eye or for intracisternal or intraspinal application. Topical administration is contemplated for transdermal delivery and also for administration to the eyes or mucosa, or for inhalation therapies. Nasal solutions of the HDAC inhibitor alone or in combination with other pharmaceutically acceptable excipients can also be administered.

E. Formulations For Other Routes of Administration

[0285] Depending upon the disease state being treated, other routes of administration, such as topical application, transdermal patches, a rectal administration, may also be used. For example, pharmaceutical dosage forms for rectal administration are rectal suppositories, capsules and tablets for systemic effect. Rectal suppositories are used herein to mean solid bodies for insertion into the rectum that melt or soften at body temperature releasing one or more pharmacologically or therapeutically active ingredients. Pharmaceutically acceptable substances utilized in rectal suppositories are bases or vehicles and agents to raise the melting point. Examples of bases include cocoa butter (theobroma oil), glycerin-gelatin, carbowax, (polyoxyethylene glycol) and appropriate mixtures of mono-, di- and triglycerides of fatty acids. Combinations of the various bases may be used. Agents to raise the melting point of suppositories include spermaceti and wax. Rectal suppositories may be prepared either by the compressed method or by molding. The typical weight of a rectal suppository is about 2 to 3 gm. Tablets and capsules for rectal administration may be manufactured using the same pharmaceutically acceptable substance and by the same methods as for formulations for oral administration.

F. Examples of Formulations

[0286] The following are particular examples of oral, intravenous and tablet formulations that may optionally be used with compounds of the present invention. It is noted that these formulations may be varied depending on the particular compound being used and the indication for which the formulation is going to be used.

ORAL FORMULATION

Compound of the Present Invention	10-100 mg
Citric Acid Monohydrate	105 mg
Sodium Hydroxide	18 mg
Flavoring	
Water	q.s. to 100 mL

INTRAVENOUS FORMULATION

Compound of the Present Invention	0.1-10 mg
Dextrose Monohydrate	q.s. to make isotonic
Citric Acid Monohydrate	1.05 mg
Sodium Hydroxide	0.18 mg
Water for Injection	q.s. to 1.0 mL

TABLET FORMULATION

Compound of the Present Invention	1%
Microcrystalline Cellulose	73%
Stearic Acid	25%
Colloidal Silica	1%

6. KITS COMPRISING HDAC INHIBITORS

[0287] The invention is also directed to kits and other articles of manufacture for treating diseases associated with HDAC. It is noted that diseases are intended to cover all conditions for which the HDAC possesses activity that contributes to the pathology and/or symptomology of the condition.

[0288] In one embodiment, a kit is provided that comprises a composition comprising at least one HDAC inhibitor of the present invention in combination with instructions. The instructions may indicate the disease state for which the composition is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition. The kit may also comprise packaging materials. The packaging material may comprise a container for housing the composition. The kit may also optionally comprise additional components, such as syringes for administration of the composition. The kit may comprise the composition in single or multiple dose forms.

[0289] In another embodiment, an article of manufacture is provided that comprises a composition comprising at least one HDAC inhibitor of the present invention in combination with packaging materials. The packaging material may comprise a container for housing the composition. The container may optionally comprise a label indicating the disease state for

which the composition is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition. The kit may also optionally comprise additional components, such as syringes for administration of the composition. The kit may comprise the composition in single or multiple dose forms.

[0290] It is noted that the packaging material used in kits and articles of manufacture according to the present invention may form a plurality of divided containers such as a divided bottle or a divided foil packet. The container can be in any conventional shape or form as known in the art which is made of a pharmaceutically acceptable material, for example a paper or cardboard box, a glass or plastic bottle or jar, a re-sealable bag (for example, to hold a "refill" of tablets for placement into a different container), or a blister pack with individual doses for pressing out of the pack according to a therapeutic schedule. The container that is employed will depend on the exact dosage form involved, for example a conventional cardboard box would not generally be used to hold a liquid suspension. It is feasible that more than one container can be used together in a single package to market a single dosage form. For example, tablets may be contained in a bottle that is in turn contained within a box. Typically the kit includes directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral, topical, transdermal and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician.

[0291] One particular example of a kit according to the present invention is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process recesses are formed in the plastic foil. The recesses have the size and shape of individual tablets or capsules to be packed or may have the size and shape to accommodate multiple tablets and/or capsules to be packed. Next, the tablets or capsules are placed in the recesses accordingly and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are individually sealed or collectively sealed, as desired, in the recesses between the plastic foil

and the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.

[0292] Another specific embodiment of a kit is a dispenser designed to dispense the daily doses one at a time in the order of their intended use. Preferably, the dispenser is equipped with a memory-aid, so as to further facilitate compliance with the regimen. An example of such a memory-aid is a mechanical counter that indicates the number of daily doses that has been dispensed. Another example of such a memory-aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.

7. COMBINATION THERAPY

[0293] A wide variety therapeutic agents may have a therapeutic additive or synergistic effect with HDAC inhibitors according to the present invention. Such therapeutic agents may additively or synergistically combine with the HDAC inhibitors to inhibit undesirable cell growth, such as inappropriate cell growth resulting in undesirable benign conditions or tumor growth.

[0294] In one embodiment, a method is provided for treating a cell proliferative disease state comprising treating cells with a compound according to the present invention in combination with an anti-proliferative agent, wherein the cells are treated with the compound according to the present invention before, at the same time, and/or after the cells are treated with the anti-proliferative agent, referred to herein as combination therapy. It is noted that treatment of one agent before another is referred to herein as sequential therapy, even if the agents are also administered together. It is noted that combination therapy is intended to cover when agents are administered before or after each other (sequential therapy) as well as when the agents are administered at the same time.

[0295] Examples of therapeutic agents that may be used in combination with HDAC inhibitors include, but are not limited to, anticancer agents, alkylating agents, antibiotic agents, antimetabolic agents, hormonal agents, plant-derived agents, and biologic agents.

[0296] Alkylating agents are polyfunctional compounds that have the ability to substitute alkyl groups for hydrogen ions. Examples of alkylating agents include, but are not limited to, bischloroethylamines (nitrogen mustards, e.g. chlorambucil, cyclophosphamide, ifosfamide, mechlorethamine, melphalan, uracil mustard), aziridines (e.g. thiotepa), alkyl alkone sulfonates (e.g. busulfan), nitrosoureas (e.g. carmustine, lomustine, streptozocin), nonclassic alkylating agents (altretamine, dacarbazine, and procarbazine), platinum compounds (carboplatin and cisplatin). These compounds react with phosphate, amino, hydroxyl, sulfhydryl, carboxyl, and imidazole groups. Under physiological conditions, these drugs ionize and produce positively charged ion that attach to susceptible nucleic acids and proteins, leading to cell cycle arrest and/or cell death. Combination therapy including a HDAC inhibitor and an alkylating agent may have therapeutic synergistic effects on cancer and reduce sides affects associated with these chemotherapeutic agents.

[0297] Antibiotic agents are a group of drugs that produced in a manner similar to antibiotics as a modification of natural products. Examples of antibiotic agents include, but are not limited to, anthracyclines (e.g. doxorubicin, daunorubicin, epirubicin, idarubicin and anthracenedione), mitomycin C, bleomycin, dactinomycin, plicatomycin. These antibiotic agents interfere with cell growth by targeting different cellular components. For example, anthracyclines are generally believed to interfere with the action of DNA topoisomerase II in the regions of transcriptionally active DNA, which leads to DNA strand scissions. Bleomycin is generally believed to chelate iron and forms an activated complex, which then binds to bases of DNA, causing strand scissions and cell death. Combination therapy including a HDAC inhibitor and an antibiotic agent may have therapeutic synergistic effects on cancer and reduce sides affects associated with these chemotherapeutic agents.

[0298] Antimetabolic agents are a group of drugs that interfere with metabolic processes vital to the physiology and proliferation of cancer cells. Actively proliferating cancer cells require continuous synthesis of large quantities of nucleic acids, proteins, lipids, and other vital cellular constituents. Many of the antimetabolites inhibit the synthesis of purine or pyrimidine nucleosides or inhibit the enzymes of DNA replication. Some antimetabolites also interfere with the synthesis of ribonucleosides and RNA and/or amino acid metabolism and protein synthesis as well. By interfering with the synthesis of vital cellular constituents, antimetabolites can delay or arrest the growth of cancer cells. Examples of antimetabolic

agents include, but are not limited to, fluorouracil (5-FU), floxuridine (5-FUdR), methotrexate, leucovorin, hydroxyurea, thioguanine (6-TG), mercaptopurine (6-MP), cytarabine, pentostatin, fludarabine phosphate, cladribine (2-CDA), asparaginase, and gemcitabine. Combination therapy including a HDAC inhibitor and a antimetabolic agent may have therapeutic synergistic effects on cancer and reduce sides affects associated with these chemotherapeutic agents.

[0299] Hormonal agents are a group of drug that regulate the growth and development of their target organs. Most of the hormonal agents are sex steroids and their derivatives and analogs thereof, such as estrogens, androgens, and progestins. These hormonal agents may serve as antagonists of receptors for the sex steroids to down regulate receptor expression and transcription of vital genes. Examples of such hormonal agents are synthetic estrogens (e.g. diethylstilbestrol), antiestrogens (e.g. tamoxifen, toremifene, fluoxymesterol and raloxifene), antiandrogens (bicalutamide, nilutamide, flutamide), aromatase inhibitors (e.g., aminoglutethimide, anastrozole and tetrazole), ketoconazole, goserelin acetate, leuprolide, megestrol acetate and mifepristone. Combination therapy including a HDAC inhibitor and a hormonal agent may have therapeutic synergistic effects on cancer and reduce sides affects associated with these chemotherapeutic agents.

[0300] Plant-derived agents are a group of drugs that are derived from plants or modified based on the molecular structure of the agents. Examples of plant-derived agents include, but are not limited to, vinca alkaloids (e.g., vincristine, vinblastine, vindesine, vinzolidine and vinorelbine), podophyllotoxins (e.g., etoposide (VP-16) and teniposide (VM-26)), taxanes (e.g., paclitaxel and docetaxel). These plant-derived agents generally act as antimitotic agents that bind to tubulin and inhibit mitosis. Podophyllotoxins such as etoposide are believed to interfere with DNA synthesis by interacting with topoisomerase II, leading to DNA strand scission. Combination therapy including a HDAC inhibitor and a plant-derived agent may have therapeutic synergistic effects on cancer and reduce sides affects associated with these chemotherapeutic agents.

[0301] Biologic agents are a group of biomolecules that elicit cancer/tumor regression when used alone or in combination with chemotherapy and/or radiotherapy. Examples of biologic agents include, but are not limited to, immuno-modulating proteins such as cytokines, monoclonal antibodies against tumor antigens, tumor suppressor genes, and cancer vaccines.

Combination therapy including a HDAC inhibitor and a biologic agent may have therapeutic synergistic effects on cancer, enhance the patient's immune responses to tumorigenic signals, and reduce potential side effects associated with this chemotherapeutic agent.

[0302] Cytokines possess profound immunomodulatory activity. Some cytokines such as interleukin-2 (IL-2, aldesleukin) and interferon have demonstrated antitumor activity and have been approved for the treatment of patients with metastatic renal cell carcinoma and metastatic malignant melanoma. IL-2 is a T-cell growth factor that is central to T-cell-mediated immune responses. The selective antitumor effects of IL-2 on some patients are believed to be the result of a cell-mediated immune response that discriminates between self and nonself. Examples of interleukins that may be used in conjunction with HDAC inhibitor include, but are not limited to, interleukin 2 (IL-2), and interleukin 4 (IL-4), interleukin 12 (IL-12).

[0303] Interferon include more than 23 related subtypes with overlapping activities, all of the IFN subtypes within the scope of the present invention. IFN has demonstrated activity against many solid and hematologic malignancies, the later appearing to be particularly sensitive.

[0304] Other cytokines that may be used in conjunction with a HDAC inhibitor include those cytokines that exert profound effects on hematopoiesis and immune functions. Examples of such cytokines include, but are not limited to erythropoietin, granulocyte-CSF (filgrastin), and granulocyte, macrophage-CSF (sargramostim). These cytokines may be used in conjunction with a HDAC inhibitor to reduce chemotherapy-induced myelopoietic toxicity.

[0305] Other immuno-modulating agents other than cytokines may also be used in conjunction with a HDAC inhibitor to inhibit abnormal cell growth. Examples of such immuno-modulating agents include, but are not limited to bacillus Calmette-Guerin, levamisole, and octreotide, a long-acting octapeptide that mimics the effects of the naturally occurring hormone somatostatin.

[0306] Monoclonal antibodies against tumor antigens are antibodies elicited against antigens expressed by tumors, preferably tumor-specific antigens. For example, monoclonal antibody HERCEPTIN® (Trastuzumab) is raised against human epidermal growth factor receptor2 (HER2) that is overexpressed in some breast tumors including metastatic breast cancer. Overexpression of HER2 protein is associated with more aggressive disease and poorer prognosis in the clinic. HERCEPTIN® is used as a single agent for the treatment of

patients with metastatic breast cancer whose tumors over express the HER2 protein. Combination therapy including HDAC inhibitor and HERCEPTIN® may have therapeutic synergistic effects on tumors, especially on metastatic cancers.

[0307] Another example of monoclonal antibodies against tumor antigens is RITUXAN® (Rituximab) that is raised against CD20 on lymphoma cells and selectively deplete normal and malignant CD20⁺ pre-B and mature B cells. RITUXAN® is used as single agent for the treatment of patients with relapsed or refractory low-grade or follicular, CD20+, B cell non-Hodgkin's lymphoma. Combination therapy including HDAC inhibitor and RITUXAN® may have therapeutic synergistic effects not only on lymphoma, but also on other forms or types of malignant tumors.

[0308] Tumor suppressor genes are genes that function to inhibit the cell growth and division cycles, thus preventing the development of neoplasia. Mutations in tumor suppressor genes cause the cell to ignore one or more of the components of the network of inhibitory signals, overcoming the cell cycle check points and resulting in a higher rate of controlled cell growth—cancer. Examples of the tumor suppressor genes include, but are not limited to, *DPC-4*, *NF-1*, *NF-2*, *RB*, *p53*, *WT1*, *BRCA1* and *BRCA2*.

[0309] *DPC-4* is involved in pancreatic cancer and participates in a cytoplasmic pathway that inhibits cell division. *NF-1* codes for a protein that inhibits Ras, a cytoplasmic inhibitory protein. *NF-1* is involved in neurofibroma and pheochromocytomas of the nervous system and myeloid leukemia. *NF-2* encodes a nuclear protein that is involved in meningioma, schwannoma, and ependymoma of the nervous system. *RB* codes for the pRB protein, a nuclear protein that is a major inhibitor of cell cycle. *RB* is involved in retinoblastoma as well as bone, bladder, small cell lung and breast cancer. *P53* codes for p53 protein that regulates cell division and can induce apoptosis. Mutation and/or inaction of p53 is found in a wide ranges of cancers. *WT1* is involved in Wilm's tumor of the kidneys. *BRCA1* is involved in breast and ovarian cancer, and *BRCA2* is involved in breast cancer. The tumor suppressor gene can be transferred into the tumor cells where it exerts its tumor suppressing functions. Combination therapy including a HDAC inhibitor and a tumor suppressor may have therapeutic synergistic effects on patients suffering from various forms of cancers.

[0310] Cancer vaccines are a group of agents that induce the body's specific immune response to tumors. Most of cancer vaccines under research and development and clinical

trials are tumor-associated antigens (TAAs). TAA are structures (i.e. proteins, enzymes or carbohydrates) which are present on tumor cells and relatively absent or diminished on normal cells. By virtue of being fairly unique to the tumor cell, TAAs provide targets for the immune system to recognize and cause their destruction. Example of TAAs include, but are not limited to gangliosides (GM2), prostate specific antigen (PSA), alpha-fetoprotein (AFP), carcinoembryonic antigen (CEA) (produced by colon cancers and other adenocarcinomas, e.g. breast, lung, gastric, and pancreas cancer s), melanoma associated antigens (MART-1, gp100, MAGE 1,3 tyrosinase), papillomavirus E6 and E7 fragments, whole cells or portions/lysates of autologous tumor cells and allogeneic tumor cells.

[0311] An adjuvant may be used to augment the immune response to TAAs. Examples of adjuvants include, but are not limited to, bacillus Calmette-Guerin (BCG), endotoxin lipopolysaccharides, keyhole limpet hemocyanin (GKLH), interleukin-2 (IL-2), granulocyte-macrophage colony-stimulating factor (GM-CSF) and cytoxan, a chemotherapeutic agent which is believed to reduce tumor-induced suppression when given in low doses.

3. HDAC ACTIVITY ASSAY

[0312] Compounds according to the present invention may be screened for activity against one or more HDACs. Provided in this example are assays for activity against HDAC1, HDAC2, HDAC6 and HDAC8.

[0313] Purified HDAC1, HDAC2, HDAC6, and HDAC8 may be obtained as follows.

[0314] For HDAC1, DNA encoding residues 1-482 of the full-length sequence of the human enzyme may be amplified by PCR and cloned into the BamHI/XbaI sites of pFastbac (Invitrogen), which incorporates a 6-histidine tag at the N-terminus. SEQ. I.D. No. 1 corresponds to residues 1-482 with the N-terminal 6-histidine tag and SEQ. I.D. No. 2 is the DNA sequence that was used to encode SEQ. I.D. No. 1.

[0315] For HDAC2, DNA encoding residues 1-488 of the full-length sequence of the human enzyme may be amplified by PCR and cloned into the BamHI/SmaI sites of pFastbac (Invitrogen), which incorporates a 6-histidine tag at the C-terminus. SEQ. I.D. No. 3 corresponds to residues 1-488 with the C-terminal 6-histidine tag and SEQ. I.D. No. 4 is the DNA sequence that was used to encode SEQ. I.D. No. 3.

[0316] For HDAC6, DNA encoding residues 73-845 of the human enzyme may be amplified by PCR and cloned into the SmaI site of pFastbac (Invitrogen), which incorporates a

6xHistidine tag at the C-terminus. SEQ. I.D. No. 5 corresponds to residues 73-845 with the C-terminal 6-histidine tag and SEQ. I.D. No. 6 is the DNA sequence that was used to encode SEQ. I.D. No. 5.

[0317] For HDAC8, DNA encoding residues 1-377 corresponding to the entire sequence of the human enzyme may be amplified by PCR and cloned into the BamHI/SmaI sites of pFastbac (Invitrogen), which incorporates a 6-histidine tag at the N-terminus. SEQ. I.D. No. 7 corresponds to residues 1-377 with the N-terminal 6-histidine tag and SEQ. I.D. No. 8 is the DNA sequence that was used to encode SEQ. I.D. No. 7.

[0318] Recombinant baculovirus incorporating the HDAC constructs may be generated by transposition using the Bac-to-Bac system (Invitrogen). High-titer viral stocks may be generated by infection of *Spodoptera frugiperda* Sf9 cells; the expression of recombinant protein may be carried out by infection of *Spodoptera frugiperda* Sf9 or *Trichoplusia ni* Hi5 cells (Invitrogen) in 10L Wave Bioreactors (Wave Biotech).

[0319] Recombinant protein may be isolated from cellular extracts by passage over ProBond resin (Invitrogen). HDAC1 and HDAC6 may then be treated with TEV protease for the removal of the N-terminal 6xHistidine affinity tag (residual uncleaved protein may be removed through a second passage over ProBond Resin). Partially purified extracts of all HDACs may then be further purified by high pressure liquid chromatography over a BioSep S3000 gel filtration resin. The purity of HDAC proteins may be determined on denaturing SDS-PAGE gel. Purified HDACs may then be concentrated to a final concentration of 4.0 mg/ml for HDAC1, 10 mg/ml for HDAC2, 4.0 mg/ml for HDAC6, and 3 mg/ml for HDAC8. The proteins may be either stored at -78 °C in a buffer containing 25mM TRIS-HCl pH 7.6, 150mM NaCl, 0.1mM EDTA and 0.25 mM TCEP or at -20 °C in the presence of glycerol (final concentration of glycerol at 50%)

[0320] The inhibitory properties of compounds relative to HDAC1, HDAC2, HDAC6 and HDAC8 may be determined using a white or black 384-well-plate format under the following reaction conditions: 25 mM Tris pH 8.0, 100 mM NaCl, 50 mM KCl, 0.1 mM EDTA, 0.01% Brij35, 0.1 mM TCEP, 50 uM tBoc-Lys(Ac)-AMC, 2% DMSO. Reaction product may be determined quantitatively by fluorescence intensity using a Fluorescence plate reader (Molecular Devices Gemini) with an excitation wavelength at 370 nm and emission at 480 nm (for white plates) or 465 nm (for black plates).

[0321] The assay reaction may be initiated as follows: 5 ul of 150 uM tBoc-Lys(Ac)AMC was added to each well of the plate, followed by the addition of 5 ul of inhibitor (2 fold serial dilutions for 11 data points for each inhibitor) containing 6% DMSO. 5 ul of either HDAC1, HDAC2, HDAC6 or HDAC8 solution may be added to initiate the reaction (final enzyme concentrations were 2.5 nM for HDAC1, 1 nM for HDAC2, 2.5 nM for HDAC6 and 10 nM for HDAC8). The reaction mixture may then be incubated at room temperature for 60 min, and quenched and developed by addition of 5 ul of 10 mM phenanthroline and 4 mg/ml trypsin (final concentration of phenanthroline is 2.5 mM, and trypsin is 1 mg/ml). Fluorescence intensities of the resulting reaction mixtures may be measured after a 30 minute incubation at room temperature.

[0322] IC₅₀ values may be calculated by non-linear curve fitting of the compound concentrations and fluorescence intensities to the standard IC₅₀ equation. As a reference point for this assay, suberanolhydroxamic acid (SAHA) showed an IC₅₀ of 63 nM for HDAC1, 69 nM for HDAC2, 108 nM for HDAC6 and 242 nM for HDAC8.


[0323] The Section below provides examples of HDAC inhibitors that were assayed according to the above assays and found to have better than 1000 nM activity against HDAC1, HDAC2, HDAC6, and HDAC8.

EXAMPLES


1. Synthetic Schemes For HDAC Inhibitors

[0324] HDAC inhibitors according to the present invention may be synthesized according to a variety of reaction schemes. Some illustrative schemes are provided herein in the examples. Other reaction schemes could be readily devised by those skilled in the art.

Scheme 1:



Scheme 2:



General procedure for the synthesis of *N*¹-substituted-phenyl-1,2-diamines (2).

[0325] To a solution of the appropriate 1-fluoro-2-nitrobenzene (1, 1.77 mmol) and the appropriate amine (1.77 mmol) in DMF (5.0 mL) was added DIEA (1.94 mmol). The reaction was heated at 50-100 °C for 24-48 hrs and then cooled to ambient temperature. The resulting mixture was poured into H₂O, extracted with EtOAc, washed with brine, and dried over MgSO₄. The organic layer was evaporated to dryness and the resulting material was purified if needed via flash chromatography to yield the desired 2-nitrophenylamines. The appropriate *N*-substituted-2-nitrophenylamine (1.10 mmol) was dissolved in MeOH/AcOH (4:1, 5.0 mL) and heated to 100 °C. Zinc dust (5.50 mmol) was added to the reaction portion wise until frothing ceased. The reaction was cooled to ambient temperature, filtered, and evaporated to dryness. The resulting *N*¹-substituted-phenyl-1,2-diamine (2) was used without further purification.

General procedure for the synthesis of 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylic acids (4).

[0326] A solution of the appropriate 3-formyl-cinnamic acid (3, 0.57 mmol) and the appropriate *N*¹-substituted-phenyl-1,2-diamine (2, 0.57 mmol) in EtOH (5.0 mL) was refluxed for 24-48 hrs. The reaction was cooled, evaporated to dryness and purified via flash chromatography to yield the desired 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylic acid (4).

General procedure for the synthesis of *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamides (5).

[0327] To a solution of the appropriate 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylic acid (4, 0.25 mmol), and HOBt (0.38 mmol) in DMF (5.0 mL) was added EDCI (0.38mmol), *O*-(tetrahydro-pyran-2-yl)-hydroxylamine (0.38 mmol), and DIEA (0.75 mmol). The reaction was stirred at ambient temperature for 18 hrs. The resulting mixture was poured into H₂O, extracted with EtOAc, washed with brine, and dried over MgSO₄. The organic layer was evaporated to dryness and the resulting material was reconstituted in MeOH (2 mL). CSA (0.28 mmol) was added to the solution. The reaction was stirred for 2 hr at ambient temperature and, without further work-up, purified by preparative LCMS to yield the desired *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamide (5).

General procedure for the synthesis of *N*-(2-amino-phenyl)-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamide (6).

[0328] To a solution of the appropriate 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylic acid (5, 0.25 mmol), and HOBt (0.38 mmol) in DMF (5.0 mL) was added EDCI (0.38mmol), 1,2-phenylenediamine (0.38 mmol), and DIEA (0.75 mmol). The reaction was stirred at ambient temperature for 18 hrs. The resulting mixture was poured into H₂O, extracted with EtOAc, washed with brine, and dried over MgSO₄. The organic layer was evaporated to dryness, and the resulting material was reconstituted in MeOH (2 mL) and purified by preparative LCMS to yield the desired *N*-(2-amino-phenyl)-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamide (6).

General procedure for the synthesis of *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-propionamides (7).

[0329] To a solution of the appropriate *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamide (6; 0.65 mmol) in MeOH (1.0 mL) was added Pd/C (10%; 2.5 mg). H₂(g) was bubbled through the stirring reaction for 1 hr. The reaction was filtered through Celite and purified via preparative LCMS to provide the desired *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-propionamide (7).


General procedure for the synthesis of 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylic acids (9).

[0330] A solution of the appropriate 3-formyl-cinnamic acid methyl ester (8, 0.75 mmol) and the appropriate N¹-substituted-phenyl-1,2-diamine (2, 0.75 mmol) in ethanol (2.0 mL) was heated at 80 °C for 24-48 hrs. The reaction was cooled, evaporated to dryness and purified via flash chromatography to yield the desired acrylic acid esters. To a solution of the appropriate acrylic acid methyl ester (0.50 mmol) in MeOH (1.0 mL) was added LiOH (1.0 mmol). The reaction was stirred at ambient temperature for 2 hrs and, poured into H₂O, and acidified to pH = 2 with HCl (6N). The resulting heterogeneous mixture was then extracted with EtOAc. The organic layers were combined, washed with brine, dried over MgSO₄, filtered, and evaporated to dryness to provide the appropriate 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylic acid (9) which were used in subsequent reactions without further purification.

General procedure for the synthesis of 3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-*N*-hydroxyacrylamides (10).

[0331] The procedure for the synthesis of *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamides (5) was used.

Scheme 3:



General procedure for the synthesis of 3-[3-(1H-benzoimidazol-2-yl)-phenyl]-*N*-(tetrahydro-pyran-2-yloxy)-acrylamides (11).

[0332] A solution of the appropriate 3-formyl-cinnamic acid (3, 0.57 mmol) and the appropriate substituted-phenyl-1,2-diamine (0.57 mmol) in EtOH (5.0 mL) was refluxed for 24 hrs. The reaction was cooled, evaporated to dryness, and the resulting acrylic acid was used without further purification. To a solution of the acrylic acid (0.25 mmol) and HOBt (0.38 mmol) in DMF (5.0 mL) was added EDCI (0.38 mmol), *O*-(tetrahydro-pyran-2-yl)-hydroxylamine (0.38 mmol), and DIEA (0.75 mmol). The reaction was stirred at ambient temperature for 18 hrs. The resulting mixture was poured into H₂O, extracted with EtOAc, washed with brine, dried over MgSO₄ and concentrated to dryness. The resulting material was purified via flash chromatography to yield the desired 3-[3-(1H-benzoimidazol-2-yl)-phenyl]-*N*-(tetrahydro-pyran-2-yloxy)-acrylamide (11).

General procedure for the synthesis of 3-[3-(1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide (12)

[0333] To a solution of the appropriate 3-[3-(1H-benzoimidazol-2-yl)-phenyl]-*N*-(tetrahydro-pyran-2-yloxy)-acrylamide (11, 0.25 mmol) in MeOH (2 mL) was added CSA (0.28 mmol). The reaction was stirred for 2 hr at ambient temperature and, without further work-up, purified by preparative LCMS to yield the desired 3-[3-(1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide (12).

General procedure for the synthesis of *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamides (13).


[0334] To a solution of NaH (20 mg, 0.83 mmol) in anhydrous DMF (1 mL) was added a solution of 3-[3-(1*H*-benzoimidazol-2-yl)-phenyl]-*N*-(tetrahydro-pyran-2-yloxy)-acrylamide (11, 0.28 mmol) in anhydrous DMF drop wise at ambient temperature. The reaction was stirred for 30 min and then the appropriate alkyl bromide (0.31 mmol) was added. The reaction was stirred for 30 min, quenched with MeOH (200 μ L) and partitioned between H₂O and EtOAc. The organic layer was washed with brine, dried with MgSO₄, and evaporated to dryness. The resulting material was reconstituted in MeOH (2 mL). CSA (0.31 mmol) was added to the solution. The reaction was stirred for 2 hr at ambient temperature and, without further work-up, purified by preparative LCMS to yield the desired *N*-hydroxy-3-[3-(1-substituted-1*H*-benzoimidazol-2-yl)-phenyl]-acrylamide (13).

[0335] As can be seen from the above reaction schemes, a wide variety of different HDAC inhibitors can be synthesized by these reaction schemes. It is noted that the invention is not intended to be limited to the particular compounds provided in this example. Rather, a wide variety of other compounds according to the present invention having HDAC inhibitory activity may be synthesized by the reaction schemes provided as well as other reaction schemes that may be devised by one of ordinary skill in the art in view of the present teachings.

2. Examples of Inhibitors According to the Present Invention

[0336] Provided in this example are particular compounds that have been found to have HDAC8 activity based on the assay provided in Example 2. It is noted that these compounds may also have activity relative to other HDACs. It is also noted that these compounds are intended to illustrate various HDAC inhibitors according to the present invention and the present invention is not intended to be limited to these compounds:

COMPOUND 1.




3-[3-(1*H*-Benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 6.69 (d, 1H), 7.45 (m, 3H), 7.82 (m, 5H), 8.14 (d,

1H), 9.30 (s, 1H), 10.90 (s, 1H). ESI-MS: m/z 280.2 (M + H)⁺.


COMPOUND 2.



N-Hydroxy-3-[3-(1-methyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 3.93 (s, 3H), 6.60 (d, 1H), 7.33 (band, 2H), 7.60-7.81 (band, 7H), 7.85 (m, 1H), 8.05 (m, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 294.3 (M + H)⁺.


COMPOUND 3.



N-Hydroxy-3-[3-(1-isopropyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide.


¹H NMR (400 MHz, DMSO-*d*₆): δ 1.58 (d, 6H), 4.82 (m, 1H), 6.57 (d, 6H), 7.15-7.68 (m, 8H), 8.12 (s, 1H), 9.06 (s, 1H), 10.78 (s, 1H). ESI-MS: m/z 322.3 (M + H).

COMPOUND 4.




(±)-3-[3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl]-N-hydroxy-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 1.21 (t, 3H), 1.76 (m, 1H), 2.23 (m, 1H), 3.23 (m, 4H), 3.52 (m, 1H), 3.77 (m, 2H), 4.79 (m, 1H), 6.60 (d, 1H), 7.41 (m, 2H), 7.54-7.91 (band, 6H), 8.14 (m, 1H), 9.68 (s, 1H). ESI-MS: m/z 391.2 (M + H)⁺.

COMPOUND 5.

N-Hydroxy-3-{3-[1-(1-methyl-piperidin-4-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.


^1H NMR (400MHz, DMSO-*d*₆): δ 2.27 (m, 2H), 2.80 (m, 5H), 3.20 (m, 2H), 3.57 (m, 2H), 4.70 (m, 1H), 6.60 (d, 1H), 7.45 (m, 2H), 7.60-7.95 (band, 6H), 8.14 (m, 1H), 10.07 (s, 1H). ESI-MS: m/z 377.1 (M + H)⁺.

COMPOUND 6.

N-Hydroxy-3-[3-(1-piperidin-4-ylmethyl)-1H-benzoimidazol-2-yl]-phenyl-acrylamide.

The product is obtained in salt form with two 10-camphorsulfonic acid (CSA).

^1H NMR (400MHz, DMSO-*d*₆): δ 0.71 (s, 6H), 1.02 (s, 6H), 1.23 (m, 2H), 1.24 (q, 4H), 1.65 (m, 2H), 1.77 (m, 4H), 1.92 (t, 2H), 2.14 (m, 1H), 2.22 (m, 2H), 2.35 (d, 2H), 2.62 (m, 2H), 2.65 (t, 2H), 2.85 (d, 2H), 3.12 (m, 2H), 4.47 (d, 2H), 6.64 (d, 1H), 7.40-8.06 (band, 11H), 10.8 (s, 1H). ESI-MS: m/z 377.1 (M + H)⁺.

COMPOUND 7.


N-Hydroxy-3-{3-[1-(2-morpholin-4-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

The product is obtained in salt form with two 10-camphorsulfonic acid (CSA).

^1H NMR (400 MHz, DMSO-*d*₆): δ 0.71 (s, 6H), 1.02 (s, 6H), 1.24 (q, 4H), 1.78 (m, 4H), 1.92 (t, 2H), 2.22 (m, 2H), 2.35 (d, 2H), 2.65 (t, 2H), 2.81 (d, 2H), 3.14-3.95 (band, 8H),

6.62 (d, 1H), 7.51-8.12 (band, 9H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: m/z 393.3 (M + H)⁺.


COMPOUND 8.



N-Hydroxy-3-[3-(1-phenyl-1H-benzimidazol-2-yl)-phenyl]-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 6.43 (d, 1H), 6.38-7.83 (band, 14H), 9.16 (s, 1H), 10.7 (s, 1H). ESI-MS: m/z 356.4 (M + H)⁺.


COMPOUND 9.



3-[3-[1-(4-Chloro-phenyl)-1H-benzimidazol-2-yl]-phenyl]-N-hydroxy-acrylamide.


¹H NMR (400 MHz, DMSO-*d*₆): δ 6.45 (d, 1H), 7.22-7.44 (band, 7H), 7.52 (d, 2H), 7.62 (d, 1H), 7.66 (d, 2H), 7.82 (d, 1H), 7.89 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 390.1 (M + H)⁺.

COMPOUND 10.




N-Hydroxy-3-[3-[1-(4-methoxy-phenyl)-1H-benzimidazol-2-yl]-phenyl]-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 3.81 (s, 3H), 6.46 (d, 1H), 7.12 (d, 2H), 7.17 (d, 1H), 7.31 (m, 2H), 7.35-7.44 (band, 5H), 7.59 (d, 1H), 7.79 (d, 1H), 7.89 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 386.2 (M + H)⁺.


COMPOUND 11.N-Hydroxy-3-[3-[1-(4-phenoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 6.46 (d, 1H), 7.10 (m, 3H), 7.26 (d, 1H), 7.34 (m, 3H), 7.40-7.50 (band, 8H), 7.63 (d, 1H), 7.79 (d, 1H), 7.82 (s, 1H), 9.12 (s, 1H), 10.80 (s, 1H).

ESI-MS: m/z 448.2 ($M + H$)⁺.

COMPOUND 12.3-[3-[1-(4-Benzylloxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl]-N-hydroxy-acrylamide.


^1H NMR (400 MHz, DMSO-*d*₆): δ 5.15 (s, 2H), 6.47 (d, 1H), 7.17 (d, 1H), 7.19 (d, 2H), 7.25-7.68 (band, 13H), 7.80 (d, 1H), 7.90 (s, 1H), 9.10 (s, 1H), 10.81 (s, 1H). ESI-MS: m/z 462.2 ($M + H$)⁺.

COMPOUND 13.3-[3-(1-Benzyl-1H-benzoimidazol-2-yl)-phenyl]-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 5.70 (s, 2H), 6.54 (d, 1H), 7.08 (d, 2H), 7.29 (m,

3H), 7.49 (m, 2H), 7.65 (t, 1H), 7.72 (d, 1H), 7.76 (d, 1H), 7.84 (d, 2H), 8.01 (s, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: m/z 370.2 ($M + H$)⁺.


COMPOUND 14.



3-{3-[1-(2-Chloro-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 5.60 (s, 2H), 6.51 (d, 1H), 6.65 (d, 1H), 7.20-7.40 (band, 4H), 7.5-7.65 (band, 4H), 7.71 (d, 1H), 7.81 (d, 1H), 7.91 (s, 1H), 9.10 (br s, 1H), 10.88 (br s, 1H). ESI-MS: m/z 404.2 ($M + H$)⁺.


COMPOUND 15.



3-{3-[1-(3-Chloro-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.


¹H NMR (400 MHz, DMSO-*d*₆): δ 5.60 (s, 2H), 6.54 (d, 1H), 6.87 (m, 1H), 7.09 (s, 1H), 7.28 (m, 3H), 7.56 (m, 3H), 7.67 (d, 1H), 7.76 (m, 2H), 7.94 (d, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: m/z 404.2 ($M + H$)⁺.

COMPOUND 16.




3-{3-[1-(4-Chloro-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 5.60 (s, 2H), 6.54 (d, 1H), 7.10 (d, 2H), 7.27 (m, 2H), 7.34 (d, 2H), 7.53 (m, 2H), 7.57 (d, 1H), 7.66 (d, 1H), 7.74 (d, 2H), 7.95 (s, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: m/z 404.2 ($M + H$)⁺.

COMPOUND 17.


N-Hydroxy-3-{3-[1-(2-trifluoromethoxy-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 5.65 (s, 2H), 6.52 (d, 1H), 6.72 (d, 1H), 7.26 (m, 3H), 7.38-7.55 (band, 4H), 7.58 (d, 1H), 7.71 (d, 1H), 7.78 (d, 2H), 7.92 (s, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: m/z 454.2 (M + H)⁺.

COMPOUND 18.


N-Hydroxy-3-{3-[1-(3-trifluoromethoxy-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 5.71 (s, 2H), 6.55 (d, 1H), 6.99 (m, 2H), 7.24 (d, 2H), 7.30 (d, 1H), 7.53 (t, 1H), 7.65 (m, 3H), 7.68 (d, 1H), 7.76 (m, 2H), 7.92 (s, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: m/z 454.2 (M + H)⁺.

COMPOUND 19.


N-Hydroxy-3-{3-[1-(4-pyrazol-1-yl-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 5.66 (s, 2H), 6.56 (m, 2H), 7.13 (d, 2H), 7.28 (m, 2H), 7.55 (m, 3H), 7.68-7.76 (band, 6H), 8.00 (s, 1H), 8.42 (d, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: m/z 436.2 (M + H)⁺.

COMPOUND 20.


(R)-N-Hydroxy-3-{3-[1-(1-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400MHz, DMSO-*d*6): δ 1.97 (d, 3H), 5.85 (q, 1H), 6.56 (d, 1H), 7.07-7.32 (band, 8H), 7.52-7.72 (band, 5H), 7.92 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 384.1 (M + H) $^+$.

COMPOUND 21.


(S)-N-Hydroxy-3-{3-[1-(1-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400MHz, DMSO-*d*6): δ 2.00 (d, 3H), 5.85 (q, 1H), 6.56 (d, 1H), 7.07-7.32 (band, 8H), 7.52-7.72 (band, 5H), 7.92 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 384.1 (M + H) $^+$.

COMPOUND 22.


(R)-N-Hydroxy-3-{3-[1-(1-p-tolyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*6): δ 1.95 (d, 3H), 2.22 (s, 3H), 5.80 (q, 1H), 6.59 (d, 1H), 7.05-7.22 (band, 7H), 7.52-7.73 (band, 4H), 7.78 (d, 1H), 7.90 (s, 1H), 9.15 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 398.2 (M + H) $^+$.

COMPOUND 23.


(R)-3-(3-{1-[1-(4-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-N-hydroxyacrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.05 (d, 3H), 5.85 (q, 1H), 6.59 (d, 1H), 6.90-7.29 (band, 8H), 7.50-7.8 (band, 4H), 7.90 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 402.2 (M + H)⁺.

COMPOUND 24.


N-Hydroxy-3-[3-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.99 (t, 2H), 4.53 (t, 2H), 6.53 (d, 1H), 6.83 (m, 2H), 7.11 (t, 3H), 7.29 (m, 2H), 7.49-7.74 (band, 7H), 9.11 (s, 1H), 10.8 (s, 1H). ESI-MS: *m/z* 384.4 (M + H)⁺.

COMPOUND 25.


3-(3-{1-[2-(3-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-N-hydroxyacrylamide

^1H NMR (400MHz, DMSO-*d*₆): δ 3.00 (m, 2H), 4.57 (m, 2H), 6.55 (m, 2H), 6.67 (d, 1H), 6.91 (m, 1H), 7.13 (q, 1H), 7.29 (m, 2H), 7.55 (m, 3H), 7.62 (s, 1H), 7.67 (m, 3H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 402.1 (M + H)⁺.

COMPOUND 26.


3-(3-{1-[2-(4-Fluoro-phenyl)-ethyl]-1H-benzimidazol-2-yl}-phenyl)-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.96 (t, 2H), 4.55 (t, 2H), 6.53 (d, 1H), 6.83-7.73 (band, 13H), 9.11 (s, 1H), 10.78 (s, 1H). ESI-MS: *m/z* 402.2 (M + H)⁺.

COMPOUND 27.

N-Hydroxy-3-(3-{1-[2-(2-methoxy-phenyl)-ethyl]-1H-benzimidazol-2-yl}-phenyl)-acrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 2.95 (m, 2H), 3.48 (s, 3H), 4.53 (m, 2H), 6.52 (d, 1H), 6.69 (m, 3H), 7.12 (m, 1H), 7.30 (m, 2H), 7.55 (m, 4H), 7.70 (m, 3H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 414.17 (M + H)⁺.


COMPOUND 28.

N-Hydroxy-3-(3-{1-[2-(3-methoxy-phenyl)-ethyl]-1H-benzimidazol-2-yl}-phenyl)-acrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 2.95 (m, 2H), 3.55 (s, 3H), 4.54 (m, 2H), 6.35 (m, 2H), 6.52 (d, 1H), 6.65 (m, 1H), 7.0 (m, 1H), 7.27 (m, 2H), 7.45-7.80 (band, 7H), 9.10 (s, 1H),

10.80 (s, 1H). ESI-MS: m/z 414.1 ($M + H$)⁺.

COMPOUND 29.




N-Hydroxy-3-(3-{1-[2-(4-methoxy-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 2.95 (m, 2H), 3.65 (s, 3H), 4.50 (m, 2H), 6.55 (d, 4H), 6.65-6.80 (band, 4H), 7.37 (m, 2H), 7.45-7.80 (band, 7H), 9.10 (s, 1H), 10.80 (s, 1H).

ESI-MS: m/z 414.1 ($M + H$)⁺.

COMPOUND 30.




3-(3-{1-[2-(2-Chloro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-N-hydroxyacrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 3.08 (m, 2H), 4.59 (m, 2H), 6.52 (d, 1H), 6.79 (d, 1H), 7.11 (m, 3H), 7.31 (m, 2H), 7.54 (m, 4H), 7.71 (m, 3H), 9.10 (s, 1H), 10.80 (s, 1H).

ESI-MS: m/z 418.1 ($M + H$)⁺.


COMPOUND 31.



3-(3-{1-[2-(3-Chloro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-N-hydroxyacrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.95 (t, 2H), 4.56 (t, 2H), 6.52 (d, 1H), 6.88 (s, 1H), 7.07-7.74 (band, 11H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: *m/z* 418.3 (M + H).


COMPOUND 32.



3-(3-(1-(2-(4-Chloro-phenyl)-ethyl)-1H-benzoimidazol-2-yl)-phenyl)-N-hydroxy-acrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 2.96 (m, 2H), 4.56 (m, 2H), 6.54 (d, 1H), 6.79 (d, 2H), 7.11 (m, 2H), 7.29 (m, 2H), 7.51 (m, 3H), 7.64-7.74 (band, 4H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 418.1 (M + H)⁺.


COMPOUND 33.



(±)-3-(3-(1-(2-(4-Fluoro-phenyl)-1-methyl-ethyl)-1H-benzoimidazol-2-yl)-phenyl)-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 1.68 (d, 3H), 3.11 (m, 2H), 4.68 (m, 1H), 6.49 (d, 1H), 6.58 (t, 2H), 6.84 (t, 2H), 7.20 (d, 1H), 7.32 (s, 1H), 7.45-7.75 (band, 5H), 9.10 (s, 1H), 10.81 (s, 1H). ESI-MS: *m/z* 416.2 (M + H)⁺.

COMPOUND 34.




(R)-N-Hydroxy-3-(3-(1-(2-phenyl-propyl)-1H-benzoimidazol-2-yl)-phenyl)-

acrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 1.10 (m, 3H), 3.10 (m, 1H), 4.47 (m, 2H), 6.56 (d, 1H), 6.75 (d, 2H), 7.10 (m, 3H), 7.30 (m, 2H), 7.50-7.80 (band, 7H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 398.2 ($M + H$) $^+$.

COMPOUND 35.




(S)-N-Hydroxy-3-{3-[1-(2-phenyl-propyl)-1H-benzoimidazol-2-yl]-phenyl}-

acrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 1.10 (m, 3H), 3.10 (m, 1H), 4.47 (m, 2H), 6.56 (d, 1H), 6.75 (d, 2H), 7.10 (m, 3H), 7.3 (m, 2H), 7.50-7.80 (band, 7H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 398.2 ($M + H$) $^+$.


COMPOUND 36.



(S)-N-Hydroxy-3-{3-[1-(1-hydroxymethyl-2-phenylethyl)-1H-benzoimidazol-2-yl]-


phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 3.08 (m, 1H), 4.05 (m, 1H), 4.40 (m, 2H), 5.24 (t, 1H), 6.43 (d, 1H), 6.55 (d, 2H), 7.00 (t, 2H), 7.10 (m, 1H), 7.20-8.00 (band, 7H), 9.11 (s, 1H), 10.81 (s, 1H). ESI-MS: m/z 414.4 ($M + H$) $^+$.

COMPOUND 37.


(R)-N-Hydroxy-3-{3-[1-(1-hydroxymethyl-2-phenylethyl)-1H-benzimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 3.09 (m, 1H), 4.06 (m, 1H), 4.40 (m, 2H), 5.24 (t, 1H), 6.43 (d, 1H), 6.55 (d, 2H), 7.00 (t, 2H), 7.10 (m, 1H), 7.20-8.00 (band, 7H), 9.11 (s, 1H), 10.81 (s, 1H). ESI-MS: m/z 414.4 ($M + H$) $^+$.

COMPOUND 38.

(R)-N-Hydroxy-3-{3-[1-(2-hydroxy-2-phenylethyl)-1H-benzimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 4.40 (d, 2H), 5.91 (t, 1H), 5.78 (br., 1H), 6.55 (d, 1H), 7.03-7.29 (band, 5H), 7.39 (t, 2H), 7.57 (m, 2H), 7.70-7.89 (band, 5H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 400.2 ($M + H$) $^+$.


COMPOUND 39.

N-Hydroxy-3-{3-[1-(2-pyridin-2-ylethyl)-1H-benzimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 3.14 (t, 2H), 4.69 (t, 2H), 6.54 (d, 1H), 6.80 (d,

1H), 7.17 (t, 1H), 7.26 (m, 2H), 7.52-7.94 (band, 7H), 8.38 (d, 1H), 9.09 (s, 1H), 10.78 (s, 1H). ESI-MS: m/z 385.1 (M + H)⁺.


COMPOUND 40.



N-Hydroxy-3-{3-[1-(2-pyridin-3-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 3.20 (m, 2H), 4.80 (m, 2H), 6.55 (d, 1H), 7.46-7.90 (band, 11H), 7.95 (m, 1H), 8.30 (m, 1H), 8.54 (d, 1H). ESI-MS: m/z 385.1 (M + H)⁺.


COMPOUND 41.



N-Hydroxy-3-{3-[1-(2-pyridin-4-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.


¹H NMR (400MHz, DMSO-*d*₆): δ 3.25 (m, 2H), 4.82 (m, 2H), 6.52 (d, 1H), 7.35-7.67 (band, 7H), 7.76-7.82 (m, 3H), 7.95 (m, 1H), 8.54 (d, 2H). ESI-MS: m/z 385.1 (M + H)⁺.

COMPOUND 42.




N-Hydroxy-3-(3-{1-[2-(1H-indol-3-yl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide.


¹H NMR (400 MHz, DMSO-*d*₆): δ 3.09 (t, 2H), 4.53 (t, 2H), 6.52 (d, 1H), 6.80 (m, 2H), 6.91 (s, 1H), 7.25-8.13 (m, 14H), 9.11 (s, 1H), 10.77 (s, 1H), 12.7 (s, 1H). ESI-MS: m/z

423.1 (M + H)⁺.COMPOUND 43.N-Hydroxy-3-[3-(1-indan-2-yl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide


¹H NMR (400MHz, DMSO-*d*₆): δ 3.56 (m, 4H), 5.51 (m, 1H), 6.62(d, 1H), 7.07-7.33 (band, 7H), 7.54-7.77 (band, 5H), 7.92 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 396.2 (M + H)⁺.

COMPOUND 44.N-Hydroxy-3-[3-[1-(3-phenyl-propyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide.


¹H NMR (400MHz, DMSO-*d*₆): δ 2.07 (m, 2H), 2.53 (m, 2H), 4.29 (m, 2H), 6.56 (d, 1H), 7.07-7.32 (band, 8H), 7.52-7.72 (band, 5H), 7.92 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 398.2 (M + H)⁺.

COMPOUND 45.3-[3-(5-Fluoro-1H-benzoimidazol-2-yl)-phenyl]-N-hydroxy-acrylamide.


¹H NMR (400 MHz, DMSO-*d*₆): δ 6.61 (d, 1H), 7.19 (m, 1H), 7.52 (m, 2H), 7.70 (m, 3H), 8.13 (d, 1H), 8.38 (s, 1H), 9.40 (s, 1H), 10.90 (s, 1H). ESI-MS: *m/z* 298.2 (M + H)⁺.

COMPOUND 46.N-Hydroxy-3-[3-(6-propylsulfanyl-1H-benzimidazol-2-yl)-phenyl]-acrylamide.


^1H NMR (400 MHz, DMSO-*d*₆): δ 0.92 (t, 3H), 1.53 (m, 2H), 2.95 (t, 2H), 6.58 (d, 1H), 7.31 (dd, 1H), 7.50 (d, 1H), 7.62 (m, 3H), 7.73 (d, 1H), 8.08 (d, 1H), 8.38 (s, 1H), 9.40 (s, 1H), 10.90 (s, 1H). ESI-MS: m/z 354.2 ($M + H$)⁺.

COMPOUND 47.N-Hydroxy-3-[3-(7-methyl-1H-benzimidazol-2-yl)-phenyl]-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.63 (s, 3H), 6.58 (d, 1H), 7.27 (t, 1H), 7.50 (d, 1H), 7.53 (s, 2H), 7.65 (t, 1H), 7.75 (d, 1H), 8.14 (d, 1H), 8.38 (s, 1H), 9.40 (s, 1H), 10.90 (s, 1H). ESI-MS: m/z 294.2 ($M + H$)⁺.

COMPOUND 48.2-[3-(2-Hydroxycarbamoyl-vinyl)-phenyl]-3H-benzimidazole-5-carboxylic acid methyl ester.


^1H NMR (400 MHz, DMSO-*d*₆): δ 3.91 (s, 3H), 6.62 (d, 1H), 7.54 (d, 1H), 7.62 (t, 1H), 7.74 (m, 2H), 7.89 (d, 1H), 8.14 (d, 1H), 8.23 (s, 1H), 8.40 (s, 1H), 9.40 (s, 1H), 10.90 (s, 1H). ESI-MS: m/z 338.2 ($M + H$)⁺.

COMPOUND 49.3-[3-(6,7-Difluoro-1H-benzimidazol-2-yl)-phenyl]-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 6.58 (d, 1H), 7.21 (m, 1H), 7.35 (m, 1H), 7.48 (d,

1H), 7.54 (t, 1H), 7.65 (d, 1H), 8.09 (d, 1H), 8.38 (s, 1H), 9.40 (s, 1H), 10.90 (s, 1H). ESI-MS: m/z 316.2 (M + H)⁺.


COMPOUND 50.



3-{3-[1-(4-Methoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid.

¹H NMR (400 MHz, DMSO-*d*₆): δ 3.72 (s, 3H), 6.23 (d, 1H), 7.06 (d, 2H), 7.10 (d, 1H), 7.23 (m, 2H), 7.29-7.39 (band, 3H), 7.42 (s, 1H), 7.49 (d, 1H), 7.62 (d, 1H), 7.69 (s, 1H), 7.73 (d, 1H). ESI-MS: m/z 371.2 (M + H)⁺.


COMPOUND 51.



3-{3-[1-(4-Benzoyloxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid.

¹H NMR (400 MHz, DMSO-*d*₆): δ 5.10 (s, 2H), 6.23 (d, 1H), 7.10 (d, 1H), 7.14 (m, 2H), 7.18-7.38 (band, 8H), 7.43 (m, 3H), 7.52 (d, 1H), 7.64 (d, 1H), 7.78 (s, 1H), 7.73 (d, 1H). ESI-MS: m/z 447.2 (M + H)⁺.


COMPOUND 52.



3-{3-[1-(4-Chloro-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid.

^1H NMR (400 MHz, DMSO-*d*₆): δ 6.35 (d, 1H), 7.25 (d, 1H), 7.33 (m, 2H), 7.41-7.53 (band, 5H), 7.66 (d, 2H), 7.72 (d, 1H), 7.77 (s, 1H), 7.82 (d, 1H). ESI-MS: *m/z* 375.2 (M + H)⁺.


COMPOUND 53.



(R)-3-(3-{1-[1-(4-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylic acid.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.05 (d, 3H), 5.89 (q, 1H), 6.65 (d, 1H), 7.15-7.33 (band, 7H), 7.68 (m, 2H), 7.76 (d, 2H), 7.96 (d, 1H), 7.98 (s, 1H). ESI-MS: *m/z* 387.2 (M + H)⁺.


COMPOUND 54.



(R)-3-{3-[1-(1-p-Tolyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid.


^1H NMR (400 MHz, DMSO-*d*₆): δ 1.95 (d, 3H), 2.23 (s, 3H), 5.80 (q, 1H), 6.55 (d, 1H), 7.04-7.27 (band, 7H), 7.63 (m, 2H), 7.71 (m, 2H), 7.90 (m, 2H). ESI-MS: *m/z* 383.2 (M + H)⁺.

COMPOUND 55.




3-{3-[1-(4-Phenoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid.

^1H NMR (400 MHz, DMSO-*d*₆): δ 6.25 (d, 1H), 7.05 (d, 2H), 7.12 (m, 2H), 7.18 (m, 2H), 7.25 (m, 2H), 7.35-7.42 (band, 5H), 7.46 (d, 1H), 7.64 (m, 3H), 7.76 (d, 1H). ESI-MS: *m/z* 433.2 (M + H)⁺.

COMPOUND 56.


N-(2-Amino-phenyl)-3-{3-[1-(4-methoxy-phenyl)-1H-benzimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 3.81 (s, 3H), 5.01 (br s, 2H), 6.58 (t, 1H), 6.76 (d, 1H), 6.93 (t, 1H), 6.97 (d, 1H), 7.13 (d, 2H), 7.16 (d, 1H), 7.26-7.43 (band, 6H), 7.50 (d, 1H), 7.63 (d, 1H), 7.80 (d, 1H), 8.05 (s, 1H), 8.21 (s, 1H) 9.42 (s, 1H). ESI-MS: m/z 461.2 ($M + H$) $^+$.

COMPOUND 57.


N-(2-Amino-phenyl)-3-{3-[1-(4-benzyloxy-phenyl)-1H-benzimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 5.00 (br s, 2H), 5.15 (s, 2H), 6.58 (t, 1H), 6.76 (d, 1H), 6.93 (t, 1H), 6.98 (d, 1H), 7.17 (d, 1H), 7.21 (d, 2H), 7.26-7.45 (band, 9H), 7.50 (m, 3H), 7.65 (d, 1H), 7.81 (d, 1H), 8.05 (s, 1H), 9.50 (s, 1H). ESI-MS: m/z 537.2 ($M + H$) $^+$.

COMPOUND 58.


N-(2-Amino-phenyl)-3-{3-[1-(4-chloro-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 5.01 (br s, 2H), 6.52 (t, 1H), 6.69 (d, 1H), 6.85 (t, 1H), 6.94 (d, 1H), 7.19 (d, 1H), 7.22-7.39 (band, 6H), 7.46 (m, 3H), 7.60 (m, 3H), 7.76 (d, 1H), 7.96 (s, 1H), 9.38 (s, 1H). ESI-MS: m/z 465.2 ($M + H$)⁺.

COMPOUND 59.


(R)-N-(2-Amino-phenyl)-3-(3-{1-[1-(4-fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 2.00 (d, 3H), 5.00 (br s, 2H), 5.89 (q, 1H), 6.59 (t, 1H), 6.77 (d, 1H), 6.93 (t, 1H), 7.04 (d, 1H), 7.10-7.28 (band, 7H), 7.38 (d, 1H), 7.62-7.74 (band, 4H), 7.84 (d, 1H), 8.01 (s, 1H), 9.45 (s, 1H). ESI-MS: m/z 477.2 ($M + H$)⁺.

COMPOUND 60.


(R)-N-(2-Amino-phenyl)-3-{3-[1-(1-p-tolyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 1.94 (d, 3H), 2.23 (s, 3H), 4.98 (br s, 2H), 5.84 (q, 1H), 6.57 (t, 1H), 6.76 (d, 1H), 6.92 (t, 1H), 7.00-7.22 (band, 8H), 7.36 (d, 1H), 7.60-7.71 (band, 4H), 7.82 (d, 1H), 7.97 (s, 1H), 9.40 (s, 1H). ESI-MS: m/z 473.2 ($M + H$)⁺.

COMPOUND 61.


N-(2-Amino-phenyl)-3-{3-[1-(4-phenoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 5.00 (br s, 2H), 6.59 (t, 1H), 6.76 (d, 1H), 6.93 (t, 1H), 6.96 (d, 1H), 7.11 (d, 2H), 7.19 (m, 3H), 7.23-7.39 (band, 4H), 7.41-7.55 (band, 7H), 7.67 (t, 1H), 7.93 (s, 1H), 8.05 (s, 1H), 8.21 (s, 1H), 9.45 (s, 1H). ESI-MS: *m/z* 523.2 (*M* + *H*)⁺.

COMPOUND 62.

N-Hydroxy-3-[4-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 3.00 (t, 2H), 4.62 (t, 2H), 6.60 (d, 1H), 6.92 (m, 2H), 7.15 (m, 3H), 7.44 (m, 2H), 7.55 (d, 1H), 7.62 (d, 2H), 7.75 (m, 3H), 7.88 (d, 1H), 9.10 (s, 1H), 10.88 (s, 1H). ESI-MS: *m/z* 384.2 (*M* + *H*)⁺.


COMPOUND 63.

N-(2-Amino-phenyl)-3-[4-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide.

^1H NMR (400 MHz, DMSO-*d*₆): δ 3.00 (t, 2H), 4.65 (t, 2H), 6.71 (t, 1H), 6.83 (d, 1H), 6.92 (m, 2H), 6.99 (t, 1H), 7.11 (d, 1H), 7.16 (m, 3H), 7.43 (m, 3H), 7.65 (m, 3H), 7.78

(m, 3H), 7.88 (d, 1H), 9.78 (s, 1H). ESI-MS: m/z 459.2 (M + H)⁺.


COMPOUND 64.



3-[4-(1-Phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylic acid.

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.90 (t, 2H), 4.45 (t, 2H), 6.56 (d, 1H), 6.84 (m, 2H), 7.06 (m, 3H), 7.20 (m, 2H), 7.52 (m, 3H), 7.61 (m, 2H), 7.71 (d, 2H). ESI-MS: m/z 369.2 (M + H)⁺.


COMPOUND 65.



N-Hydroxy-3-[3-[1-(2-piperidin-1-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.19-1.29 (m, 6H), 2.10-2.20 (m, 4H), 2.83 (t, 2H), 4.49 (t, 2H), 6.56 (d, 1H), 7.22-7.30 (band, 2H), 7.56-7.82 (band, 6H), 7.96 (s, 1H), 9.09 (br., s, 1H), 10.86 (br., s, 1H). ESI-MS: m/z 391.4 (M + H)⁺.

COMPOUND 66.




N-Hydroxy-3-[3-[1-(trans-4-hydroxy-cyclohexyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.20 (m, 2H), 2.95 (m, 4H), 2.48 (m, 2H), 3.64 (m,

1H), 4.42 (m, 1H), 4.62 (d, 1H), 6.54 (d, 1H), 7.23 (m, 2H), 7.56-7.86 (band, 6H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: m/z 378.2 (M + H)⁺.


COMPOUND 67.



3-[3-(1-tert-Butyl-1H-benzimidazol-2-yl)-phenyl]-N-hydroxy-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.58 (d, 9H), 6.53 (d, 1H), 7.25-7.96 (band, 8H), 9.06 (s, 1H), 10.78 (s, 1H). ESI-MS: m/z 336.3 (M + H)⁺.


COMPOUND 68.



(±)-3-[3-(1-Cyclohexyl-1H-benzimidazol-2-yl)-phenyl]-N-hydroxy-acrylamide


¹H NMR (400 MHz, DMSO-*d*₆): δ 0.88 (m, 2H), 1.28 (m, 4H), 1.57 (m, 4H), 4.14 (m, 1H), 6.53 (d, 1H), 7.25-7.91 (band, 9H), 9.06 (s, 1H), 10.71 (s, 1H). ESI-MS: m/z 362.2 (M + H)⁺.

COMPOUND 69.




3-[2-[3-(2-Hydroxycarbamoyl-vinyl)-phenyl]-benzimidazol-1-yl]-piperidine-1-carboxylic acid tert-butyl ester

¹H NMR (400MHz, DMSO-*d*₆): δ 1.24-1.42 (band, 11H), 1.78 (d, 1H), 2.09 (m, 1H), 2.56 (m, 1H), 2.97 (m, 1H), 3.65 (m, 1H), 3.89 (m, 1H), 4.27 (m, 1H), 6.59 (d, 1H), 7.29 (m, 2H), 7.56-7.99 (band, 7H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: m/z 463.2 (M + H)⁺.

COMPOUND 70.


(±)-N-Hydroxy-3-[3-(1-piperidin-3-yl)-1H-benzimidazol-2-yl]-phenyl]-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 0.82 (m, 1H), 1.24 (m, 1H), 1.44 (m, 1H), 1.76 (m, 1H), 2.06 (m, 1H), 2.45 (m, 1H), 2.67 (m, 1H), 2.89 (m, 1H), 3.14 (m, 1H), 3.14 (m, 1H), 4.37 (m, 1H), 6.57-6.61 (d, 1H), 7.26 (m, 2H), 7.56-7.99 (band, 7H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: *m/z* 363.2 (M + H)⁺.

COMPOUND 71.


3-[3-[1-(2-Diethylamino-ethyl)-1H-benzimidazol-2-yl]-phenyl]-N-hydroxy-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 0.63 (t, 6H), 2.25 (q, 4H), 2.62 (t, 2H), 4.36 (t, 1H), 6.59 (d, 1H), 7.28 (m, 2H), 7.55-8.10 (band, 7H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: *m/z* 379.2 (M + H)⁺.

COMPOUND 72.


(±)-N-Hydroxy-3-[3-[1-(1-methyl-piperidin-3-yl)-1H-benzimidazol-2-yl]-phenyl]-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.50 (m, 1H), 1.78 (d, 1H), 2.02 (m, 1H), 2.25 (m, 5H), 2.74 (m, 2H), 2.94 (m, 1H), 4.42 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.56-7.88 (band, 7H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: *m/z* 376.19 (M + H)⁺.

COMPOUND 73.


(±)-N-Hydroxy-3-{3-[1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 0.95 (m, 6H), 1.45 (m, 1H), 1.78 (d, 1H), 2.02 (m, 1H), 2.26-2.31 (m, 2H), 2.68-2.76 (m, 2H), 2.94-2.96 (d, 2H), 4.35-4.37 (m, 1H), 6.57-6.61 (d, 1H), 7.23-7.30 (m, 2H), 7.55-7.89 (band, 7H), 9.09 (s, 1H), 10.86 (s, 1H). ESI-MS: *m/z* 405.2 (M + H)⁺.

COMPOUND 74.


(±)-N-Hydroxy-3-(3-{1-[1-(2-hydroxy-ethyl)-piperidin-3-yl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide.

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.49-1.55 (m, 1H), 1.78 (d, 1H), 2.02-2.05 (m, 1H), 2.12-2.18 (t, 1H), 2.28-2.33 (m, 1H), 2.45 (t, 2H), 2.83-2.89 (t, 2H), 3.02 (m, 1H), 3.45-3.49 (m, 2H), 4.40 (m, 1H), 6.57-6.61 (d, 1H), 7.23-7.29 (m, 2H), 7.55-7.71 (band, 4H), 7.78 (m, 1H), 7.86-7.89 (m, 2H). ESI-MS: *m/z* 407.2 (M + H)⁺.

COMPOUND 75.


(±)-3-{3-[1-(1-Ethyl-pyrrolidin-2-ylmethyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.69 (t, 3H), 1.29 (m, 1H), 1.44 (m, 3H), 2.00 (m, 2H), 2.22 (m, 1H), 2.85 (m, 2H), 4.31 (m, 2H), 6.59 (d, 1H), 7.27 (m, 2H), 7.55-7.75 (band, 5H), 7.90 (d, 1H), 8.12 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 391.1 ($M + H$) $^+$.

COMPOUND 76.


(±)-3-{3-[1-(1-Ethyl-pyrrolidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.10 (t, 3H), 2.33 (m, 4H), 2.63 (m, 2H), 3.19 (m, 2H), 5.10 (m, 1H), 6.59 (d, 1H), 7.27 (m, 2H), 7.55-7.71 (band, 4H), 7.77 (d, 1H), 7.85 (s, 1H), 8.23-8.25 (d, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 377.1 ($M + H$) $^+$.

COMPOUND 77.


(±)-3-{3-[1-(2-Dimethylamino-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 2.00 (s, 6H), 2.55 (t, 2H), 4.38 (t, 2H), 6.59 (d, 1H), 7.27 (m, 2H), 7.55-7.80 (band, 6H), 7.99 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 351.1 ($\text{M} + \text{H}$) $^+$.

COMPOUND 78.


(±)-3-{3-[1-(2-Dimethylamino-1-methyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 1.62 (d, 3H), 1.87 (s, 6H), 2.50 (m, 1H), 3.05 (m, 1H), 4.60 (m, 1H), 6.57 (d, 1H), 7.24 (m, 2H), 7.54-7.82 (band, 6H), 7.86 (s, 1H), 9.11 (s, 1H), 10.79 (s, 1H). ESI-MS: m/z 365.1 ($\text{M} + \text{H}$) $^+$.

COMPOUND 79.


(±)-3-{3-[1-(2-Diisopropylamino-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxyacrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.62 (d, 12H), 2.57 (t, 2H), 2.75 (m, 2H), 4.28 (t, 2H), 6.59 (d, 1H), 7.28 (m, 2H), 7.55-7.85 (band, 6H), 8.00 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 407.1 ($M + H$) $^+$.

COMPOUND 80.


(±)-3-{3-[1-(1-Aza-bicyclo[2.2.2]oct-2-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxyacrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.53 (m, 1H), 1.64 (m, 2H), 2.06 (m, 1H), 2.30 (m, 1H), 2.73 (m, 2H), 2.90 (m, 1H), 3.20 (m, 1H), 3.35 (m, 1H), 3.60 (m, 1H), 4.60 (m, 1H), 6.57 (d, 1H), 7.30 (m, 2H), 7.53-7.90 (band, 7H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 389.1 ($M + H$) $^+$.

COMPOUND 81.


(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-6,7-difluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.96 (t, 3H), 1.47 (m, 1H), 1.79 (m, 1H), 1.96 (m, 4H), 2.37 (m, 2H), 2.83 (m, 1H), 3.07 (m, 1H), 4.39 (m, 1H), 6.59 (d, 1H), 7.39 (m, 1H), 7.58 (m, 2H), 7.70 (d, 2H), 7.81 (m, 1H), 7.92 (s, 1H), 9.10 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 427.1 ($M + H$) $^+$.

COMPOUND 82.


(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-5,6-difluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide-trifluoroacetic acid.

^1H NMR (400 MHz, DMSO- d_6): δ 1.18 (t, 3H), 1.60 (m, 1H), 2.14 (m, 2H), 3.22 (m, 3H), 3.55 (m, 1H), 3.82 (m, 3H), 4.72 (m, 1H), 6.59 (d, 1H), 7.61 (m, 3H), 7.85 (m, 3H), 8.27 (m, 1H), 9.40 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 427.1 ($M + H$) $^+$.

COMPOUND 83.


(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4,6-difluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.98 (t, 3H), 1.42 (m, 1H), 1.78 (m, 1H), 2.10 (m, 2H), 2.35 (m, 3H), 2.71 (m, 2H), 3.00 (m, 1H), 4.36 (m, 1H), 6.59 (d, 1H), 7.18 (m, 1H), 7.57 (d, 1H), 7.69 (m, 2H), 7.81 (m, 1H), 7.85 (s, 1H), 7.92 (s, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 427.1 ($M + H$) $^+$.

COMPOUND 84.


(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-5,6,7-trifluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.95 (t, 3H), 1.47 (m, 1H), 1.78 (m, 1H), 2.03 (m, 3H), 2.38 (m, 3H), 2.85 (m, 1H), 3.10 (m, 1H), 4.42 (m, 1H), 6.59 (d, 1H), 7.62 (d, 1H), 7.66 (m, 2H), 7.74 (m, 1H), 7.84 (m, 2H), 9.11 (s, 1H), 10.81 (s, 1H). ESI-MS: m/z 445.1 ($M + H$) $^+$.

COMPOUND 85.


(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4,6,7-trifluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.95 (t, 3H), 1.46 (m, 1H), 1.78 (m, 1H), 2.03 (m, 3H), 2.37 (m, 3H), 2.85 (m, 1H), 3.10 (m, 1H), 4.42 (m, 1H), 6.64 (d, 1H), 7.54 (m, 2H), 7.67 (m, 2H), 7.83 (m, 1H), 7.88 (s, 1H), 9.12 (s, 1H), 10.81 (s, 1H). ESI-MS: m/z 445.1 ($M + H$) $^+$.

COMPOUND 86.

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4,5,6,7-tetrafluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.


^1H NMR (400 MHz, DMSO- d_6): δ 0.95 (t, 3H), 1.46 (m, 1H), 1.79 (m, 1H), 2.04 (m, 3H), 2.37 (m, 3H), 2.85 (m, 1H), 3.17 (m, 1H), 4.45 (m, 1H), 6.60 (d, 1H), 7.58 (d, 1H), 7.67 (m, 2H), 7.85 (m, 1H), 7.88 (s, 1H), 9.12 (s, 1H), 10.81 (s, 1H). ESI-MS: m/z 463.1 ($M + H$) $^+$.

COMPOUND 87.

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4-fluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.96 (t, 3H), 1.48 (m, 1H), 1.79 (m, 1H), 2.05 (m, 2H), 2.32 (m, 3H), 2.71 (m, 1H), 2.84 (m, 1H), 3.04 (m, 1H), 4.38 (m, 1H), 6.60 (d, 1H), 7.08 (m, 1H), 7.27 (m, 1H), 7.57 (d, 1H), 7.65 (m, 2H), 7.73 (d, 1H), 7.80 (m, 1H), 7.87 (s, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 409.1 ($M + H$) $^+$.


COMPOUND 88.



(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-5-fluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.


^1H NMR (400 MHz, DMSO- d_6): δ 0.96 (t, 3H), 1.48 (m, 1H), 1.79 (m, 1H), 2.05 (m, 2H), 2.32 (m, 3H), 2.71 (m, 1H), 2.84 (m, 1H), 3.03 (m, 1H), 4.38 (m, 1H), 6.59 (d, 1H), 7.14 (m, 1H), 7.49 (m, 1H), 7.57 (d, 1H), 7.65 (m, 2H), 7.79 (m, 1H), 7.89 (s, 1H), 7.90-7.93 (m, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 409.1 ($M + H$) $^+$.

COMPOUND 89.




(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-6-fluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.96 (t, 3H), 1.47 (m, 1H), 1.78 (m, 1H), 2.11 (m, 2H), 2.32 (m, 3H), 2.71 (m, 1H), 2.83 (m, 1H), 3.00 (m, 1H), 4.36 (m, 1H), 6.59 (d, 1H), 7.12 (m, 1H), 7.56 (d, 1H), 7.63 (m, 2H), 7.70 (m, 1H), 7.77 (m, 2H), 7.84 (s, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 409.1 ($M + H$) $^+$.

COMPOUND 90.


(R)-N-Hydroxy-3-{3-[1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazole-2-yl]-phenyl}-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.95 (d, 6H), 1.45 (m, 1H), 1.79 (m, 1H), 2.01 (m, 1H), 2.29 (m, 2H), 2.74 (m, 2H), 3.04 (m, 2H), 4.35 (m, 1H), 6.59 (d, 1H), 7.25 (m, 2H), 7.56 (d, 1H), 7.63 (m, 2H), 7.69 (m, 1H), 7.79 (m, 1H), 7.84 (s, 1H), 7.88 (d, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 405.2 ($M + H$) $^+$.

COMPOUND 91.


(R)-3-{3-[4-Fluoro-1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.95 (d, 6H), 1.45 (m, 1H), 1.79 (m, 1H), 2.02 (m, 1H), 2.27 (m, 2H), 2.73 (m, 2H), 2.98 (m, 2H), 4.35 (m, 1H), 6.60 (d, 1H), 7.08 (m, 1H), 7.26 (m, 1H), 7.57 (d, 1H), 7.65 (m, 2H), 7.74 (d, 1H), 7.80 (m, 1H), 7.86 (s, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 423.1 ($M + H$) $^+$.

COMPOUND 92.


(R)-3-{3-[5-Fluoro-1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.95 (d, 6H), 1.44 (m, 1H), 1.79 (m, 1H), 2.04 (m, 1H), 2.27 (m, 2H), 2.73 (m, 2H), 2.98 (m, 2H), 4.35 (m, 1H), 6.59 (d, 1H), 7.14 (m, 1H), 7.50 (m, 1H), 7.56 (d, 1H), 7.64 (m, 2H), 7.79 (m, 1H), 7.84 (s, 1H), 7.92 (m, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 423.1 ($M + H$) $^+$.

COMPOUND 93.


(R)-3-{3-[4-Fluoro-1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.49 (m, 1H), 1.78 (m, 1H), 2.06 (m, 2H), 2.24 (m, 4H), 2.72 (m, 2H), 2.95 (m, 1H), 4.42 (m, 1H), 6.60 (d, 1H), 7.08 (m, 1H), 7.26 (m, 1H), 7.58 (d, 1H), 7.65 (m, 2H), 7.73 (d, 1H), 7.80 (m, 1H), 7.88 (s, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 395.1 ($M + H$) $^+$.

COMPOUND 94.


(R)-3-{3-[5-Fluoro-1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.49 (m, 1H), 1.77 (m, 1H), 2.03 (m, 2H), 2.24 (m, 4H), 2.72 (m, 2H), 2.99 (m, 1H), 4.43 (m, 1H), 6.59 (d, 1H), 7.14 (m, 1H), 7.50 (m, 1H), 7.57 (d, 1H), 7.64 (m, 2H), 7.78 (m, 1H), 7.85 (s, 1H), 7.91 (m, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 395.1 ($M + H$) $^+$.

COMPOUND 95.


(R)-3-{5-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-2-fluoro-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 0.96 (t, 3H), 1.51 (m, 1H), 1.79 (m, 1H), 2.04 (m, 2H), 2.34 (m, 3H), 2.75 (m, 1H), 2.86 (m, 1H), 3.03 (m, 1H), 4.37 (m, 1H), 6.70 (d, 1H), 7.26 (m, 2H), 7.48 (m, 1H), 7.58 (d, 1H), 7.71 (m, 2H), 7.87 (m, 1H), 7.95 (d, 1H), 9.15 (s, 1H), 10.90 (s, 1H). ESI-MS: m/z 409.1 ($M + H$) $^+$.

COMPOUND 96.


(R)-3-[3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-4-fluoro-phenyl]-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.01 (t, 3H), 1.48 (m, 1H), 1.79 (m, 1H), 1.99 (m, 2H), 2.32 (m, 3H), 2.71 (m, 1H), 2.85 (m, 1H), 2.98 (m, 1H), 4.12 (m, 1H), 6.53 (d, 1H), 7.30 (m, 2H), 7.53 (m, 2H), 7.72 (d, 1H), 7.90 (m, 3H), 9.10 (s, 1H), 10.77 (s, 1H). ESI-MS: m/z 409.1 ($M + H$) $^+$.

COMPOUND 97.


(R)-3-[4-Fluoro-3-[5-fluoro-1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl]-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.47 (m, 1H), 1.75 (m, 1H), 1.90 (m, 1H), 2.05 (m, 1H), 2.19 (m, 4H), 2.69 (m, 2H), 2.87 (m, 1H), 4.11 (m, 1H), 6.53 (d, 1H), 7.19 (m, 1H), 7.54 (m, 3H), 7.87 (m, 2H), 7.96 (m, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 413.1 ($M + H$) $^+$.

COMPOUND 98.


(R)-3-(3-{1-[1-(2,2-Difluoro-ethyl)-piperidin-3-yl]-4-fluoro-1H-benzoimidazol-2-yl}-phenyl)-N-hydroxy-acrylamide.

^1H NMR (400 MHz, DMSO- d_6): δ 1.49 (m, 1H), 1.78 (m, 1H), 2.05 (m, 1H), 2.32 (m, 2H), 2.83 (m, 3H), 3.08 (m, 2H), 4.40 (m, 1H), 6.13 (m, 1H), 6.60 (d, 1H), 7.08 (m, 1H), 7.26 (m, 1H), 7.58 (d, 1H), 7.66 (m, 2H), 7.75 (d, 1H), 7.81 (m, 1H), 7.87 (s, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 445.1 ($M + H$) $^+$.

COMPOUND 99.


(±)-3-{3-[1-(2-Dimethylamino-2-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 1.92 (s, 6H), 3.46 (t, 1H), 4.52 (q, 1), 5.02 (q, 1H), 6.55 (d, 1H), 6.83 (d, 2H), 7.05-7.28 (band, 5H), 7.52-7.72 (band, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 427.2 ($M + H$) $^+$.

COMPOUND 100.


(±)-3-{3-[1-(1-Benzyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 1.50 (m, 1H), 1.78 (d, 1H), 2.03 (br s, 2H), 2.14 (m, 1H), 2.33 (m, 1H), 2.82 (m, 2H), 2.95 (m, 1H), 3.55 (m, 1H), 4.40 (m, 1H), 6.60 (d, 1H), 7.23-7.34 (band, 7H), 7.53-7.69 (band, 4H), 7.75-7.87 (band, 3H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 453.2 ($M + H$) $^+$.

COMPOUND 101.


(±)-N-Hydroxy-3-{3-[1-(1-propyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 0.79 (t, 3H), 1.45 (m, 3H), 1.78 (d, 1H), 2.04 (m, 2H), 2.29 (m, 3H), 2.78 (m, 2H), 3.02 (m, 1H), 4.40 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.55-7.71 (band, 4H), 7.78 (m, 1H), 7.86 (m, 2H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 405.2 ($M + H$) $^+$.

COMPOUND 102.


(±)-N-Hydroxy-3-{3-[1-(1-isobutyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 0.82 (d d, 6H), 1.46 (m, 1H), 1.77 (m, 2H), 1.97 (m, 1H), 2.09 (m, 3H), 2.30 (m, 1H), 2.73 (m, 2H), 3.04 (d, 1H), 4.38 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.55-7.89 (band, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 419.2 (M + H)⁺.

COMPOUND 103.


(±)-3-{3-[1-(1-Allyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxyacrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 1.49 (m, 1H), 1.79 (d, 1H), 2.09 (m, 2H), 2.30 (m, 1H), 2.80 (m, 2H), 2.97 (m, 3H), 4.40 (m, 1H), 5.15 (M, 2H), 5.79 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.56-7.71 (band, 4H), 7.82 (m, 3H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 403.2 (M + H)⁺.

COMPOUND 104.


(±)-3-{3-[1-(1-Acetyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxyacrylamide.

^1H NMR (400MHz, CDCl_3 -*d*): δ 1.60 (m, 1H), 1.94 (m, 2H), 2.29 (s, 3H), 2.73 (m, 1H), 3.28 (t, 1H), 3.60 (t, 1H), 3.99 (d, 1H), 4.57 (m, 1H), 5.39 (d, 1H), 6.88 (d, 1H), 7.34 (m, 2H), 7.51-7.69 (band, 5H), 7.89 (m, 2H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 405.1 (*M* + *H*) $^+$.

COMPOUND 105.


(S)-N-Hydroxy-3-{3-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400MHz, $\text{DMSO-}d_6$): δ 1.50 (m, 1H), 1.78 (d, 1H), 2.02 (m, 1H), 2.24 (m, 5H), 2.74 (br s, 2H), 2.94 (br s, 1H), 4.42 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.56-7.88 (band, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: *m/z* 377.1 (*M* + *H*) $^+$.

COMPOUND 106.


(S)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxyacrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 0.96 (t, 3H), 1.50 (m, 1H), 1.78 (d, 1H), 2.05 (m, 2H), 2.33 (m, 3H), 2.75 (t, 1H), 2.84 (d, 1H), 3.02 (m, 1H), 4.40 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.51-7.88 (band, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 391.2 ($M + H$) $^+$.

COMPOUND 107.

(R)-N-Hydroxy-3-{3-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}acrylamide.


^1H NMR (400MHz, DMSO- d_6): δ 1.50 (m, 1H), 1.78 (d, 1H), 2.02 (m, 2H), 2.21-2.33 (m, 4H), 2.75 (m, 2H), 2.94 (m, 1H), 4.42 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.56-7.88 (band, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 377.1 ($M + H$) $^+$.

COMPOUND 108.

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxyacrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 0.96 (t, 3H), 1.50 (m, 1H), 1.78 (d, 1H), 2.05 (m, 2H), 2.33 (m, 3H), 2.75 (t, 1H), 2.84 (d, 1H), 3.02 (d, 1H), 4.40 (m, 1H), 6.59 (d, 1H), 7.26 (m, 2H), 7.55-7.88 (band, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 391.2 (M + H)⁺.


COMPOUND 109.



(R)-N-Hydroxy-3-{3-[3-(1-isopropyl-piperidin-3-yl)-3H-imidazo[4,5-b]pyridin-2-yl]-phenyl}-acrylamide.


^1H NMR (400MHz, DMSO-*d*₆): δ 0.95 (br s, 6H), 1.43 (m, 1H), 1.78 (m, 1H), 1.99 (m, 1H), 2.19 (m, 1H), 2.75 (m, 3H), 2.92 (m, 1H), 3.25 (m, 1H), 4.35 (m, 1H), 6.58 (d, 1H), 7.31 (m, 1H), 7.57 (d, 1H), 7.65 (m, 2H), 7.81 (d, 1H), 7.89 (s, 1H), 8.11 (d, 1H), 8.41 (d, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 406.2 (M + H)⁺.

COMPOUND 110.




(S)-N-Hydroxy-3-{3-[3-(1-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide.

^1H NMR (400MHz, DMSO-*d*₆): δ 1.70 (m, 1H), 1.90 (m, 1H), 2.20 (m, 1H), 2.49 (m, 3H), 3.30 (m, 2H), 4.70 (m, 1H), 6.60 (d, 1H), 7.31 (m, 2H), 7.51-7.90 (m, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 363.2 (M + H)⁺.

COMPOUND 111.

(R)-N-Hydroxy-3-[3-(1-piperidin-3-yl)-1H-benzimidazol-2-yl]-phenyl-acrylamide.


^1H NMR (400MHz, DMSO- d_6): δ 1.39 (m, 1H), 1.70 (m, 1H), 2.00 (m, 1H), 2.45 (m, 3H), 2.80 (m, 1H), 3.05 (m, 1H), 4.30 (m, 1H), 6.60 (d, 1H), 7.31 (m, 2H), 7.51-7.90 (m, 7H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 363.2 ($M + H$) $^+$.

COMPOUND 112.

(±)-N-Hydroxy-3-{3-[1-(1-propyl-piperidin-3-yl)-1H-benzimidazol-2-yl]-phenyl}-

propionamide.

^1H NMR (400MHz, DMSO- d_6): δ 0.98 (t, 3H), 1.71 (m, 3H), 2.01 (m, 2H), 2.31 (m, 1H), 2.97 (m, 2H), 3.20 (m, 1H), 3.49 (m, 1H), 3.98 (m, 2H), 4.68 (m, 1H), 7.35 (m, 2H), 7.48 (d, 1H), 7.54 (m, 3H), 7.74 (m, 1H), 8.08 (m, 1H), 9.11 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 407.2 ($M + H$) $^+$.


COMPOUND 113.

(±)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzimidazol-2-yl]-phenyl}-N-hydroxy-

propionamide.

^1H NMR (400MHz, DMSO- d_6): δ 1.29 (t, 3H), 1.75 (m, 1H), 1.98 (m, 2H), 2.31 (m, 2H), 2.48 (m, 2H), 2.95 (m, 2H), 3.20 (m, 2H), 3.42 (m, 1H), 3.82 (m, 1H), 3.95 (m, 1H), 4.68 (m, 1H), 7.35 (m, 2H), 7.48 (d, 1H), 7.54 (m, 3H), 7.74 (m, 1H), 8.08 (m, 1H), 9.68 (s, 1H), 10.80 (s, 1H). ESI-MS: m/z 393.2 ($M + H$) $^+$.


COMPOUND 114.



N-Hydroxy-3-[3-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-propionamide.


^1H NMR (400MHz, DMSO- d_6): δ 2.28 (t, 2H), 2.97 (t, 2H), 3.05 (t, 2H), 4.62 (t, 2H), 6.85 (m, 2H), 7.15 (m, 3H), 7.31 (s, 1H), 7.38 (m, 1H), 7.48 (m, 4H), 7.77 (m, 1H), 7.96 (d, 1H), 8.70 (s, 1H), 10.45 (s, 1H). ESI-MS: m/z 386.2 ($M + H$) $^+$.

COMPOUND 115.




N-Hydroxy-3-[3-[3-(1-methyl-piperidin-3-yl)-3H-imidazo[4,5-b]pyridin-2-yl]-phenyl]-acrylamide.

^1H NMR (400MHz, DMSO- d_6): δ 1.49 (m, 1H), 1.76 (m, 1H), 1.95 (m, 2H), 2.20 (s, 3H), 2.70 (m, 2H), 2.93 (m, 1H), 3.09 (m, 1H), 4.41 (m, 1H), 6.58 (d, 1H), 7.31 (m, 1H), 7.56-7.60 (d, 1H), 7.66 (m, 2H), 7.81 (d, 1H), 7.90 (s, 1H), 8.11 (d, 1H), 8.41 (d, 1H). ESI-MS: m/z 377.19 ($M + H$) $^+$.

COMPOUND 116.


(R)-3-{3-[3-(1-Ethyl-piperidin-3-yl)-3H-imidazo[4,5-b]pyridin-2-yl]-phenyl}-N-hydroxy-acrylamide.

ESI-MS: m/z 392.2 (M + H)⁺.

COMPOUND 117.


N-Hydroxy-3-[5-(1-phenethyl-1H-benzoimidazol-2-yl)-pyridin-3-yl]-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 2.97 (t, 2H), 4.61 (t, 2H), 6.65 (m, 3H), 7.02 (m, 3H), 7.32 (d t, 2H), 7.52 (d, 1H), 7.80 (m, 3H), 8.61 (s, 1H), 8.83 (s, 1H). ESI-MS: m/z 385.2 (M + H)⁺.

COMPOUND 118.

(R)-N-Hydroxy-3-[5-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-pyridin-3-yl]-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 1.56 (m, 1H), 1.77 (m, 1H), 2.05 (m, 2H), 2.27 (m, 4H), 2.75 (m, 2H), 3.02 (m, 1H), 4.35 (m, 1H), 6.70 (d, 1H), 7.31 (m, 2H), 7.61 (d, 1H), 7.72 (m, 1H), 7.90 (d, 1H), 8.29 (s, 1H), 8.83 (s, 1H), 8.98 (s, 1H). ESI-MS: m/z 378.2 (M + H)⁺.

COMPOUND 119.

(R)-N-Hydroxy-3-{5-[3-(1-methyl-piperidin-3-yl)-3H-imidazo[4,5-b]pyridin-2-yl]}-pyridin-3-yl}-acrylamide.

¹H NMR (400MHz, DMSO-*d*₆): δ 1.51 (m, 1H), 1.92 (m, 3H), 2.15 (s, 3H), 2.71 (m, 2H), 3.02 (m, 2H), 4.33 (m, 1H), 6.78 (br d, 1H), 7.33 (m, 2H), 8.13 (d, 1H), 8.27 (s, 1H), 8.42 (d, 1H), 8.79 (s, 1H), 8.95 (s, 1H). ESI-MS: *m/z* 379.2 (M + H)⁺.

[0337] As used herein the symbols and conventions used in these processes, schemes and examples are consistent with those used in the contemporary scientific literature, for example, the Journal of the American Chemical Society or the Journal of Biological Chemistry. Standard single-letter or three-letter abbreviations are generally used to designate amino acid residues, which are assumed to be in the L-configuration unless otherwise noted. Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification. Specifically, the following abbreviations may be used in the examples and throughout the specification:

g (grams);	mg (milligrams);
L (liters);	mL (milliliters);
μL (microliters);	psi (pounds per square inch);
M (molar);	mM (millimolar);
i.v. (intravenous);	Hz (Hertz);
MHz (megahertz);	mol (moles);
mmol (millimoles);	RT (ambient temperature);
min (minutes);h (hours);	
mp (melting point);	TLC (thin layer chromatography);
Tr (retention time);	RP (reverse phase);
MeOH (methanol);	i-PrOH (isopropanol);
TEA (triethylamine);	TFA (trifluoroacetic acid);

TFAA (trifluoroacetic anhydride); THF (tetrahydrofuran);
DMSO (dimethylsulfoxide); EtOAc (ethyl acetate);
DME (1,2-dimethoxyethane); DCM (dichloromethane);
DCE (dichloroethane); DMF (N,N-dimethylformamide);
DMPU (N,N'-dimethylpropyleneurea); CDI (1,1-carbonyldiimidazole);
IBCF (isobutyl chloroformate); HOAc (acetic acid);
HOSu (N-hydroxysuccinimide); HOBT (1-hydroxybenzotriazole);
Et₂O (diethyl ether); EDCI (ethylcarbodiimide hydrochloride);
BOC (tert-butyloxycarbonyl); Fmoc (9-fluorenylmethoxycarbonyl);
DCC (dicyclohexylcarbodiimide); CBZ (benzyloxycarbonyl);
Ac (acetyl); atm (atmosphere);
TMSE (2-(trimethylsilyl)ethyl); TMS (trimethylsilyl);
TIPS (triisopropylsilyl); TBS (t-butyldimethylsilyl);
DMAP (4-dimethylaminopyridine); Me (methyl);
OMe (methoxy); Et (ethyl);
Et (ethyl); tBu (tert-butyl);
HPLC (high pressure liquid chromatography);
BOP (bis(2-oxo-3-oxazolidinyl)phosphinic chloride);
TBAF (tetra-n-butylammonium fluoride);
mCPBA (meta-chloroperbenzoic acid).

[0338] All references to ether or Et₂O are to diethyl ether; brine refers to a saturated aqueous solution of NaCl. Unless otherwise indicated, all temperatures are expressed in °C (degrees Centigrade). All reactions conducted under an inert atmosphere at RT unless otherwise noted.

[0339] ¹H NMR spectra were recorded on a Bruker Avance 400. Chemical shifts are expressed in parts per million (ppm). Coupling constants are in units of hertz (Hz). Splitting patterns describe apparent multiplicities and are designated as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), br (broad).

[0340] Low-resolution mass spectra (MS) and compound purity data were acquired on a Waters ZQ LC/MS single quadrupole system equipped with electrospray ionization (ESI) source, UV detector (220 and 254 nm), and evaporative light scattering detector (ELSD).

Thin-layer chromatography was performed on 0.25 mm E. Merck silica gel plates (60F-254), visualized with UV light, 5% ethanolic phosphomolybdic acid, Ninhydrin or p-anisaldehyde solution. Flash column chromatography was performed on silica gel (230-400 mesh, Merck).

[0341] It will be apparent to those skilled in the art that various modifications and variations can be made to the compounds, compositions, kits, and methods of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents.


What is claimed is:

1. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen, and

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a

metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


2. A compound according to claim 1 wherein the compound comprises the formula



3. A compound of claim 1, wherein Q is a substituted or unsubstituted phenyl ring.
4. A compound of claim 1, wherein Q is a substituted or unsubstituted heteroaryl.
5. A compound of claim 1, wherein Q is a substituted or unsubstituted heteroaryl selected from the group consisting of furan, thiophene, pyrrole, pyrazole, triazole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, benzofuran, isobenzofuran, benzothiophene, isobenzothiophene, indole, isobenzazole, quinoline, isoquinoline, cinnoline, quinazoline, naphthyridine, pyridopyridine, quinoxaline, phthalazine, benthiazole, and triazine.
6. A compound according to claim 1, wherein at least one X in the six membered ring is a substituted carbon atom.
7. A compound according to claim 1, wherein at least one of the X substituents in the six membered ring is -CF.
8. A compound according to claim 1, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino,


alkylamino, or an alkoxy group.

9. A compound according to claim 1, wherein M is selected from the group consisting of:




10. A compound according to claim 1, wherein M comprises a hydroxamic acid moiety.

11. A compound according to claim 1, wherein -Q-L-M is




12. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR_{12} and N ;

each Y is independently selected from the group consisting of O , S and NR_{12} ;

R_1 is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_1 is not halo, cyano, nitro and thio in the case where the ring atom to which R_1 is bound is nitrogen;

R_2, R_3, R_4 and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted; and

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;


Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a


metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

13. A compound according to claim 12, wherein the compound comprises the formula




14. A compound according to claim 12, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
15. A compound according to claim 12, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
16. A compound according to claim 12, wherein M is selected from the group consisting of:




17. A compound according to claim 12, wherein M comprises a hydroxamic acid

moiety.

18. A compound according to claim 12, wherein -Q-L-M is



19. A compound comprising a formula selected from the group consisting of:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;


R_6 , R_7 , R_8 , and R_9 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound.

20. A compound according to claim 19, wherein the compound comprises the formula




21. A compound according to claim 19, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.
22. A compound according to claim 19, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
23. A compound according to claim 19, wherein M is selected from the group

consisting of:




24. A compound according to claim 19, wherein M comprises a hydroxamic acid moiety.

25. A compound according to claim 19, wherein -L-M is



26. A compound comprising a formula selected from the group consisting of:



wherein

each X is independently selected from the group consisting of CR_{12} and N ;

each Y is independently selected from the group consisting of O , S and NR_{12} ;

R_1 is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_1 is not halo, cyano, nitro and thio in the case where the ring atom to which R_1 is bound is nitrogen;

R_6 , R_7 , R_8 , and R_9 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and

thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound.

27. A compound according to claim 26, wherein the compound comprises the formula




28. A compound according to claim 26, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.

29. A compound according to claim 26, wherein M is selected from the group consisting of:





30. A compound according to claim 26, wherein M comprises a hydroxamic acid moiety.

31. A compound according to claim 26, wherein -L-M is



32. A compound according to claim 26, wherein at least one of R₆, R₇, R₈, and R₉ is fluorine.
33. A compound according to claim 26, wherein at least one X in the six membered ring is a substituted carbon atom.
34. A compound according to claim 26, wherein at least one of the X substituents in the six membered ring is -CF.
35. A compound comprising a formula selected from the group consisting of:





wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;


R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


36. A compound according to claim 35, wherein the compound comprises the formula




37. A compound according to claim 35, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
38. A compound according to claim 35, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.
39. A compound according to claim 35, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
40. A compound according to claim 35, wherein M is selected from the group consisting of:



41. A compound according to claim 35, wherein M comprises a hydroxamic acid moiety.
42. A compound according to claim 35, wherein $-\text{L}-\text{M}$ is




43. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁ is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁ is not halo, cyano, nitro and thio in the case where the ring atom to which R₁ is bound is nitrogen;

R₁₀ and R₁₁ are taken together to form a substituted or unsubstituted aromatic ring;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


44. A compound according to claim 43, wherein the substituted or unsubstituted aromatic ring formed when R₁₀ and R₁₁ are taken together is selected from the group consisting of substituted or unsubstituted aryl and heteroaryl.
45. A compound according to claim 43, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
46. A compound according to claim 43, wherein M is selected from the group consisting of:



47. A compound according to claim 43, wherein M comprises a hydroxamic acid moiety.
48. A compound according to claim 43, wherein -L-M is



49. A compound comprising the formula



wherein

R_1 is selected from the group consisting of hydrogen, halo, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_1 is not halo, cyano, nitro and thio in the case where the ring atom to which R_1 is bound is nitrogen;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the remainder of the compound.

50. A compound according to claim 49, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.


51. A compound according to claim 49, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides

(-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.

52. A compound according to claim 49, wherein M is selected from the group consisting of:



53. A compound according to claim 49, wherein M comprises a hydroxamic acid moiety.
54. A compound according to claim 49, wherein -L-M is




55. A compound comprising the formula:



wherein

Z-Q- is selected from the group consisting of




R_1 is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the remainder of the compound.

56. A compound according to claim 55, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
57. A compound according to claim 55, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
58. A compound according to claim 55, wherein M is selected from the group consisting of:



59. A compound according to claim 55, wherein M comprises a hydroxamic acid moiety.

60. A compound according to claim 55, wherein -L-M is




61. A compound comprising the formula:



wherein

Z-Q- is selected from the group consisting of




R₁ is selected from the group consisting of (C₁₋₄)alkyl, phenyl, 1-piperidin-4-ylmethyl, 2-morpholin-4-yl-ethyl, 2-halo-phenyl, 2-halo-phen(C₁₋₄)alkyl, 3-halo-phen(C₁₋₄)alkyl, 2-CF₃O-phen(C₁₋₄)alkyl, 3-CF₃O-phen(C₁₋₄)alkyl, 3-halo-phenyl, 4-halo-phenyl, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 4-phenoxy-phenyl, 4-benzyloxyphenyl, 4-pyrazol-1-yl-

benzyl, 1-p-tolyl-ethyl, pyrrolidin-3-yl, 1-(C₁₋₄)alkyl-pyrrolidin-2-yl, 1-(C₁₋₄)alkyl-pyrrolidin-2-yl; 2-di(C₁₋₄)alkylamino-ethyl, 2-di(C₁₋₄)alkylamino-1-methyl-ethyl, 2-di(C₁₋₄)alkylamino-ethyl, 2-hydroxy-2-phenyl-ethyl, 2-pyridin-2-yl-ethyl, 2-pyridin-3-yl-ethyl, 2-pyridin-4-yl-ethyl, 2-(1H-indol-3-yl)-ethyl, 3-indolyl(C₁₋₄)alkyl, 1-indan-2-yl, *R*- α -(HOCH₂)-phen(C₁₋₄)alkyl, *S*- α -(HOCH₂)-phen(C₁₋₄)alkyl, *S*- β -(HOCH₂)-phen(C₁₋₄)alkyl, *R*- β -(CH₃)-phen(C₁₋₄)alkyl, 6-propylsulfanyl, *trans*-4-hydroxy-cyclohexyl, 1-aza-bicyclo[2.2.2]oct-2-yl, 1-(C₁₋₄)alkyl-piperidin-3-yl, 1-(2,2-difluoro-ethyl)-piperidin-3-yl, (2-di(C₁₋₄)alkylamino-2-phenyl-ethyl), 1-benzyl-piperidin-3-yl, 1-allyl-piperidin-3-yl, 1-acetyl-piperidin-3-yl, piperidin-3-yl, and phen(C₁₋₄)alkyl;


R₂, R₃, R₄, and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, cyano, and nitro;

M is selected from the group consisting of:




and L is E, Z or mixtures of E/Z -CH₂=CH₂-.

62. A compound according to claim 61 wherein at least one of R₂, R₃, R₄, or R₅ is fluorine.
63. A compound according to claim 61, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
64. A compound according to claim 61, wherein M is selected from the group consisting of:



65. A compound according to claim 61, wherein M comprises a hydroxamic acid moiety.

66. A compound according to claim 61, wherein -L-M is



67. A compound selected from the group consisting of:

3-[3-(1H-Benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide;

N-Hydroxy-3-[3-(1-methyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

N-Hydroxy-3-[3-(1-isopropyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

(±)-3-[3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl]-*N*-hydroxy-acrylamide;

N-Hydroxy-3-[3-[1-(1-methyl-piperidin-4-yl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide;

N-Hydroxy-3-[3-(1-piperidin-4-ylmethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

N-Hydroxy-3-[3-[1-(2-morpholin-4-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide;

N-Hydroxy-3-[3-(1-phenyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

3-[3-[1-(4-Chloro-phenyl)-1H-benzoimidazol-2-yl]-phenyl]-*N*-hydroxy-acrylamide;

N-Hydroxy-3-[3-[1-(4-methoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide;

N-Hydroxy-3-[3-[1-(4-phenoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide;

3-{3-[1-(4-Benzyloxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

3-[3-(1-Benzyl-1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide;

3-{3-[1-(2-Chloro-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

3-{3-[1-(3-Chloro-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

3-{3-[1-(4-Chloro-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

N-Hydroxy-3-{3-[1-(2-trifluoromethoxy-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

N-Hydroxy-3-{3-[1-(3-trifluoromethoxy-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

N-Hydroxy-3-{3-[1-(4-pyrazol-1-yl-benzyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[1-(1-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*S*)-*N*-Hydroxy-3-{3-[1-(1-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[1-(1-*p*-tolyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*R*)-3-(3-{1-[1-(4-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-acrylamide;

N-Hydroxy-3-[3-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

3-(3-{1-[2-(3-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-acrylamide;

3-(3-{1-[2-(4-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-acrylamide;

N-Hydroxy-3-(3-{1-[2-(2-methoxy-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide;

N-Hydroxy-3-(3-{1-[2-(3-methoxy-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide;

N-Hydroxy-3-(3-{1-[2-(4-methoxy-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide;

3-(3-{1-[2-(2-Chloro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-acrylamide;

3-(3-{1-[2-(3-Chloro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-

acrylamide;

3-(3-{1-[2-(4-Chloro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-acrylamide;

(±)-3-(3-{1-[2-(4-Fluoro-phenyl)-1-methyl-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-*N*-hydroxy-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[1-(2-phenyl-propyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*S*)-*N*-Hydroxy-3-{3-[1-(2-phenyl-propyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*S*)-*N*-Hydroxy-3-{3-[1-(1-hydroxymethyl-2-phenylethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[1-(1-hydroxymethyl-2-phenylethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[1-(2-hydroxy-2-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

N-Hydroxy-3-{3-[1-(2-pyridin-2-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

N-Hydroxy-3-{3-[1-(2-pyridin-3-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

N-Hydroxy-3-{3-[1-(2-pyridin-4-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

N-Hydroxy-3-(3-{1-[2-(1H-indol-3-yl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylamide;

N-Hydroxy-3-[3-(1-indan-2-yl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

N-Hydroxy-3-{3-[1-(3-phenyl-propyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

3-[3-(5-Fluoro-1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide;

N-Hydroxy-3-[3-(6-propylsulfanyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

N-Hydroxy-3-[3-(7-methyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;

2-[3-(2-Hydroxycarbamoyl-vinyl)-phenyl]-3H-benzoimidazole-5-carboxylic acid methyl ester;

3-[3-(6,7-Difluoro-1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide;

3-{3-[1-(4-Methoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid;
3-{3-[1-(4-Benzoyloxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid;
3-{3-[1-(4-Chloro-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid;
(R)-3-(3-{1-[1-(4-Fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-phenyl)-acrylic acid;
(R)-3-(3-{1-(1-p-Tolyl-ethyl)-1H-benzoimidazol-2-yl}-phenyl)-acrylic acid;
3-{3-[1-(4-Phenoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-acrylic acid;
N-(2-Amino-phenyl)-3-{3-[1-(4-methoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
N-(2-Amino-phenyl)-3-{3-[1-(4-benzyloxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
N-(2-Amino-phenyl)-3-{3-[1-(4-chloro-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
(R)-*N*-(2-Amino-phenyl)-3-(3-{1-[1-(4-fluoro-phenyl)-ethyl]-1H-benzoimidazol-2-yl}-
phenyl)-acrylamide;
(R)-*N*-(2-Amino-phenyl)-3-{3-[1-(1-p-tolyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
N-(2-Amino-phenyl)-3-{3-[1-(4-phenoxy-phenyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
N-Hydroxy-3-[4-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;
N-(2-Amino-phenyl)-3-[4-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;
3-[4-(1-Phenethyl-1H-benzoimidazol-2-yl)-phenyl]-acrylic acid;
N-Hydroxy-3-{3-[1-(2-piperidin-1-yl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
N-Hydroxy-3-{3-[1-(*trans*-4-hydroxy-cyclohexyl)-1H-benzoimidazol-2-yl]-phenyl}-
acrylamide;
3-[3-(1-tert-Butyl-1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide;
(±)-3-[3-(1-Cyclohexyl-1H-benzoimidazol-2-yl)-phenyl]-*N*-hydroxy-acrylamide;
3-{2-[3-(2-Hydroxycarbamoyl-vinyl)-phenyl]-benzoimidazol-1-yl}-piperidine-1-
carboxylic acid tert-butyl ester;
(±)-*N*-Hydroxy-3-[3-(1-piperidin-3-yl-1H-benzoimidazol-2-yl)-phenyl]-acrylamide;
3-{3-[1-(2-Diethylamino-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-

acrylamide;

(±)-*N*-Hydroxy-3-{3-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-

acrylamide;

(±)-*N*-Hydroxy-3-{3-[1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-

acrylamide;

(±)-*N*-Hydroxy-3-(3-{1-[1-(2-hydroxy-ethyl)-piperidin-3-yl]-1H-benzoimidazol-2-yl}-

phenyl)-acrylamide;

(±)-3-{3-[1-(1-Ethyl-pyrrolidin-2-ylmethyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-

hydroxy-acrylamide;

(±)-3-{3-[1-(1-Ethyl-pyrrolidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-

acrylamide;

(±)-3-{3-[1-(2-Dimethylamino-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-

acrylamide;

(±)-3-{3-[1-(2-Dimethylamino-1-methyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-

hydroxy-acrylamide;

(±)-3-{3-[1-(2-Diisopropylamino-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-

acrylamide;

(±)-3-{3-[1-(1-Aza-bicyclo[2.2.2]oct-2-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-

hydroxy-acrylamide;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-6,7-difluoro-1H-benzoimidazol-2-yl]-phenyl}-*N*-

hydroxy-acrylamide;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-5,6-difluoro-1H-benzoimidazol-2-yl]-phenyl}-*N*-

hydroxy-acrylamide-trifluoroacetic acid;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4,6-difluoro-1H-benzoimidazol-2-yl]-phenyl}-*N*-

hydroxy-acrylamide;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-5,6,7-trifluoro-1H-benzoimidazol-2-yl]-phenyl}-

N-hydroxy-acrylamide;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4,6,7-trifluoro-1H-benzoimidazol-2-yl]-phenyl}-

N-hydroxy-acrylamide;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4,5,6,7-tetrafluoro-1H-benzoimidazol-2-yl]-

phenyl}-*N*-hydroxy-acrylamide;

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-4-fluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-5-fluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-6-fluoro-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-N-Hydroxy-3-{3-[1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazole-2-yl]-phenyl}-acrylamide;

(R)-3-{3-[4-Fluoro-1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-{3-[5-Fluoro-1-(1-isopropyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-{3-[4-Fluoro-1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-{3-[5-Fluoro-1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-{5-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-2-fluoro-phenyl}-N-hydroxy-acrylamide;

(R)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-4-fluoro-phenyl}-N-hydroxy-acrylamide;

(R)-3-{4-Fluoro-3-[5-fluoro-1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(R)-3-(3-{1-[1-(2,2-Difluoro-ethyl)-piperidin-3-yl]-4-fluoro-1H-benzoimidazol-2-yl}-phenyl)-N-hydroxy-acrylamide;

(±)-3-{3-[1-(2-Dimethylamino-2-phenyl-ethyl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(±)-3-{3-[1-(1-Benzyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-N-hydroxy-acrylamide;

(±)-N-Hydroxy-3-{3-[1-(1-propyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(±)-*N*-Hydroxy-3-{3-[1-(1-isobutyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(±)-3-{3-[1-(1-Allyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

(±)-3-{3-[1-(1-Acetyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

(*S*)-*N*-Hydroxy-3-{3-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*S*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-acrylamide;

(*R*)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

(*R*)-*N*-Hydroxy-3-{3-[3-(1-isopropyl-piperidin-3-yl)-3H-imidazo[4,5-*b*]pyridin-2-yl]-phenyl}-acrylamide;

(*S*)-*N*-Hydroxy-3-[3-(1-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide;

(*R*)-*N*-Hydroxy-3-[3-(1-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl]-acrylamide;

(±)-*N*-Hydroxy-3-{3-[1-(1-propyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-propionamide;

(±)-3-{3-[1-(1-Ethyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-phenyl}-*N*-hydroxy-propionamide;

N-Hydroxy-3-[3-(1-phenethyl-1H-benzoimidazol-2-yl)-phenyl]-propionamide;

N-Hydroxy-3-{3-[3-(1-methyl-piperidin-3-yl)-3H-imidazo[4,5-*b*]pyridin-2-yl]-phenyl}-acrylamide;


(*R*)-3-{3-[3-(1-Ethyl-piperidin-3-yl)-3H-imidazo[4,5-*b*]pyridin-2-yl]-phenyl}-*N*-hydroxy-acrylamide;

N-Hydroxy-3-[5-(1-phenethyl-1H-benzoimidazol-2-yl)-pyridin-3-yl]-acrylamide;

(*R*)-*N*-Hydroxy-3-{5-[1-(1-methyl-piperidin-3-yl)-1H-benzoimidazol-2-yl]-pyridin-3-yl}-acrylamide; and

(*R*)-*N*-Hydroxy-3-{5-[3-(1-methyl-piperidin-3-yl)-3H-imidazo[4,5-*b*]pyridin-2-yl]-pyridin-3-yl}-acrylamide.

68. A compound comprising a formula selected from the group consisting of:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;


each Y is independently selected from the group consisting of O, S and NR_{12} ;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the ring.

69. A compound according to claim 68, wherein the compound comprises the formula




70. A compound according to claim 68, wherein V are selected so that the ring is a unsubstituted or substituted piperidin-3-yl moiety.


71. A compound according to claim 68, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
72. A compound according to claim 68, wherein M is selected from the group consisting of:



73. A compound according to claim 68, wherein M comprises a hydroxamic acid moiety.
74. A compound according to claim 68, wherein -L-M is




75. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;


each X is independently selected from the group consisting of CR_{12} and N;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the ring.

76. A compound according to claim 75, wherein the compound comprises the formula




77. A compound according to claim 75, wherein V are selected so that the ring is an N-substituted piperidin-3-yl moiety.

78. A compound according to claim 75, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
79. A compound according to claim 75, wherein M is selected from the group consisting of:



80. A compound according to claim 75, wherein M comprises a hydroxamic acid moiety.
81. A compound comprising a formula selected from the group consisting of:



wherein

each V is independently selected from the group consisting of $\text{C}(\text{R}_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


82. A compound according to claim 81, wherein the compound comprises the formula



83. A compound according to claim 81, wherein V are selected so that the ring is an N-substituted piperdin-3-yl moiety.


84. A compound according to claim 81, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.

85. A compound according to claim 81, wherein M is selected from the group consisting of:



86. A compound according to claim 81, wherein M comprises a hydroxamic acid moiety.

87. A compound according to claim 81, wherein -L-M is




88. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

89. A compound according to claim 88 wherein the compound comprises the formula




90. A compound according to claim 88, wherein V are selected so that the ring is an N-substituted piperidin-3-yl moiety.
91. A compound of claim 88, wherein Q is a substituted or unsubstituted phenyl ring.
92. A compound of claim 88, wherein Q is a substituted or unsubstituted heteroaryl.
93. A compound of claim 88, wherein Q is a substituted or unsubstituted heteroaryl selected from the group consisting of furan, thiophene, pyrrole, pyrazole, triazole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, benzofuran, isobenzofuran, benzothiophene, isobenzothiophene, indole, isobenzazole, quinoline, isoquinoline, cinnoline, quinazoline, naphthyridine, pyridopyridine, quinoxaline, phthalazine, benthiazole, and triazine.
94. A compound according to claim 88, wherein at least one X in the six membered ring is a substituted carbon atom.
95. A compound according to claim 88, wherein at least one of the X substituents in the six membered ring is -CF.
96. A compound according to claim 88, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
97. A compound according to claim 88, wherein M is selected from the group consisting of:




98. A compound according to claim 88, wherein M comprises a hydroxamic acid moiety.

99. A compound according to claim 88, wherein -Q-L-M is




100. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_2 , R_3 , R_4 and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

101. A compound according to claim 100, wherein the compound comprises the formula




102. A compound according to claim 100, wherein V are selected so that the ring is an N-substituted piperdin-3-yl moiety.
103. A compound according to claim 100, wherein at least one of R₂, R₃, R₄, or R₅ is fluorine.
104. A compound according to claim 100, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
105. A compound according to claim 100, wherein M is selected from the group consisting of:




106. A compound according to claim 100, wherein M comprises a hydroxamic acid moiety.

107. A compound according to claim 100, wherein -Q-L-M is



108. A compound comprising a formula selected from the group consisting of:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;


R_6 , R_7 , R_8 , and R_9 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

109. A compound according to claim 108, wherein the compound comprises the formula




110. A compound according to claim 108, wherein V are selected so that the ring is an N-substituted piperdin-3-yl moiety.
111. A compound according to claim 108, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.


112. A compound according to claim 108, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
113. A compound according to claim 108, wherein M is selected from the group consisting of:



114. A compound according to claim 108, wherein M comprises a hydroxamic acid moiety.
115. A compound according to claim 108, wherein -L-M is



116. A compound comprising a formula selected from the group consisting of:



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_6 , R_7 , R_8 , and R_9 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent

and the ring.

117. A compound according to claim 116, wherein the compound comprises the formula




118. A compound according to claim 116, wherein V are selected so that the ring is an N-substituted piperidin-3-yl moiety.
119. A compound according to claim 116, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
120. A compound according to claim 116, wherein M is selected from the group consisting of:





121. A compound according to claim 116, wherein M comprises a hydroxamic acid moiety.

122. A compound according to claim 116, wherein -L-M is



123. A compound according to claim 116, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.
124. A compound according to claim 116, wherein at least one X in the six membered ring is a substituted carbon atom.
125. A compound according to claim 116, wherein at least one of the X substituents in the six membered ring is -CF.
126. A compound comprising a formula selected from the group consisting of:





wherein

each V is independently selected from the group consisting of C(R₁₂)₂ and NR₁₂ where at least one V is NR₁₂;

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;


R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.

127. A compound according to claim 126, wherein the compound comprises the formula




128. A compound according to claim 126, wherein V are selected so that the ring is an N-substituted piperdin-3-yl moiety.
129. A compound according to claim 126, wherein at least one of R₂, R₃, R₄, or R₅ is fluorine.
130. A compound according to claim 126, wherein at least one of R₆, R₇, R₈, and R₉ is fluorine.
131. A compound according to claim 126, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
132. A compound according to claim 126, wherein M is selected from the group consisting of:



133. A compound according to claim 126, wherein M comprises a hydroxamic acid

moiety.

134. A compound according to claim 126, wherein -L-M is




135. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;


R_{10} and R_{11} are taken together to form a substituted or unsubstituted aromatic ring;
each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

Q is a substituted or unsubstituted aromatic ring;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

136. A compound according to claim 135, wherein V are selected so that the ring is an N-substituted piperdin-3-yl moiety.
137. A compound according to claim 135, wherein the substituted or unsubstituted aromatic ring formed when R_{10} and R_{11} are taken together is selected from the group consisting of substituted or unsubstituted aryl and heteroaryl.
138. A compound according to claim 135, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
139. A compound according to claim 135, wherein M is selected from the group consisting of:




140. A compound according to claim 135, wherein M comprises a hydroxamic acid moiety.

141. A compound according to claim 135, wherein -L-M is



142. A compound comprising the formula



wherein

each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

each X is independently selected from the group consisting of CR_{12} and N;

each Y is independently selected from the group consisting of O, S and NR_{12} ;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro,


and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 0-10 atoms separation between M and the remainder of the compound.

143. A compound according to claim 142, wherein V are selected so that the ring is an N-substituted piperidin-3-yl moiety.
144. A compound according to claim 142, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
145. A compound according to claim 142, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
146. A compound according to claim 142, wherein M is selected from the group consisting of:




147. A compound according to claim 142, wherein M comprises a hydroxamic acid moiety.

148. A compound according to claim 142, wherein -L-M is




149. A compound comprising the formula:



wherein

Z-Q- is selected from the group consisting of



each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12} where at least one V is NR_{12} ;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of


hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and


L is a substituent providing between 2-10 atoms separation between M and the Q substituent.

150. A compound according to claim 149, wherein V are selected so that the ring is an N-substituted piperdin-3-yl moiety.
151. A compound according to claim 149, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
152. A compound according to claim 149, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
153. A compound according to claim 149, wherein M is selected from the group consisting of:




154. A compound according to claim 149, wherein M comprises a hydroxamic acid moiety.

155. A compound according to claim 149, wherein -L-M is




156. A compound comprising the formula:



wherein

Z-Q- is selected from the group consisting of




each V is independently selected from the group consisting of $C(R_{12})_2$ and NR_{12}

where at least one V is NR_{12} ;

R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, cyano, and nitro;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

M is selected from the group consisting of




and L is E, Z or mixtures of E/Z $-\text{CH}_2=\text{CH}_2-$.

157. A compound according to claim 156, wherein V are selected so that the ring is an N-substituted piperidin-3-yl moiety.
158. A compound according to claim 156 wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
159. A compound according to claim 156, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
160. A compound according to claim 156, wherein M is selected from the group consisting of:




161. A compound according to claim 156, wherein M comprises a hydroxamic acid moiety.

162. A compound according to claim 156, wherein -L-M is



163. A compound comprising a formula selected from the group consisting of:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl,

heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between M and the ring.


164. A compound according to claim 163, wherein the compound comprises a formula selected from the group consisting of



165. A compound according to claim 163, wherein the compound comprises a formula selected from the group consisting of




166. A compound according to claim 163, wherein the compound comprises the formula




167. A compound according to claim 163, wherein R₁₄ comprises a member selected

from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.


168. A compound according to claim 163, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
169. A compound according to claim 163, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
170. A compound according to claim 163, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
171. A compound according to claim 163, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
172. A compound according to claim 163, wherein M is selected from the group consisting of:



173. A compound according to claim 163, wherein M comprises a hydroxamic acid moiety.
174. A compound according to claim 163, wherein $-L-M$ is




175. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or

unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between M and the ring.


176. A compound according to claim 175, wherein the compound comprises a formula selected from the group consisting of




177. A compound according to claim 175, wherein the compound comprises a formula selected from the group consisting of



178. A compound according to claim 175 wherein the compound comprises the formula




179. A compound according to claim 175, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
180. A compound according to claim 175, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
181. A compound according to claim 175, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
182. A compound according to claim 175, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
183. A compound according to claim 175, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
184. A compound according to claim 175, wherein M is selected from the group consisting of:




185. A compound according to claim 175, wherein M comprises a hydroxamic acid moiety.

186. A compound according to claim 175, wherein -L-M is




187. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

Q is a substituted or unsubstituted aromatic ring;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


188. A compound according to claim 187, wherein the compound comprises a formula selected from the group consisting of



189. A compound according to claim 187, wherein the compound comprises a formula selected from the group consisting of




190. A compound according to claim 187, wherein the compound comprises the formula




191. A compound according to claim 190, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
192. A compound according to claim 190, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
193. A compound according to claim 190, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
194. A compound according to claim 190, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
195. A compound of claim 187, wherein Q is a substituted or unsubstituted phenyl ring.

196. A compound of claim 187, wherein Q is a substituted or unsubstituted heteroaryl.
197. A compound of claim 187, wherein Q is a substituted or unsubstituted heteroaryl selected from the group consisting of furan, thiophene, pyrrole, pyrazole, triazole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, benzofuran, isobenzofuran, benzothiophene, isobenzothiophene, indole, isobenzazole, quinoline, isoquinoline, cinnoline, quinazoline, naphthyridine, pyridopyridine, quinoxaline, phthalazine, benthiazole, and triazine.
198. A compound according to claim 187, wherein at least one X in the six membered ring is a substituted carbon atom.
199. A compound according to claim 187, wherein at least one of the X substituents in the six membered ring is -CF.
200. A compound according to claim 187, wherein M comprises a member selected from the group consisting of trifluoroacetyl (-C(O)-CF₃), -NH-P(O)OH-CH₃, sulfonamides (-SO₂NH₂), hydroxysulfonamides (-SO₂NHOH), thiols(-SH), and carbonyl groups having the formula -C(O)-R₁₃ wherein R₁₃ is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
201. A compound according to claim 187, wherein M is selected from the group consisting of:



202. A compound according to claim 187, wherein M comprises a hydroxamic acid moiety.

203. A compound according to claim 187, wherein -Q-L-M is




204. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₂, R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted; and

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

Q is a substituted or unsubstituted aromatic ring;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.


205. A compound according to claim 204, wherein the compound comprises a formula selected from the group consisting of



206. A compound according to claim 204, wherein the compound comprises a formula selected from the group consisting of




207. A compound according to claim 204, wherein the compound comprises the formula




208. A compound according to claim 204, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
209. A compound according to claim 204, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
210. A compound according to claim 204, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
211. A compound according to claim 204, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
212. A compound according to claim 204, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
213. A compound according to claim 204, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.

214. A compound according to claim 204, wherein M is selected from the group consisting of:




215. A compound according to claim 204, wherein M comprises a hydroxamic acid moiety.

216. A compound according to claim 204, wherein -Q-L-M is



217. A compound comprising a formula selected from the group consisting of:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.


218. A compound according to claim 217, wherein the compound comprises a formula selected from the group consisting of



219. A compound according to claim 217, wherein the compound comprises a formula selected from the group consisting of




220. A compound according to claim 217, wherein the compound comprises the formula




221. A compound according to claim 217, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
222. A compound according to claim 217, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
223. A compound according to claim 217, wherein R_{14} is a substituted or unsubstituted $-\text{C}(\text{O})\text{C}_{1-6}$ alkyl.


224. A compound according to claim 217, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
225. A compound according to claim 217, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.
226. A compound according to claim 217, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
227. A compound according to claim 217, wherein M is selected from the group consisting of:



228. A compound according to claim 217, wherein M comprises a hydroxamic acid moiety.
229. A compound according to claim 217, wherein $-\text{L}-\text{M}$ is



230. A compound comprising a formula selected from the group consisting of:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₆, R₇, R₈, and R₉ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;

R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.


231. A compound according to claim 230, wherein the compound comprises a formula selected from the group consisting of



232. A compound according to claim 230, wherein the compound comprises a formula selected from the group consisting of




233. A compound according to claim 230, wherein the compound comprises the formula




234. A compound according to claim 230, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
235. A compound according to claim 230, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
236. A compound according to claim 230, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.

237. A compound according to claim 230, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
238. A compound according to claim 230, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})\text{OH}-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
239. A compound according to claim 230, wherein M is selected from the group consisting of:




240. A compound according to claim 230, wherein M comprises a hydroxamic acid moiety.
241. A compound according to claim 230, wherein $-\text{L}-\text{M}$ is



242. A compound according to claim 230, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.

243. A compound according to claim 230, wherein at least one X in the six membered ring is a substituted carbon atom.
244. A compound according to claim 230, wherein at least one of the X substituents in the six membered ring is -CF.
245. A compound comprising a formula selected from the group consisting of:



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are each independently selected from the group

consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;


each R_{12} is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R_{12} is not halo, cyano, nitro, and thio in the case where the ring atom to which R_{12} is bound is nitrogen;

R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the ring.


246. A compound according to claim 245, wherein the compound comprises a formula selected from the group consisting of




247. A compound according to claim 245, wherein the compound comprises a formula selected from the group consisting of



248. A compound according to claim 245, wherein the compound comprises the formula




249. A compound according to claim 245, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
250. A compound according to claim 245, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
251. A compound according to claim 245, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
252. A compound according to claim 245, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
253. A compound according to claim 245, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
254. A compound according to claim 245, wherein at least one of R_6 , R_7 , R_8 , and R_9 is fluorine.
255. A compound according to claim 245, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
256. A compound according to claim 245, wherein M is selected from the group consisting of:




257. A compound according to claim 245, wherein M comprises a hydroxamic acid moiety.

258. A compound according to claim 245, wherein -L-M is




259. A compound comprising the formula



wherein

Z is selected from the group consisting of



wherein

each X is independently selected from the group consisting of CR₁₂ and N;

each Y is independently selected from the group consisting of O, S and NR₁₂;

R₁₀ and R₁₁ are taken together to form a substituted or unsubstituted aromatic ring;

each R₁₂ is independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted, with the proviso that R₁₂ is not halo, cyano, nitro, and thio in the case where the ring atom to which R₁₂ is bound is nitrogen;


R₁₄ is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

Q is a substituted or unsubstituted aromatic ring;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the Q substituent.

260. A compound according to claim 259, wherein the compound comprises a formula selected from the group consisting of




261. A compound according to claim 259, wherein the compound comprises a formula selected from the group consisting of




262. A compound according to claim 259, wherein R₁₄ comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.

263. A compound according to claim 259, wherein R₁₄ is a substituted or unsubstituted C₁₋₆ alkyl.


264. A compound according to claim 259, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
265. A compound according to claim 259, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
266. A compound according to claim 259, wherein the substituted or unsubstituted aromatic ring formed when R_{10} and R_{11} are taken together is selected from the group consisting of substituted or unsubstituted aryl and heteroaryl.
267. A compound according to claim 259, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
268. A compound according to claim 259, wherein M is selected from the group consisting of:



269. A compound according to claim 259, wherein M comprises a hydroxamic acid moiety.
270. A compound according to claim 259, wherein $-L-M$ is



271. A compound comprising the formula



wherein


R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 0-10 atoms separation between the M substituent and the remainder of the compound.

272. A compound according to claim 271, wherein the compound comprises a formula selected from the group consisting of




273. A compound according to claim 271, wherein the compound comprises a formula selected from the group consisting of



274. A compound according to claim 271, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
275. A compound according to claim 271, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
276. A compound according to claim 271, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
277. A compound according to claim 271, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
278. A compound according to claim 271, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.


279. A compound according to claim 271, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-\text{C}(\text{O})-\text{CF}_3$), $-\text{NH}-\text{P}(\text{O})(\text{OH})-\text{CH}_3$, sulfonamides ($-\text{SO}_2\text{NH}_2$), hydroxysulfonamides ($-\text{SO}_2\text{NHOH}$), thiols ($-\text{SH}$), and carbonyl groups having the formula $-\text{C}(\text{O})-\text{R}_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.

280. A compound according to claim 271, wherein M is selected from the group consisting of:




281. A compound according to claim 271, wherein M comprises a hydroxamic acid moiety.

282. A compound according to claim 271, wherein $-\text{L}-\text{M}$ is




283. A compound comprising the formula:



wherein

$\text{Z}-\text{Q}-$ is selected from the group consisting of



wherein


R_2 , R_3 , R_4 , and R_5 are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, thio, cyano, nitro, and a carbonyl group, each substituted or unsubstituted;

R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;


M is a substituent capable of complexing with a deacetylase catalytic site and/or a metal ion; and

L is a substituent providing between 2-10 atoms separation between the M substituent and the Q substituent.


284. A compound according to claim 283, wherein the compound comprises a formula selected from the group consisting of



285. A compound according to claim 283, wherein the compound comprises a formula selected from the group consisting of




286. A compound according to claim 283, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
287. A compound according to claim 283, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
288. A compound according to claim 283, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
289. A compound according to claim 283, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
290. A compound according to claim 283, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
291. A compound according to claim 283, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)-CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
292. A compound according to claim 283, wherein M is selected from the group consisting of:



293. A compound according to claim 283, wherein M comprises a hydroxamic acid moiety.

294. A compound according to claim 283, wherein -L-M is




295. A compound comprising the formula:



wherein

Z-Q- is selected from the group consisting of




wherein

R₂, R₃, R₄, and R₅ are each independently selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, cyano, and nitro;


R_{14} is selected from the group consisting of hydrogen, halo, alkyl, alkoxy, aryl, heteroaryl, aminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryloxy, heteroaryloxy, arylalkyl, heteroarylalkyl, amino, and a carbonyl group, each substituted or unsubstituted;

M is selected from the group consisting of




and L is E, Z or mixtures of E/Z -CH₂=CH₂-.


296. A compound according to claim 295, wherein the compound comprises a formula selected from the group consisting of



297. A compound according to claim 295, wherein the compound comprises a formula selected from the group consisting of



298. A compound according to claim 295, wherein R_{14} comprises a member selected from the group consisting of hydrogen and a substituent that is convertible *in vivo* to hydrogen.
299. A compound according to claim 295, wherein R_{14} is a substituted or unsubstituted C_{1-6} alkyl.
300. A compound according to claim 295, wherein R_{14} is a substituted or unsubstituted $-C(O)C_{1-6}$ alkyl.
301. A compound according to claim 295, wherein R_{14} is selected from the group consisting of H, methyl, ethyl, propyl, isopropyl, butyl, acetyl, and BOC.
302. A compound according to claim 295, wherein at least one of R_2 , R_3 , R_4 , or R_5 is fluorine.
303. A compound according to claim 295, wherein M comprises a member selected from the group consisting of trifluoroacetyl ($-C(O)CF_3$), $-NH-P(O)OH-CH_3$, sulfonamides ($-SO_2NH_2$), hydroxysulfonamides ($-SO_2NHOH$), thiols ($-SH$), and carbonyl groups having the formula $-C(O)-R_{13}$ wherein R_{13} is hydroxylamino, hydroxyl, amino, alkylamino, or an alkoxy group.
304. A compound according to claim 295, wherein M is selected from the group consisting of:



305. A compound according to claim 295, wherein M comprises a hydroxamic acid moiety.
306. A compound according to claim 295, wherein -L-M is




FIGURE 1




FIGURE 2A




FIGURE 2A (cont.)




FIGURE 2A (cont.)




FIGURE 2A (cont.)




FIGURE 2A (cont.)




FIGURE 2B




FIGURE 2C




FIGURE 2C (cont.)




FIGURE 2D




FIGURE 3

SEQ. ID No. 1

MSYYHHHHHDYDIPTTENLYFQGAMEPGGSMAQTQGTTRRKVCYYYYDGDVGNYYYYGQGHF
MKPHRIRMTHNLLNLYGLYRKMEIYRPHKANAEEMTKYHSDDYIKFLRSIRPDNMSEYSK
QMQRFNVGEDCPVFDGLFEFCQLSTGGSVASAVKLNKQQTDI AVNWAGGLHHAKKSEASG
FCYVNDIVLAILELLKYHQRVLYIDIDIHHGDGVEEAFYTTDRVMTVSFHKYGEYFPGTG
DLRDIGAGKGKYYAVNYPLRDGIDDES YEAFKPVMSKVMEMFQPSAVVLQCGSDSLSGD
RLGCFNLTIKGHAKCVEFVKSFNLPMLMLGGGGYTIRNVARCWYETETAVALDTEIPNELP
YNDYFEYFGPDFKLHISPSNMTNQNTNEYLEKIKQRLFENLRMLPHAPGVQMQAIPEDAI
PEESGDEDEDDPKRISICSSDKRIACEEEFSDSEEEGEGGRKNSSNFKKAKRVKTEDEK
EKDPEEKKEVT EEEKTKEEKPEAKGVKEEVKLA

FIGURE 4

SEQ. ID No. 2

ATGTCGTACTACCATCACCATCACCATCACGATTACGATATCCCAACGACCGAAAACCTG
TATTTTCAGGGCGCCATGGAACCCGGGGGATCCATGGCGCAGACGCAGGGCACCCGGAGG
AAAGTCTGTTACTACTACGACGGGGATGTTGGAAATTACTATTATGGACAAGGCCACCCA
ATGAAGCCTCACC GAATCCGCATGACTCATAATTTGCTGCTCAACTATGGTCTCTACCGA
AAAATGGAAATCTATCGCCCTCACAAGCCAATGCTGAGGAGATGACCAAGTACCACAGC
GATGACTACATTAAATTCTTGCGCTCCATCCGTCCAGATAACATGTCGGAGTACAGCAAG
CAGATGCAGAGATTCAACGTTGGTGAGGACTGTCCAGTATTCGATGGCCTGTTTGAGTTC
TGTCAGTTGTCTACTGGTGGTTCTGTGGCAAGTGCTGTGAACTTAATAAGCAGCAGACG
GACATCGCTGTGAATTGGGCTGGGGCCTGCACCATGCAAAGAAGTCCGAGGCATCTGGC
TTCTGTTACGTCAATGATATCGTCTTGCCATCCTGGAAGTCTAAAGTATCACCAGAGG
GTGCTGTACATTGACATTGATATTCACCATGGTGACGGCGTGGAAGAGGCCTTCTACACC
ACGGACCGGGTCATGACTGTGTCTTTTCATAAGTATGGAGAGTACTTCCAGGAAC TGGG
GACCTACGGGATATCGGGGCTGGCAAAGGCAAGTATTATGCTGTAACTACCCGCTCCGA
GACGGGATTGATGACGAGTCCTATGAGGCCATTTTCAAGCCGGTCATGTCAAAGTAATG
GAGATGTTCCAGCCTAGTGCGGTGGTCTTACAGTGTGGCTCAGACTCCCTATCTGGGGAT
CGGTTAGGTTGCTTCAATCTAACTATCAAAGGACACGCCAAGTGTGTGGAATTTGTCAAG
AGCTTTAACCTGCCTATGCTGATGCTGGGAGGCGGTGGTTACACCATTCGTAACGTTGCC
CGGTGCTGGACATATGAGACAGCTGTGGCCCTGGATACGGAGATCCCTAATGAGCTTCCA
TACAATGACTACTTTGAATACTTTGGACCAGATTTCAAGCTCCACATCAGTCCTTCCAAT
ATGACTAACCAGAACACGAATGAGTACCTGGAGAAGATCAAACAGCGACTGTTTGAGAAC
CTTAGAATGCTGCCGCACGCACCTGGGGTCCAAATGCAGGCGATTCTTGAGGACGCCATC
CCTGAGGAGAGTGCGCATGAGGACGAAGACGACCCTGACAAGCGCATCTCGATCTGCTCC
TCTGACAAACGAATTGCCTGTGAGGAAGAGTTCTCCGATTCTGAAGAGGAGGGAGAGGGG
GGCCGCAAGAACTCTTCAAATTCAAAAAAGCCAAGAGAGTCAAAACAGAGGATGAAAAA
GAGAAAGACCCAGAGGAGAAGAAAGAGTACCGAAGAGGAGAAAACCAAGGAGGAGAAG
CCAGAAGCCAAAGGGTCAAGGAGGAGGTCAAGTTGGCCTGA

FIGURE 5

SEQ. ID No. 3

MGSMAYSQGGGKKKVCYYYDGDIGNYYYQGHPMKPHRIRMTHNLLLNYGLYRKMEIYRP
HKATAEEMTKYHSDEYIKFLRSIRPDNMSEYSKQMQRFNVGEDCPVFDGLFEFCQLSTGG
SVAGAVKLNRRQQTDMAVNWAGGLHHAKKSEASGFCYVNDIVLAILLELLKYHQRVLYIDID
IHHGDBGVEEAFYTTDRVMTVSFHKYGEYFPGTGDRLRDIGAGKGKYYAVNFPMRDGIIDDES
YGQIFKPIISKVMEYQPSAVVLQCGADSLSGDRLGCFNLTVKGHAKCDEVVKTFFNLPLL
MLGGGGYTIRNVARCWYETAVALDCEIPNELPYNDYFEYFGPDFKLHISPSNMTNQNTF
EYMEKIKQRLFENLRMLPHAPGVQMQAIPEDAVHEDSGDEDGEDPDKRISIRASDKRIAC
DEEFSDESEDEGEGGRNVADHKKGAKKARIEEDKKETEDKKTVDVKEEDKSKDNSGEKTD
TGKTKSEQLSNPGHHHHHH

FIGURE 6

SEQ. ID No. 4

ATGGGATCCATGGCGTACAGTCAAGGAGGCGGCAAAAAAAAAAGTCTGCTACTACTACGAC
GGTGATATTGGAAATTATTATTATGGACAGGGTCATCCCATGAAGCCTCATAGAATCCGC
ATGACCCATAACTTGCTGTAAATTATGGCTTATACAGAAAAATGGAAATATATAGGCC
CATAAAGCCACTGCCGAAGAAATGACAAAATATCACAGTGATGAGTATATCAAATTTCTA
CGGTCAATAAGACCAGATAACATGTCTGAGTATAGTAAGCAGATGCAGAGATTTAATGTT
GGAGAAGATTGTCCAGTGTTTGATGGACTCTTTGAGTTTTGTCAGCTCTCAACTGGCGGT
TCAGTTGCTGGAGCTGTGAAGTTAAACCGACAACAGACTGATATGGCTGTTAATTGGGCT
GGAGGATTACATCATGCTAAGAAATCAGAAGCATCAGGATTCTGTTACGTTAATGATATT
GTGCTTGCCATCCTTGAATTACTAAAGTATCATCAGAGAGTCTTATATATTGATATAGAT
ATTCATCATGGTGATGGTGTTGAAGAAGCTTTTTTATACAACAGATCGTGTAATGACGGTA
TCATTCCATAAATATGGGGAATACTTTCCTGGCAGAGAGACTTGAGGGATATTGGTGCT
GGAAAAGGCAAATACTATGCTGTCAATTTTCCAATGAGAGATGGTATAGATGATGAGTCA
TATGGGCAGATATTTAAGCCTATTATCTCAAAGGTGATGGAGATGTATCAACCTAGTGCT
GTGGTATTACAGTGTGGTGACAGACTCATTATCTGGTGATAGACTGGGTTGTTTCAATCTA
ACAGTCAAAGGTCATGCTAAATGTGTAGAAGTTGTAAAACTTTTAACTTACCATTACTG
ATGCTTGAGGAGGTGGCTACACAATCCGTAATGTTGCTCGATGTTGGACATATGAGACT
GCAGTTGCCCTTGATTGTGAGATTCCCAATGAGTTGCCATATAATGATTACTTTGAGTAT
TTTGGACCAGACTTCAAACCTGCATATTAGTCCTTCAAACATGACAAACAGAACACTCCA
GAATATATGGAAAAGATAAAACAGCGTTTGTGTTGAAAATTTGCGCATGTTACCTCATGCA
CCTGGTGTCCAGATGCAAGCTATTCCAGAAGATGCTGTTTCATGAAGACAGTGGAGATGAA
GATGGAGAAGATCCAGACAAGAGAATTTCTATTGAGCATCAGACAAGCGGATAGCTTGT
GATGAAGAATTCTCAGATTCTGAGGATGAAGGAGAAGGAGGTGGAAGAAATGTGGCTGAT
CATAAGAAAGGAGCAAAGAAAGCTAGAATTGAAGAAGATAAGAAAGAAACAGAGGACAAA
AAACAGACGTTAAGGAAGAAAGATAAATCCAAGGACAACAGTGGTGAAAAAACAGATACC
AAAGGAACCAAATCAGAACAGCTCAGCAACCCCGGCATCACCATCACCATCACTAA

FIGURE 7

SEQ. ID No. 5

MPGMDLNLEAEALAGTGLVLDEQLNEFHCLWDDSFPEGPERLHAIKEQLIQEGLLDRCVS
FQARFAEKEELMLVHSLEYIDLMETTQYMNEGELRVLADTYDSVYLHPNSYSCACLASGS
VLRVDAVLGAERNGMAIIRPPGHHAQHSMDGYCMFNHVAVAAARYAQQKHRIRRVLI
DWDVHHGQGTQFTFDQDPSVLYFSIHRYEQGRFWPHLKASNWSTTGFGQGQGYTINVPWN
QVGMRDADYIAAFLHVLLPVALEFQPPQLVLVAAGFDALQGDPKGEMAATPAGFAQLTHLL
MGLAGGKLILSLEGGYNLRALAEGVSASLHTLLGDPCPMLESPGAPCRSAQASVSCALEA
LEPFWEVLVRSTETVERDNMEEDNVEESEESEEGPWEPVLPILTWPVLQSRTGLVYDQNM
NHCNLWDSHHPEVPQRIIRIMCRLEELGLAGRCLTLTPRPATEAELLTCHSAEYVGHRLA
TEKMKTRELHRESSNFDSIYICPSTFACAQLATGAACRLVEAVLSGEVLNGAAVVRPPGH
HAEQDAACGFCFFNSVAVAAARHAQTISGHALRILIVDWDVHHGNGTQHMFEDDPSVLYVS
LHRYDHGTTFFPMGDEGASSQIGRAAGTGFTVNVANGPRMGDADYLAAWHRLVLPYAYEF
NPELVLSAGFDAARGDPLGGCQVSPEGYAHLTHLLMGLASGRIILILEGGYNLTSISES
MAACTRSLLGDPPLLTLPPLSGALASITETIQVHRRYWRSRLVMKVEDREGPGHHHH
HH

FIGURE 9

SEQ. ID No. 7

MHHHHHHPMEEPEEPADSGQSLVPVYIYSPEYVSMCDLAKIPKRASMVHSLIEAYALHK
QMRIVKPKVASMEEMAAFTDAYLQHLQKVSQEGDDHDPDSIEYGLGYDCPATEGIFDYA
AAIGGATITAAQCLIDGMCKVAINWSGGWHHAKKDEASGFCYLNDAVLGILRLRRKFERI
LYVDLDLHHGDGVEDAFSFTSKVMTVSLHKFSPGFFPGTGDVSDVGLGKGRIYSVNVPIQ
DGIQDEKYYQICESVLKEVYQAFNPKAVVLQLGADTIAGDPMCSFNMTVPVGIGKCLKYIL
QWQLATLILGGGGYNLANTARCWTYLTGVILGKTLSSSEIPDHEFFTAYGPDYVLEITPSC
RPDRNEPHRIQQILNYIKGNLKHVV

FIGURE 8

SEQ. ID No. 6

ATGCCCCGGGATGGATCTGAACCTTGAGGCTGAAGCACTGGCTGGCACTGGCTTGGTGTG
GATGAGCAGTTAAATGAATTCCATTGCCTCTGGGATGACAGCTTCCCGGAAGGCCCTGAG
CGGCTCCATGCCATCAAGGAGCAACTGATCCAGGAGGGCCTCCTAGATCGCTGCGTGTCC
TTTCAGGCCCGGTTTGCTGAAAAGGAAGAGCTGATGTTGGTTCACAGCCTAGAATATATT
GATCTGATGGAAACAACCCAGTACATGAATGAGGGAGAACTCCGTGTCTAGCAGACACC
TACGACTCAGTTTATCTGCATCCGAACTCATACTCCTGTGCCTGCCTGGCCTCAGGCTCT
GTCCTCAGGCTGGTGGATGCGGTCTGGGGGCTGAGATCCGGAATGGCATGGCCATCATT
AGGCCTCCTGGACATCACGCCCAGCACAGTCTTATGGATGGCTATTGCATGTTCAACCAC
GTGGCTGTGGCAGCCCCGCTATGCTCAACAGAAACACCGCATCCGGAGGGTCCTTATCGTA
GATTGGGATGTGCACCACGGTCAAGGAACACAGTTACCTTCGACCAGGACCCCACTGTC
CTCTATTTCTCCATCCACCGCTACGAGCAGGGTAGGTTCTGGCCCCACCTGAAGGCCTCT
AACTGGTCCACCACAGGTTTCGGCCAAGGCCAAGGATATACCATCAATGTGCCTTGGAAC
CAGGTGGGGATGCGGGATGCTGACTACATTGCTGCTTTCCTGCACGTCCTGCTGCCAGTC
GCCCTCGAGTTCAGCCTCAGCTGGTCTGGTGGCTGCTGGATTTGATGCCCTGCAAGGG
GACCCCAAGGGTGAGATGGCCGCCACTCCGGCAGGGTTCGCCCAGCTAACCCACCTGCTC
ATGGGTCTGGCAGGAGGCAAGCTGATCCTGTCTCTGGAGGGTGGCTACAACCTCCGCGCC
CTGGCTGAAGGCGTCAGTGCTTCGCTCCACACCCTTCTGGGAGACCCTTGCCCCATGCTG
GAGTCACCTGGTGCCCCCTGCCGGAGTGCCCAAGGCTTCACTTCTGTGCTCTGGAAGCC
CTTGAGCCCTTCTGGGAGGTTCTTGTGAGATCAACTGAGACCGTGGAGAGGGACAACATG
GAGGAGGACAATGTAGAGGAGAGCGAGGAGGAAGGACCCTGGGAGCCCCCTGTGCTCCCA
ATCCTGACATGGCCAGTGCTACAGTCTCGCACAGGGCTGGTCTATGACCAAAATATGATG
AATCACTGCAACTTGTGGGACAGCCACCACCCTGAGGTACCCAGCGCATCTTGCGGATC
ATGTGCCGTCTGGAGGAGCTGGGCCTTGCCGGGCGCTGCCTCACCCCTGACACCGCGCCCT
GCCACAGAGGCTGAGCTGCTCACCTGTACAGTGCTGAGTACGTGGGTCTCTCCGGGCC
ACAGAGAAAATGAAAACCCGGGAGCTGCACCGTGAGAGTTCCAACCTTTGACTCCATCTAT
ATCTGCCCCAGTACCTTTCGCTGTGCACAGCTTGCCACTGGCGCTGCCTGCCGCCTGGTG
GAGGCTGTGCTCTCAGGAGAGGTTCTGAATGGTGTGCTGTGTTGCGTCCCCCAGGACAC
CACGCAGAGCAGGATGCAGCTTGCGGTTTTTGTCTTTTCAACTCTGTGGCTGTGGCTGCT
CGCCATGCCCAGACTATCAGTGGGCATGCCCTACGGATCCTGATTGTGGATTGGGATGTC
CACCACGGTAATGGAACCTCAGCACATGTTTGAGGATGACCCAGTGTGCTATATGTGTCC
CTGCACCGCTATGATCATGGCACCTTCTTCCCCATGGGGGATGAGGGTGCCAGCAGCCAG
ATCGGCCGGGCTGCGGGCACAGGCTTACCGTCAACGTGGCATGGAACGGGCCCCGCATG
GGTGATGCTGACTACCTAGCTGCCTGGCATCGCTGGTGTCTCCATTGCCCTACGAGTTT
AACCAGAACTGGTGCTGGTCTCAGCTGGCTTTGATGCTGCACGGGGGGATCCGCTGGGG
GGCTGCCAGGTGTACCTGAGGGTTATGCCACCTCACCCACCTGCTGATGGGCCTTGCC
AGTGGCCGATTATCCTTATCCTAGAGGGTGGCTATAACCTGACATCCATCTCAGAGTCC
ATGGCTGCCTGCACTCGCTCCCTCCTTGAGACCCACCACCCCTGCTGACCCTGCCACGG
CCCCACTATCAGGGGCCCTGGCCTCAATCACTGAGACCATCCAAGTCCATCGCAGATAC
TGGCGCAGCTTACGGGTCATGAAGGTAGAAGACAGAGAAGGACCCGGGCATCACCATCAC
CATCACTAA

FIGURE 10

SEQ. ID No. 8

ATGCACCATCACCATCACCATCCCATGGAGGAGCCGGAGGAACCGGCGGACAGTGGGCAG
TCGCTGGTCCCGGTTTATATCTATAGTCCCGAGTATGTCAGTATGTGTGACTCCCTGGCC
AAGATCCCCAAACGGGCCAGTATGGTGCATTCTTTGATTGAAGCATATGCACTGCATAAG
CAGATGAGGATAGTTAAGCCTAAAGTGGCCTCCATGGAGGAGATGGCCGCCTTCCACACT
GATGCTTATCTGCAGCATCTCCAGAAGGTGAGCCAAGAGGGCGATGATGATCATCCGGAC
TCCATAGAATATGGGCTAGGTTATGACTGCCCAGCCACTGAAGGGATATTTGACTATGCA
GCAGCTATAGGAGGGGCTACGATCACAGCTGCCCAATGCCTGATTGACGGAATGTGCAAA
GTAGCAATTAACTGGTCTGGAGGGTGGCATCATGCAAAGAAAGATGAAGCATCTGGTTTT
TGTTATCTCAATGATGCTGTCTTGGGAATATTACGATTGCGACGGAATTTGAGCGTATT
CTCTACGTGGATTTGGATCTGCACCATGGAGATGGTGTAGAAGACGCATTTCAGTTTCACC
TCCAAAGTCATGACCGTGTCCCTGCACAAATTCTCCCCAGGATTTTCCCAGGAACAGGT
GACGTGTCTGATGTTGGCCTAGGGAAGGGACGGTACTACAGTGTAATGTGCCCATTTCAG
GATGGCATACAAGATGAAAAATATTACCAGATCTGTGAAAGTGTAATAAGGAAGTATAC
CAAGCCTTTAATCCCAAAGCAGTGGTCTTACAGCTGGGAGCTGACACAATAGCTGGGGAT
CCCATGTGCTCCTTTAACATGACTCCAGTGGGAATTGGCAAGTGTCTTAAGTACATCCTT
CAATGGCAGTTGGCAACACTCATTTTGGGAGGAGGAGGCTATAACCTTGCCAACACGGCT
CGATGCTGGACATACTTGACCGGGGTCATCCTAGGGAAAACACTATCCTCTGAGATCCCA
GATCATGAGTTTTTTCAGCATATGGTCTGATTATGTGCTGGAAATCACGCCAAGCTGC
CGGCCAGACCGCAATGAGCCCCACCGAATCCAACAAATCCTCAACTACATCAAAGGGAAT
CTGAAGCATGTGGTCTAG

SEQUENCE LISTING

<110> Syrrx, Inc.
 <120> HISTONE DEACETYLASE INHIBITORS
 <130> SYR-HDAC-5005-PCT
 <140> Not Yet Assigned
 <141> 2004-03-17
 <150> US 60/455,437
 <151> 2003-03-17
 <150> US 60/531,203
 <151> 2003-12-19
 <160> 8
 <170> PatentIn version 3.2
 <210> 1
 <211> 513
 <212> PRT
 <213> Custom

 <220>
 <221> Residues 1-482 of HDAC1 and a 6-histidine tag at the N-terminus
 <222> (1)..(513)
 <400> 1
 Met Ser Tyr Tyr His His His His His His Asp Tyr Asp Ile Pro Thr
 1 5 10 15
 Thr Glu Asn Leu Tyr Phe Gln Gly Ala Met Glu Pro Gly Gly Ser Met
 20 25 30
 Ala Gln Thr Gln Gly Thr Arg Arg Lys Val Cys Tyr Tyr Tyr Asp Gly
 35 40 45
 Asp Val Gly Asn Tyr Tyr Tyr Gly Gln Gly His Pro Met Lys Pro His
 50 55 60
 Arg Ile Arg Met Thr His Asn Leu Leu Leu Asn Tyr Gly Leu Tyr Arg
 65 70 75 80
 Lys Met Glu Ile Tyr Arg Pro His Lys Ala Asn Ala Glu Glu Met Thr
 85 90 95
 Lys Tyr His Ser Asp Asp Tyr Ile Lys Phe Leu Arg Ser Ile Arg Pro
 100 105 110
 Asp Asn Met Ser Glu Tyr Ser Lys Gln Met Gln Arg Phe Asn Val Gly
 115 120 125
 Glu Asp Cys Pro Val Phe Asp Gly Leu Phe Glu Phe Cys Gln Leu Ser
 130 135 140

Thr Gly Gly Ser Val Ala Ser Ala Val Lys Leu Asn Lys Gln Gln Thr
 145 150 155 160
 Asp Ile Ala Val Asn Trp Ala Gly Gly Leu His His Ala Lys Lys Ser
 165 170 175
 Glu Ala Ser Gly Phe Cys Tyr Val Asn Asp Ile Val Leu Ala Ile Leu
 180 185 190
 Glu Leu Leu Lys Tyr His Gln Arg Val Leu Tyr Ile Asp Ile Asp Ile
 195 200 205
 His His Gly Asp Gly Val Glu Glu Ala Phe Tyr Thr Thr Asp Arg Val
 210 215 220
 Met Thr Val Ser Phe His Lys Tyr Gly Glu Tyr Phe Pro Gly Thr Gly
 225 230 235 240
 Asp Leu Arg Asp Ile Gly Ala Gly Lys Gly Lys Tyr Tyr Ala Val Asn
 245 250 255
 Tyr Pro Leu Arg Asp Gly Ile Asp Asp Glu Ser Tyr Glu Ala Ile Phe
 260 265 270
 Lys Pro Val Met Ser Lys Val Met Glu Met Phe Gln Pro Ser Ala Val
 275 280 285
 Val Leu Gln Cys Gly Ser Asp Ser Leu Ser Gly Asp Arg Leu Gly Cys
 290 295 300
 Phe Asn Leu Thr Ile Lys Gly His Ala Lys Cys Val Glu Phe Val Lys
 305 310 315 320
 Ser Phe Asn Leu Pro Met Leu Met Leu Gly Gly Gly Gly Tyr Thr Ile
 325 330 335
 Arg Asn Val Ala Arg Cys Trp Thr Tyr Glu Thr Ala Val Ala Leu Asp
 340 345 350
 Thr Glu Ile Pro Asn Glu Leu Pro Tyr Asn Asp Tyr Phe Glu Tyr Phe
 355 360 365
 Gly Pro Asp Phe Lys Leu His Ile Ser Pro Ser Asn Met Thr Asn Gln
 370 375 380
 Asn Thr Asn Glu Tyr Leu Glu Lys Ile Lys Gln Arg Leu Phe Glu Asn
 385 390 395 400
 Leu Arg Met Leu Pro His Ala Pro Gly Val Gln Met Gln Ala Ile Pro

405 410 415

Glu Asp Ala Ile Pro Glu Glu Ser Gly Asp Glu Asp Glu Asp Asp Pro
420 425 430

Asp Lys Arg Ile Ser Ile Cys Ser Ser Asp Lys Arg Ile Ala Cys Glu
435 440 445

Glu Glu Phe Ser Asp Ser Glu Glu Glu Gly Glu Gly Gly Arg Lys Asn
450 455 460

Ser Ser Asn Phe Lys Lys Ala Lys Arg Val Lys Thr Glu Asp Glu Lys
465 470 475 480

Glu Lys Asp Pro Glu Glu Lys Lys Glu Val Thr Glu Glu Glu Lys Thr
485 490 495

Lys Glu Glu Lys Pro Glu Ala Lys Gly Val Lys Glu Glu Val Lys Leu
500 505 510

A7a

<210>	2
<211>	1542
<212>	DNA
<213>	Custom

```
<220>
<221> DNA sequence encoding residues 1-482 of HDAC1 and a 6-histidine
tag at the N-terminus
<222> (1)..(1542)
```

<400>	2								
atgtcgtact	accatcacca	tcaccatcac	gattacgata	tcccaacgac	cgaaaacctg				60
tattttcagg	gcgccatgga	acccggggga	tccatggcgc	agacgcaggg	caccgcgagg				120
aaagtctgtt	actactacga	cgggggatgtt	ggaaattact	attatggaca	aggccacca				180
atgaagcctc	accgaatccg	catgactcat	aatttgctgc	tcaactatgg	tctctaccga				240
aaaatggaaa	tctatcgccc	tcacaaagcc	aatgctgagg	agatgaccaa	gtaccacagc				300
gatgactaca	ttaaattctt	gcgctccatc	cgtccagata	acatgtcggg	gtacagcaag				360
cagatgcaga	gattcaacgt	tgggtgaggac	tgtccagtat	tcgatggcct	gtttgagttc				420
tgtcagttgt	ctactggtgg	ttctgtggca	agtgtgtgta	aacttaataa	gcagcagacg				480
gacatcgctg	tgaattgggc	tgggggcctg	caccatgcaa	agaagtccga	ggcatctggc				540
ttctgttacg	tcaatgatat	cgtcttggcc	atcctggaac	tgctaaagta	tcaccagagg				600
gtgtgtgtaca	ttgacattga	tattcaccat	ggtgacggcg	tggaagaggc	cttctacacc				660
acggaccggg	tcatgactgt	gtcctttcat	aagtatggag	agtacttccc	aggaactggg				720

```

gacctacggg atatcggggc tggcaaaggc aagtattatg ctgttaacta cccgctccga      780
gacgggattg atgacgagtc ctatgaggcc attttcaagc cggcatgctc caaagtaatg      840
gagatgttcc agcctagtagc ggtggtctta cagtgtggct cagactccct atctggggat      900
cggttaggtt gcttcaatct aactatcaaa ggacacgcca agtgtgtgga atttgtcaag      960
agctttaacc tgcctatgct gatgctggga ggcgggtggt acaccattcg taacgttgcc     1020
cggtgctgga catatgagac agctgtggcc ctggatacgg agatccctaa tgagcttcca     1080
tacaatgact actttgaata ctttggaacca gatttcaagc tccacatcag tccttccaat     1140
atgactaacc agaacacgaa tgagtacctg gagaagatca aacagcgact gtttgagaac     1200
cttagaatgc tgccgcacgc acctggggtc caaatgcagg cgattcctga ggacgccatc     1260
cctgaggaga gtggcgatga ggacgaagac gacctgaca agcgcctctc gatctgctcc     1320
tctgacaaac gaattgcctg tgaggaagag ttctccgatt ctgaagagga gggagagggg     1380
ggccgcaaga actcttccaa cttcaaaaaa gccaaagagag tcaaaacaga ggatgaaaaa     1440
gagaaagacc cagaggagaa gaaagaagtc accgaagagg agaaaaccaa ggaggagaag     1500
ccagaagcca aaggggtcaa ggaggaggtc aagttggcct ga                          1542

```

```

<210> 3
<211> 498
<212> PRT
<213> Custom

```

```

<220>
<221> Residues 1-488 of HDAC2 and a 6-histidine tag at the C-terminus
<222> (1)..(498)

```

```

<400> 3

```

```

Met Gly Ser Met Ala Tyr Ser Gln Gly Gly Gly Lys Lys Lys Val Cys
 1           5           10           15

```

```

Tyr Tyr Tyr Asp Gly Asp Ile Gly Asn Tyr Tyr Tyr Gly Gln Gly His
 20           25           30

```

```

Pro Met Lys Pro His Arg Ile Arg Met Thr His Asn Leu Leu Leu Asn
 35           40           45

```

```

Tyr Gly Leu Tyr Arg Lys Met Glu Ile Tyr Arg Pro His Lys Ala Thr
 50           55           60

```

```

Ala Glu Glu Met Thr Lys Tyr His Ser Asp Glu Tyr Ile Lys Phe Leu
 65           70           75           80

```

```

Arg Ser Ile Arg Pro Asp Asn Met Ser Glu Tyr Ser Lys Gln Met Gln
 85           90           95

```

```

Arg Phe Asn Val Gly Glu Asp Cys Pro Val Phe Asp Gly Leu Phe Glu
100          105          110

```

Phe Cys Gln Leu Ser Thr Gly Gly Ser Val Ala Gly Ala Val Lys Leu
 115 120 125
 Asn Arg Gln Gln Thr Asp Met Ala Val Asn Trp Ala Gly Gly Leu His
 130 135 140
 His Ala Lys Lys Ser Glu Ala Ser Gly Phe Cys Tyr Val Asn Asp Ile
 145 150 155 160
 Val Leu Ala Ile Leu Glu Leu Leu Lys Tyr His Gln Arg Val Leu Tyr
 165 170 175
 Ile Asp Ile Asp Ile His His Gly Asp Gly Val Glu Glu Ala Phe Tyr
 180 185 190
 Thr Thr Asp Arg Val Met Thr Val Ser Phe His Lys Tyr Gly Glu Tyr
 195 200 205
 Phe Pro Gly Thr Gly Asp Leu Arg Asp Ile Gly Ala Gly Lys Gly Lys
 210 215 220
 Tyr Tyr Ala Val Asn Phe Pro Met Arg Asp Gly Ile Asp Asp Glu Ser
 225 230 235 240
 Tyr Gly Gln Ile Phe Lys Pro Ile Ile Ser Lys Val Met Glu Met Tyr
 245 250 255
 Gln Pro Ser Ala Val Val Leu Gln Cys Gly Ala Asp Ser Leu Ser Gly
 260 265 270
 Asp Arg Leu Gly Cys Phe Asn Leu Thr Val Lys Gly His Ala Lys Cys
 275 280 285
 Val Glu Val Val Lys Thr Phe Asn Leu Pro Leu Leu Met Leu Gly Gly
 290 295 300
 Gly Gly Tyr Thr Ile Arg Asn Val Ala Arg Cys Trp Thr Tyr Glu Thr
 305 310 315 320
 Ala Val Ala Leu Asp Cys Glu Ile Pro Asn Glu Leu Pro Tyr Asn Asp
 325 330 335
 Tyr Phe Glu Tyr Phe Gly Pro Asp Phe Lys Leu His Ile Ser Pro Ser
 340 345 350
 Asn Met Thr Asn Gln Asn Thr Pro Glu Tyr Met Glu Lys Ile Lys Gln
 355 360 365
 Arg Leu Phe Glu Asn Leu Arg Met Leu Pro His Ala Pro Gly Val Gln

370 375 380
 Met Gln Ala Ile Pro Glu Asp Ala Val His Glu Asp Ser Gly Asp Glu
 385 390 395 400
 Asp Gly Glu Asp Pro Asp Lys Arg Ile Ser Ile Arg Ala Ser Asp Lys
 405 410 415
 Arg Ile Ala Cys Asp Glu Glu Phe Ser Asp Ser Glu Asp Glu Gly Glu
 420 425 430
 Gly Gly Arg Arg Asn Val Ala Asp His Lys Lys Gly Ala Lys Lys Ala
 435 440 445
 Arg Ile Glu Glu Asp Lys Lys Glu Thr Glu Asp Lys Lys Thr Asp Val
 450 455 460
 Lys Glu Glu Asp Lys Ser Lys Asp Asn Ser Gly Glu Lys Thr Asp Thr
 465 470 475 480
 Lys Gly Thr Lys Ser Glu Gln Leu Ser Asn Pro Gly His His His His
 485 490 495

His His

<210> 4
 <211> 1497
 <212> DNA
 <213> Custom

<220>
 <221> DNA sequence encoding residues 1-488 of HDAC2 and a 6-histidine
 tag at the C-terminus
 <222> (1)..(1497)

<400> 4
 atgggatcca tggcgtacag tcaaggaggc ggcaaaaaaa aagtctgcta ctactacgac 60
 ggtgatattg gaaattatta ttatggacag ggtcatccca tgaagcctca tagaatccgc 120
 atgaccata acttgctgtt aaattatggc ttatacagaa aaatggaaat atataggccc 180
 cataaagcca ctgccgaaga aatgacaaaa taccacagtg atgagtatat caaatttcta 240
 cgggtcaataa gaccagataa catgtctgag tatagtaagc agatgcagag atttaatgtt 300
 ggagaagatt gtccagtgtt tgatggactc tttgagtttt gtcagctctc aactggcggc 360
 tcagttgctg gagctgtgaa gttaaaccga caacagactg atatggctgt taattgggct 420
 ggaggattac atcatgctaa gaaatcagaa gcatcaggat tctgttacgt taatgatatt 480
 gtgcttgcca tccttgaatt actaaagtat catcagagag tcttatatat tgatatagat 540
 attcatcatg gtgatggtgt tgaagaagct ttttatacaa cagatcgtgt aatgacggta 600


```

tcattccata aatatgggga atactttcct ggcacaggag acttgagga tattggtgct 660
ggaaaaggca aatactatgc tgtcaatttt ccaatgagag atggtataga tgatgagtca 720
tatgggcaga tatttaagcc tattatctca aaggatgatg agatgtatca acctagtgc 780
gtggtattac agtggtggtc agactcatta tctggtgata gactgggttg tttcaatcta 840
acagtcaaag gtcagtctaa atgtgtagaa gttgtaaaaa cttttaactt accattactg 900
atgcttgagag gaggtggcta cacaatccgt aatgttgctc gatgttgac atatgagact 960
gcagttgcc ttgatttgta gattcccaat gagttgccat ataataatta ctttgagtat 1020
tttgaccag acttcaaact gcatattagt ctttcaaaca tgacaaacca gaacactcca 1080
gaatatatgg aaaagataaa acagcgtttg ttgtaaaatt tgcgcatgtt acctcatgca 1140
cctggtgtcc agatgcaagc tattccagaa gatgctgttc atgaagacag tggagatgaa 1200
gatggagaag atccagacaa gagaatttct attcgagcat cagacaagcg gatagcttgt 1260
gatgaagaat tctcagattc tgaggatgaa ggagaaggag gtcgaagaaa tgtggctgat 1320
cataagaaag gagcaaagaa agctagaatt gaagaagata agaaagaaac agaggacaaa 1380
aaaacagacg ttaaggaaga agataaatcc aaggacaaca gtggtgaaaa aacagatacc 1440
aaaggaacca aatcagaaca gctcagcaac cccgggcac accatcacca tcactaa 1497

```

```

<210> 5
<211> 782
<212> PRT
<213> Custom

```

```

<220>
<221> Residues 73-845 of HDAC6 and a 6-histidine tag at the C-terminus
<222> (1)..(782)

```

```

<400> 5

```

```

Met Pro Gly Met Asp Leu Asn Leu Glu Ala Glu Ala Leu Ala Gly Thr
1          5          10          15

```

```

Gly Leu Val Leu Asp Glu Gln Leu Asn Glu Phe His Cys Leu Trp Asp
          20          25          30

```

```

Asp Ser Phe Pro Glu Gly Pro Glu Arg Leu His Ala Ile Lys Glu Gln
          35          40          45

```

```

Leu Ile Gln Glu Gly Leu Leu Asp Arg Cys Val Ser Phe Gln Ala Arg
          50          55          60

```

```

Phe Ala Glu Lys Glu Glu Leu Met Leu Val His Ser Leu Glu Tyr Ile
65          70          75          80

```

```

Asp Leu Met Glu Thr Thr Gln Tyr Met Asn Glu Gly Glu Leu Arg Val
          85          90          95

```

Leu Ala Asp Thr Tyr Asp Ser Val Tyr Leu His Pro Asn Ser Tyr Ser
 100 105 110
 Cys Ala Cys Leu Ala Ser Gly Ser Val Leu Arg Leu Val Asp Ala Val
 115 120 125
 Leu Gly Ala Glu Ile Arg Asn Gly Met Ala Ile Ile Arg Pro Pro Gly
 130 135 140
 His His Ala Gln His Ser Leu Met Asp Gly Tyr Cys Met Phe Asn His
 145 150 155 160
 Val Ala Val Ala Ala Arg Tyr Ala Gln Gln Lys His Arg Ile Arg Arg
 165 170 175
 Val Leu Ile Val Asp Trp Asp Val His His Gly Gln Gly Thr Gln Phe
 180 185 190
 Thr Phe Asp Gln Asp Pro Ser Val Leu Tyr Phe Ser Ile His Arg Tyr
 195 200 205
 Glu Gln Gly Arg Phe Trp Pro His Leu Lys Ala Ser Asn Trp Ser Thr
 210 215 220
 Thr Gly Phe Gly Gln Gly Gln Gly Tyr Thr Ile Asn Val Pro Trp Asn
 225 230 235 240
 Gln Val Gly Met Arg Asp Ala Asp Tyr Ile Ala Ala Phe Leu His Val
 245 250 255
 Leu Leu Pro Val Ala Leu Glu Phe Gln Pro Gln Leu Val Leu Val Ala
 260 265 270
 Ala Gly Phe Asp Ala Leu Gln Gly Asp Pro Lys Gly Glu Met Ala Ala
 275 280 285
 Thr Pro Ala Gly Phe Ala Gln Leu Thr His Leu Leu Met Gly Leu Ala
 290 295 300
 Gly Gly Lys Leu Ile Leu Ser Leu Glu Gly Gly Tyr Asn Leu Arg Ala
 305 310 315 320
 Leu Ala Glu Gly Val Ser Ala Ser Leu His Thr Leu Leu Gly Asp Pro
 325 330 335
 Cys Pro Met Leu Glu Ser Pro Gly Ala Pro Cys Arg Ser Ala Gln Ala
 340 345 350
 Ser Val Ser Cys Ala Leu Glu Ala Leu Glu Pro Phe Trp Glu Val Leu
 355 360 365

Val Arg Ser Thr Glu Thr Val Glu Arg Asp Asn Met Glu Glu Asp Asn
 370 375 380
 Val Glu Glu Ser Glu Glu Glu Gly Pro Trp Glu Pro Pro Val Leu Pro
 385 390 395 400
 Ile Leu Thr Trp Pro Val Leu Gln Ser Arg Thr Gly Leu Val Tyr Asp
 405 410 415
 Gln Asn Met Met Asn His Cys Asn Leu Trp Asp Ser His His Pro Glu
 420 425 430
 Val Pro Gln Arg Ile Leu Arg Ile Met Cys Arg Leu Glu Glu Leu Gly
 435 440 445
 Leu Ala Gly Arg Cys Leu Thr Leu Thr Pro Arg Pro Ala Thr Glu Ala
 450 455 460
 Glu Leu Leu Thr Cys His Ser Ala Glu Tyr Val Gly His Leu Arg Ala
 465 470 475 480
 Thr Glu Lys Met Lys Thr Arg Glu Leu His Arg Glu Ser Ser Asn Phe
 485 490 495
 Asp Ser Ile Tyr Ile Cys Pro Ser Thr Phe Ala Cys Ala Gln Leu Ala
 500 505 510
 Thr Gly Ala Ala Cys Arg Leu Val Glu Ala Val Leu Ser Gly Glu Val
 515 520 525
 Leu Asn Gly Ala Ala Val Val Arg Pro Pro Gly His His Ala Glu Gln
 530 535 540
 Asp Ala Ala Cys Gly Phe Cys Phe Phe Asn Ser Val Ala Val Ala Ala
 545 550 555 560
 Arg His Ala Gln Thr Ile Ser Gly His Ala Leu Arg Ile Leu Ile Val
 565 570 575
 Asp Trp Asp Val His His Gly Asn Gly Thr Gln His Met Phe Glu Asp
 580 585 590
 Asp Pro Ser Val Leu Tyr Val Ser Leu His Arg Tyr Asp His Gly Thr
 595 600 605
 Phe Phe Pro Met Gly Asp Glu Gly Ala Ser Ser Gln Ile Gly Arg Ala
 610 615 620
 Ala Gly Thr Gly Phe Thr Val Asn Val Ala Trp Asn Gly Pro Arg Met
 625 630 635 640

Gly Asp Ala Asp Tyr Leu Ala Ala Trp His Arg Leu Val Leu Pro Ile
645 650 655

Ala Tyr Glu Phe Asn Pro Glu Leu Val Leu Val Ser Ala Gly Phe Asp
660 665 670

Ala Ala Arg Gly Asp Pro Leu Gly Gly Cys Gln Val Ser Pro Glu Gly
675 680 685

Tyr Ala His Leu Thr His Leu Leu Met Gly Leu Ala Ser Gly Arg Ile
690 695 700

Ile Leu Ile Leu Glu Gly Gly Tyr Asn Leu Thr Ser Ile Ser Glu Ser
705 710 715 720

Met Ala Ala Cys Thr Arg Ser Leu Leu Gly Asp Pro Pro Pro Leu Leu
725 730 735

Thr Leu Pro Arg Pro Pro Leu Ser Gly Ala Leu Ala Ser Ile Thr Glu
740 745 750

Thr Ile Gln Val His Arg Arg Tyr Trp Arg Ser Leu Arg Val Met Lys
755 760 765

Val Glu Asp Arg Glu Gly Pro Gly His His His His His His
770 775 780

<210> 6
<211> 2349
<212> DNA
<213> Custom

<220>
<221> DNA encoding residues 73-845 of HDAC6 and a 6-histidine tag at
the C-terminus
<222> (1)..(2349)

<400> 6	
atgcccgga tggatctgaa ccttgaggct gaagcactgg ctggcactgg cttggtgttg	60
gatgagcagt taaatgaatt ccattgcctc tgggatgaca gcttcccga aggccctgag	120
cggctccatg ccatcaagga gcaactgatc caggagggcc tcctagatcg ctgcgtgtcc	180
tttcaggccc ggtttgctga aaaggaagag ctgatgttgg ttcacagcct agaatatatt	240
gatctgatgg aaacaacca gtacatgaat gagggagaac tccgtgtcct agcagacacc	300
tacgactcag tttatctgca tccgaactca tactcctgtg cctgcctggc ctgaggctct	360
gtcctcaggc tggatgatgc ggtcctgggg gctgagatcc ggaatggcat ggccatcatt	420
aggcctctcg gacatcacgc ccagcacagt cttatggatg gctattgcat gttcaaccac	480
gtggctgtgg cagcccgcta tgctcaacag aaacaccgca tccggagggt cttatcgta	540

gattgggatg tgcaccacgg tcaaggaaca cagtccacct tcgaccagga cccagtgctc 600
 ctctatttct ccatccaccg ctacgagcag ggtaggttct ggccccacct gaaggcctct 660
 aactgggtcca ccacagggtt cgccaaggc caaggatata ccatcaatgt gccttggaac 720
 caggtgggga tgcgggatgc tgactacatt gctgctttcc tgcacgtcct gctgccagtc 780
 gccctcgagt tccagcctca gctggtcctg gtggctgctg gatttgatgc cctgcaaggg 840
 gacccaagg gtgagatggc cgccactccg gcagggttcg cccagctaac ccacctgctc 900
 atgggtctgg caggaggcaa gctgatacctg tctctggagg gtggctacaa cctccgcgcc 960
 ctggctgaag gcgtcagtgc ttcgctccac acccttctgg gagaccctg ccccatgctg 1020
 gagtcacctg gtgccccctg ccggagtgc caggcttcag tttcctgtgc tctggaagcc 1080
 cttgagccct tctgggaggt tcttgtaga tcaactgaga ccgtggagag ggacaacatg 1140
 gaggaggaca atgtagagga gagcgaggag gaaggacctt gggagcccc tgtgctccca 1200
 atcctgacat ggccagtgtc acagtctcgc acagggttg tctatgacca aaatatgatg 1260
 aatcactgca acttgtagga cagccaccac cctgaggtac cccagcgcac cttgcggatc 1320
 atgtgccgtc tggaggagct gggccttgcc gggcgctgcc tcacctgac accgcgccct 1380
 gccacagagg ctgagctgct cacctgtcac agtgctgagt acgtgggtca tctccgggcc 1440
 acagagaaaa tgaaaacccg ggagctgcac cgtgagagtt ccaactttga ctccatctat 1500
 atctgcccc gtaccttcgc ctgtgcacag cttgccactg gcgctgcctg ccgcctgggtg 1560
 gaggctgtgc tctcaggaga ggttctgaat ggtgctgctg tgggtgcgtcc cccaggacac 1620
 cacgcagagc aggatgcagc ttgcggtttt tgctttttca actctgtggc tgtggctgct 1680
 cgccatgccc agactatcag tgggcatgcc ctacggatcc tgattgtgga ttgggatgtc 1740
 caccacggta atggaactca gcacatgttt gaggatgacc ccagtgtgct atatgtgtcc 1800
 ctgcaccgct atgatcatgg caccttcttc cccatggggg atgaggggtg cagcagccag 1860
 atcgccgggg ctgcggggcac aggccttcacc gtcaacgtgg catggaacgg gccccgcatg 1920
 ggtgatgctg actacctagc tgcctggcat cgcctgggtg tttccattgc ctacgagttt 1980
 aaccagaac tgggtgctggt ctgagctggc ttgatgctg cacgggggga tccgctgggg 2040
 ggctgccagg tgtcacctga gggttatgcc cacctcacc acctgctgat gggccttgcc 2100
 agtggccgca ttatccttat cctagagggt ggctataacc tgacatccat ctgagagtc 2160
 atggctgcct gcactcgtc cctccttgga gaccaccac ccctgctgac cctgccacgg 2220
 cccccactat caggggccct ggcctcaatc actgagacca tccaagtcca tcgcagatac 2280
 tggcgagct tacgggtcat gaaggtagaa gacagagaag gacccgggca tcaccatcac 2340
 catcactaa 2349

<210> 7
 <211> 385
 <212> PRT

<213> Custom

<220>
 <221> Residues 1-377 of HDAC8 and a 6-histidine tag at the N-terminus
 <222> (1)..(385)

<400> 7

Met His His His His His His Pro Met Glu Glu Pro Glu Glu Pro Ala
 1 5 10 15

Asp Ser Gly Gln Ser Leu Val Pro Val Tyr Ile Tyr Ser Pro Glu Tyr
 20 25 30

Val Ser Met Cys Asp Ser Leu Ala Lys Ile Pro Lys Arg Ala Ser Met
 35 40 45

Val His Ser Leu Ile Glu Ala Tyr Ala Leu His Lys Gln Met Arg Ile
 50 55 60

Val Lys Pro Lys Val Ala Ser Met Glu Glu Met Ala Ala Phe His Thr
 65 70 75 80

Asp Ala Tyr Leu Gln His Leu Gln Lys Val Ser Gln Glu Gly Asp Asp
 85 90 95

Asp His Pro Asp Ser Ile Glu Tyr Gly Leu Gly Tyr Asp Cys Pro Ala
 100 105 110

Thr Glu Gly Ile Phe Asp Tyr Ala Ala Ala Ile Gly Gly Ala Thr Ile
 115 120 125

Thr Ala Ala Gln Cys Leu Ile Asp Gly Met Cys Lys Val Ala Ile Asn
 130 135 140

Trp Ser Gly Gly Trp His His Ala Lys Lys Asp Glu Ala Ser Gly Phe
 145 150 155 160

Cys Tyr Leu Asn Asp Ala Val Leu Gly Ile Leu Arg Leu Arg Arg Lys
 165 170 175

Phe Glu Arg Ile Leu Tyr Val Asp Leu Asp Leu His His Gly Asp Gly
 180 185 190

Val Glu Asp Ala Phe Ser Phe Thr Ser Lys Val Met Thr Val Ser Leu
 195 200 205

His Lys Phe Ser Pro Gly Phe Phe Pro Gly Thr Gly Asp Val Ser Asp
 210 215 220

Val Gly Leu Gly Lys Gly Arg Tyr Tyr Ser Val Asn Val Pro Ile Gln
 225 230 235 240

Asp Gly Ile Gln Asp Glu Lys Tyr Tyr Gln Ile Cys Glu Ser Val Leu
 245 250 255

Lys Glu Val Tyr Gln Ala Phe Asn Pro Lys Ala Val Val Leu Gln Leu
 260 265 270

Gly Ala Asp Thr Ile Ala Gly Asp Pro Met Cys Ser Phe Asn Met Thr
 275 280 285

Pro Val Gly Ile Gly Lys Cys Leu Lys Tyr Ile Leu Gln Trp Gln Leu
 290 295 300

Ala Thr Leu Ile Leu Gly Gly Gly Gly Tyr Asn Leu Ala Asn Thr Ala
 305 310 315 320

Arg Cys Trp Thr Tyr Leu Thr Gly Val Ile Leu Gly Lys Thr Leu Ser
 325 330 335

Ser Glu Ile Pro Asp His Glu Phe Phe Thr Ala Tyr Gly Pro Asp Tyr
 340 345 350

Val Leu Glu Ile Thr Pro Ser Cys Arg Pro Asp Arg Asn Glu Pro His
 355 360 365

Arg Ile Gln Gln Ile Leu Asn Tyr Ile Lys Gly Asn Leu Lys His Val
 370 375 380

Val
 385

<210> 8
 <211> 1158
 <212> DNA
 <213> Custom

<220>
 <221> DNA encoding residues 1-377 of HDAC8 and a 6-histidine tag at the
 N-terminus
 <222> (1)..(1158)

<400> 8
 atgcaccatc accatcacca tcccatggag gagccggagg aaccggcgga cagtgggcag 60
 tcgctggtcc cggtttatat ctatagtccc gagtatgtca gtatgtgtga ctccctggcc 120
 aagatcccca aacgggccag tatggtgcat tctttgattg aagcatatgc actgcataag 180
 cagatgagga tagttaagcc taaagtggcc tccatggagg agatggccgc cttccacact 240
 gatgcttatc tgcagcatct ccagaaggct agccaagagg gcatgatga tcatccggac 300
 tccatagaat atgggctagg ttatgactgc ccagccactg aagggatatt tgactatgca 360
 gcagctatag gaggggctac gatcacagct gcccaatgcc tgattgacgg aatgtgcaaa 420

gtagcaatta actggtcttg aggggtggcat catgcaaaga aagatgaagc atctggtttt	480
tggtatctca atgatgctgt cctgggaata ttacgattgc gacggaaatt tgagcgtatt	540
ctctacgtgg atttggatct gcaccatgga gatggtgtag aagacgcatt cagtttcacc	600
tccaaagtca tgaccgtgtc cctgcacaaa ttctccccag gatttttccc aggaacaggt	660
gacgtgtctg atgttggcct aggggaaggga cgggtactaca gtgtaaatgt gcccattcag	720
gatggcatac aagatgaaaa atattaccag atctgtgaaa gtgtactaaa ggaagtatac	780
caagccttta atcccaaagc agtgggtctta cagctgggag ctgacacaat agctggggat	840
cccatgtgct cctttaacat gactccagtg ggaattggca agtgtcttaa gtacatcctt	900
caatggcagt tggcaacact cattttggga ggaggaggct ataaccttgc caacacggct	960
cgatgctgga catacttgac cggggtcac ctagggaaaa cactatcctc tgagatccca	1020
gatcatgagt ttttcacagc atatggctct gattatgtgc tggaaatcac gccaagctgc	1080
cggccagacc gcaatgagcc ccaccgaatc caacaaatcc tcaactacat caaagggaat	1140
ctgaagcatg tgggtctag	1158